






A
PA−target enrichment by degree decile
Top two deciles; stratified−permutation p=1.0e−4
top−2 deciles
OR 3.47, p=3.8e−6
expected (all hubs)
% PA targets
Cross−species−degree decile (10 = highest)


B
Orthogonal InTxDB comparison
Shared hubs; Spearman rho=0.65, p=1.6e−01
CFH
TP53
PLG
RHOA
RAC1
CDC42
co−targeting species
IntAct
InTxDB


C
Leave−one−pathogen−out transfer
AUC range 0.738−0.755
transfer AUC−ROC
0.740
0.749
0.755
0.746
0.750
0.748
0.738
full
leave Yersinia
leave Bacillus
leave Francisella
leave Salmonella
leave Chlamydia
leave Escherichia


D
Hard−negative transfer by host degree
Spearman rho=−0.98, p=3.4e−5
transfer AUC−ROC
Negative−pool median host degree


E
Negative−set resampling
mean 0.701; range 0.63−0.77; full model 0.740
transfer AUC−ROC
negative−set resampling (seed)


F
Human−lung three−pathogen response
n=5 donors; FC=2^log2FC; * nominal paired p<0.05
COL1A1
TF
CAMP
FN1
ELANE
MUC1
NLRC4
TLR5
Fold change vs uninfected
Ctrl
PA
IAV
BCG
Uninfected
PA
IAV
BCG