






A
FliC exposure in mouse MDSC (bracket: WT vs dFliC; * p<0.05)
Myd88 M−MDSC
Myd88 PMN−MDSC
Tlr5 M−MDSC
Tlr5 PMN−MDSC
mean counts
NT
WT
dFliC
*


B
PA remodels BALF composition (scRNA)
sham: AlvMac 87.5% | cDC 10.1%
PA: Neu 89.6% | AlvMac 2.6% | cDC 2.2%
% BALF cells (WT)
sham
PA
AlvMac
cDC
Mono
Neu
T


C
Lower TLR5 within AlvMac and cDC
Descriptive pooled libraries; n=1 per condition
sham
PA
TLR5 mean log−norm expr
0.082
0.016
0.088
0.003
AlvMac
cDC


D
Pooled BALF libraries
One library per condition; descriptive only
WT sham
WT PA
DKO sham
DKO PA
Tlr5-detected cells (%)
AlvMac
cDC


E
MYD88 dynamics are cell−type−dependent (not co−suppressed)
Mouse lung (SP−AKI)
Mouse lung (meta−DEG)
MDSC M−
Human lung explants
MDSC PMN−
MYD88 log2FC (PA vs control)


F
WT−vs−dFliC equivalence
Gray lines: B10.5 and B11.0 log2FC margins; TLR5 equiv
Tlr5 M−MDSC
Tlr5 PMN−MDSC
Myd88 M−MDSC
Myd88 PMN−MDSC
WT vs dFliC log2FC contrast (limma−voom; df=12)
WT vs dFliC log2FC contrast (limma−voom)