





A TLR5 meta−analysis, nine datasets
Pooled −0.86 (95% CI −1.50 to −0.22); I2=96.1%; PI −2.81 to 1.09
Mouse lung, GSE233206
Mouse lung, GSE192890 (acute)
Mouse lung, SP−AKI
Human lung explant
Human cornea
Mouse MDSC
Human airway organoids
Primary HBE (GSE97036)
Persistent Calu−3, 120 h
Pooled (REML)
Prediction interval
−2.41 (−2.85, −1.97)
−1.15 (−1.56, −0.74)
−0.90 (−1.22, −0.59)
−0.55 (−1.18, 0.08)
−2.08 (−3.25, −0.92)
−0.39 (−1.41, 0.62)
−0.02 (−0.19, 0.16)
−1.22 (−1.51, −0.94)
0.73 (0.36, 1.10)
−0.86 (−1.50, −0.22)
−0.86 (−2.81, 1.09)
−2
0
2
TLR5 log2FC (PA vs control)


B Fixed receptor/adaptor panel
TLR5 is the only FDR−significant negative effect
NLRP3
TLR2
NOD2
MYD88
TLR6
TIRAP
TLR4
TLR9
NLRC4
IRAK1
IRAK4
NOD1
TLR5
−4
−2
0
2
4
6
Pooled log2FC (REML)


C Leave−one−dataset−out
Mouse lung, GSE233206
Mouse lung, GSE192890 (acute)
Mouse lung, SP−AKI
Human lung explant
Human cornea
Mouse MDSC
Human airway organoids
Primary HBE (GSE97036)
Persistent Calu−3, 120 h
−2.0
−1.5
−1.0
−0.5
0.0
Pooled TLR5 log2FC with dataset excluded


D Sensitivity analyses
No Egger inference with fewer than 10 studies
Primary (9 datasets)
Hartung−Knapp
Exclude persistent
Exclude GSE97036
Persistent endpoint 24 h
MDSC effects split
Exclude reanalysed SRPs
−2
−1
0
Pooled TLR5 log2FC


E Receptor/adaptor panel (added dataset)
GSE201465, donor−paired DESeq2; TLR2, TLR4 and MYD88 induced
log2FC (PA vs control)
1.5
1.0
0.5
0.0
−0.5
ns
***
**
TLR6
NOD1
NOD2
TLR5
NLRC4
IRAK1
IRAK4
TIRAP
MYD88
TLR2
TLR4