

A
FliC exposure in mouse MDSC (bracket: WT vs dFliC; * p<0.05)
Myd88 M−MDSC
Myd88 PMN−MDSC
Tlr5 M−MDSC
Tlr5 PMN−MDSC
mean counts
NT
WT
dFliC
B
PA remodels BALF composition (scRNA)
sham: AlvMac 87.5% | cDC 10.1%
PA: Neu 89.6% | AlvMac 2.6% | cDC 2.2%
% BALF cells (WT)
sham
PA
AlvMac
cDC
Mono
Neu
T
B
Epi
NK
Other
pDC
C
Lower TLR5 within AlvMac and cDC
Descriptive pooled libraries; n=1 per condition
sham
PA
TLR5 mean log−norm expr
0.082
0.016
0.088
0.003
AlvMac
cDC
D
UMI−adjusted Tlr5 detection
PA vs sham; descriptive pooled−library model
Odds ratio for Tlr5 detection
OR = 0.33 (95% CI 0.14−0.74), p=0.00702
PA infection
E
MYD88 dynamics are cell−type−dependent (not co−suppressed)
Mouse lung (SP−AKI)
Mouse lung (meta−DEG)
MDSC M−
Human lung explants
MDSC PMN−
MYD88 log2FC (PA vs control)
F
WT−vs−dFliC equivalence analysis
TLR5 equivalence not established; TOST bound +/−0.5
equivalence bound +/−0.5
Tlr5 M−MDSC
−0.14 [−2.7, 2.4]; pTOST=0.38
Tlr5 PMN−MDSC
3.22 [1.2, 5.3]; pTOST=0.99
Myd88 M−MDSC
0.07 [−0.3, 0.5]; pTOST=0.017
Myd88 PMN−MDSC
1.66 [1.2, 2.1]; pTOST=1
WT vs dFliC log2FC contrast (limma−voom)