

On the Equilibrium of White Dwarfs

Yiqi Xie*

School of Physics, Peking University

July 6, 2017

Abstract

A model for non-rotating spherical helium white dwarfs was built and its equilibrium was solved. Electron degenerate pressure and Newtonian gravitational field were taken into consideration, based on which a key equation on the distribution of electron chemical potential was derived. We first checked the equation at the non-relativistic limit and the ultra-relativistic limit, then a general solution was obtained through numerical methods determined with adequate deliberation. We also calculated the density distribution and the total mass for a spectrum of white dwarfs, from which the contribution of relativistic effect to the complete collapse of the star was illustrated. The upper mass limit for a white dwarf to stand such collapse, namely the Chandrasekhar limit, has been derived as $1.44M_{\odot}$ in this paper, which is consistent with the number formerly provided by Chandrasekhar himself.

1 Introduction

White dwarfs, in observation, are abnormally faint for their color^[1]. They are the core remnants, the final evolutionary stage, of stars whose mass is at a low or medium level^[2]. A typical white dwarf is composed mostly of helium^[3], the product of hydrogen fusion reaction happened in its former evolutionary stages, and for some cases, heavier elements such as carbon and oxygen might also be found. Unable to fuse these elements by itself, a white dwarf cannot generate enough thermal pressure in against of gravity, and as a result, it collapses into a highly compressed state. The mass of a typical white dwarf is comparable to that of Sun, while its volume is comparable to that of Earth^{[4][5]}. Under such high density, the electron degeneracy pressure arises in place of thermal pressure^[6], so that the star stabilizes at a new equilibrium. Here we list some characteristic physical properties of a white dwarf in contrast with the Sun in Table 1.1.

Table 1.1: White Dwarf Sketch^{[4][5]}

Parameter	White Dwarf	Sun (in Contrast)
Mass M / g	$\sim 10^{33}$	$\sim 10^{33}$
Radius R / cm	$\sim 10^9$	$\sim 10^{11}$
Density ρ / $\text{g} \cdot \text{cm}^{-3}$	$\sim 10^6$	$\sim 10^0$
Temperature T / K	$\sim 10^7$	$\sim 10^7$

As mentioned above, the equilibrium between degenerate pressure and gravitational force holds the white dwarf in its shape. From such equilibrium it is possible to derive the constraints for a white dwarf's mass, radius, etc. Exact work has been done by Chandrasekhar in 1931-1935^{[7][8]}, where he stressed that the influence of relativistic effect in leading a white dwarf to collapse completely as its mass reaching to a limit - later been named after him - of $1.44M_{\odot}$. In following sections of this paper we will reformulate this work from our own perspective and see if we can derive Chandrasekhar's exact result.

*Student ID: 1300011383. Email: xieyiqi@pku.edu.cn

2 Modeling a White Dwarf

2.1 Thermodynamics for Degenerate Electrons

For the beginning let us discuss a little further on the degenerate electrons of white dwarf matters. To make it simple, we neglect the electron-electron interaction and the electron-nuclei interaction, so that the system of electrons could be treated as ideal gas - it is quite a proper model since we are dealing with matter at $\sim 10^7\text{K}$ where everything gets highly ionized.

According to quantum field theory, electron is a fermion that obeys the Pauli exclusion principle. When highly compressed in space, electrons, by intuition, have to distinguish each other in their momentum, thus even at the temperature of absolute zero, most electrons are still fastly moving around, resulting in a non-zero pressure, namely the degeneracy pressure.

Degenerate electron gas at finite temperature experiences both ordinary thermal pressure and degeneracy pressure. However, when the temperature is not too high, degeneracy pressure still steadily dominates^[3]. In a more specific way, the above condition for temperature writes

$$k_B T \ll \epsilon_F, \quad (2.1)$$

where ϵ_F , or Fermi energy, represents the chemical potential of the electron gas at $T = 0$. The value of ϵ_F can be calculated from the number density n of the gas by

$$\epsilon_F = \sqrt{p_F^2 c^2 + m_e^2 c^4} - m_e c^2, \quad p_F = (3\pi^2 n)^{1/3} \hbar. \quad (2.2)$$

Note that for safety we have applied relativistic dispersion relation in (2.2), where p_F is the momentum corresponding to ϵ_F . Since we are dealing with white dwarfs made of helium, that every electron is bonded with one proton plus one neutron, we may substitute number density n by mass density ρ through

$$n \approx \frac{\rho}{2m_p}. \quad (2.3)$$

Recall the information in Table 1.1, we take $\rho \sim 10^6 \text{g} \cdot \text{cm}^{-3}$, and by (2.2) and (2.3) the calculation will give $\epsilon_F \sim 10^5 \text{eV}$. On the other hand we see $k_B T \sim 10^3 \text{eV}$, so that the condition in (2.1) is well satisfied. Therefore, when dealing with the thermodynamics of degenerate electron gas of a white dwarf, one may take $T = 0$ and consider degeneracy pressure as the only mechanism against gravity. In addition, it is important to note that ϵ_F is comparable to the rest energy of an electron, so we shall conclude that the application of relativistic dispersion relation in (2.2) is necessary.

Till now we haven't inspected any specific behavior of degeneracy pressure yet - actually we do not need to know many details except for its relation to density or to Fermi energy. Recall Gibbs-Duhem equation:

$$Nd\mu = -SdT + VdP. \quad (2.4)$$

With T been considered zero in our case, the dT term get crossed out, reducing the equation to

$$nd\mu = dP, \quad (2.5)$$

where for now μ is the same of ϵ_F and P stands totally for the degeneracy pressure. We will see that (2.5) is good enough for later discussion.

2.2 Equation of Equilibrium

We consider only non-rotating white dwarfs, where centrifugal force does not exist. Non-rotating white dwarfs basically live on an equilibrium merely between the electron degeneracy pressure and the gravitational force. Witnessed the special relativistic nature of the densed electron gas, one may also be cautious on the general relativistic effect of the gravity. Luckily such effect is negletable, for the escape velocity of a typical white dwarf - roughly estimated as the order of $\sqrt{GM/R} \sim 10^8 \text{cm} \cdot \text{s}^{-1}$ - is still much lower than the speed of light^[9]. Thus we consider that the form of Newtonian gravitational field is fairly acceptable, and the equation of equilibrium writes

$$\nabla P + \rho \nabla \phi = 0, \quad \nabla^2 \phi = 4\pi G \rho. \quad (2.6)$$

In former section we have clarified the relations between μ , P and ρ . Putting (2.2), (2.3) and (2.5) together with (2.6), reducing for several steps, and taking spherical symmetry, we can now express the equation of equilibrium in the spatial distribution of μ :

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\mu}{dr} \right) + \frac{16Gm_p^2}{3\pi\hbar^3 c^3} \cdot \mu^{3/2} (\mu + 2m_e c^2)^{3/2} = 0. \quad (2.7)$$

where r represents the radial distance.

We consider (2.7) as the key equation to solve in this paper. Categorically it is a second order differential equation wanting two boundary conditions to define its solution, one of which should apparently be that density decreases to zero on the surface, and the other of which comes from the requirement of symmetry that the gradient of any distributed physical parameter should converge to zero at the center. So we add

$$\mu \Big|_{r=R} = 0, \quad \frac{d\mu}{dr} \Big|_{r=0} = 0. \quad (2.8)$$

Once we solved $\mu(r)$, the distribution of density can be easily derived through

$$\rho = \frac{2m_p}{3\pi^2\hbar^3 c^3} \cdot \mu^{3/2} (\mu + 2m_e c^2)^{3/2}, \quad (2.9)$$

as a result from (2.2) and (2.3), and by integral a constraint on M and R arises as

$$M = \int_0^R 4\pi\rho r^2 dr = -\frac{R^2}{2Gm_p} \frac{d\mu}{dr} \Big|_{r=R}. \quad (2.10)$$

Now we have completed a model for non-rotating spherical helium white dwarfs and derived the equation of equilibrium as specific as possible. For the following sections we will move on to the solution of (2.7)-(2.10). Here we construct three constants as below that will later be frequently used:

$$\begin{cases} R_0 = \left(\frac{3\pi\hbar^3}{64Gm_p^2 m_e^2 c} \right)^{1/2} \approx 1.93 \times 10^8 \text{cm} \\ M_0 = \left(\frac{3\pi\hbar^3 c^3}{64G^3 m_p^4} \right)^{1/2} \approx 1.42 \times 10^{33} \text{g} \\ \rho_0 = \frac{16m_p m_e^3 c^3}{\pi^2 \hbar^3} \approx 4.71 \times 10^7 \text{g} \cdot \text{cm}^{-3} \end{cases} \quad (2.11)$$

We may note that all these constans are characteristic for a typical white dwarf.

3 Solution to the Equation of State

3.1 Preliminary Analysis

For tentative treatment, we take noramlization

$$\xi = \frac{r}{R}, \quad f = \frac{\mu}{2m_e c^2}. \quad (3.1)$$

As a result, (2.7) with (2.8) turns into

$$\begin{cases} \frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{df}{d\xi} \right) + \left(\frac{R}{R_0} \right)^2 f^{3/2} (f+1)^{3/2} = 0 \\ f \Big|_{\xi=1} = 0, \quad \frac{df}{d\xi} \Big|_{\xi=0} = 0 \end{cases} \quad (3.2)$$

From the first sight we realize that this is a indefinite equation with one variable parameter R . Since the only restriction on R is positivity, our equation is doomed to suffer from numerical deformation when R varies over the full range of $(0, +\infty)$. Such deformation would cause difficulties in solution, and for some cases we have to adjust our normalization on μ . More specifically:

- For $R/R_0 \sim 1$: this is the only good case for (3.2), where $f \sim 1$ and the relativistic nature of the electron gas remains authentic;
- For $R/R_0 \gg 1$: this is where f has to be very small in order to balance the equation, and the electron gas reaches the non-relativistic end. Better choice of normalization on μ might be $f_n = (R/R_0)^4 \cdot f$, so that (3.2) becomes

$$\begin{cases} \frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{df_n}{d\xi} \right) + f_n^{3/2} \left[\left(\frac{R_0}{R} \right)^4 f_n + 1 \right]^{3/2} = 0 \\ f_n \Big|_{\xi=1} = 0, \quad \frac{df_n}{d\xi} \Big|_{\xi=0} = 0 \end{cases}; \quad (3.3)$$

- For $R/R_0 \ll 1$: this is where f has to be very large in order to balance the equation, and the electron gas reaches the ultra-relativistic end. Better choice of normalization on μ might be $f_u = R/R_0 \cdot f$, so that (3.2) becomes

$$\begin{cases} \frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{df_u}{d\xi} \right) + f_u^{3/2} \left(f_u + \frac{R}{R_0} \right)^{3/2} = 0 \\ f_u \Big|_{\xi=1} = 0, \quad \frac{df_u}{d\xi} \Big|_{\xi=0} = 0 \end{cases}. \quad (3.4)$$

Though our solution might be intricate in its relation to R , we may first let $R \rightarrow +\infty$ and $R \rightarrow 0$ and solve the cases of non-relativistic limit and of ultra-relativistic limit respectively. We hope that results at these two limits could set up the boundaries for the whole problem and give us some clue to the solution of the general case.

3.2 Non-Relativistic Limit and Ultra-Relativistic Limit

Taking $R/R_0 \rightarrow +\infty$ in (3.3) and $R/R_0 \rightarrow 0$ in (3.4), we obtain their asymptotic forms as below

$$\begin{cases} \frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{df_n}{d\xi} \right) + f_n^{3/2} = 0 \\ f_n \Big|_{\xi=1} = 0, \quad \frac{df_n}{d\xi} \Big|_{\xi=0} = 0 \end{cases}, \quad \text{and} \quad \begin{cases} \frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{df_u}{d\xi} \right) + f_u^3 = 0 \\ f_u \Big|_{\xi=1} = 0, \quad \frac{df_u}{d\xi} \Big|_{\xi=0} = 0 \end{cases}. \quad (3.5)$$

In mathematics they are known as the Lane-Emden equations indexed 3/2 and 3 respectively. Density distribution and total mass are related to the solutions in

$$\begin{cases} \rho_n = \frac{1}{3} \rho_0 \left(\frac{R_0}{R} \right)^6 \cdot f_n^{3/2} \\ M_n = -M_0 \left(\frac{R_0}{R} \right)^3 \cdot \frac{df_n}{d\xi} \Big|_{\xi=1} \end{cases}, \quad \text{and} \quad \begin{cases} \rho_u = \frac{1}{3} \rho_0 \left(\frac{R_0}{R} \right)^3 \cdot f_u^3 \\ M_u = -M_0 \cdot \frac{df_u}{d\xi} \Big|_{\xi=1} \end{cases}. \quad (3.6)$$

Analytical solutions to (3.5) do not exist. One may solve them through numerical approaches (we used MATLAB). Here we only complete their boundary values¹ in Table 3.1, based on which we state:

Table 3.1: Solution Feature at the Limits

	f_n	f_u
$f(0)$	178.22	6.90
$f'(1)$	-132.38	-2.02

¹More details could be found in Section 3.3 where the general solution over a spectrum of R is illustrated.

- The ratio of central density to average density is

$$\rho_c/\bar{\rho} = \begin{cases} 5.99 & \text{non-relativistic} \\ 54.18 & \text{ultra-relativistic} \end{cases}, \quad (3.7)$$

indicating that the emergence of relativistic effects will put a significant improvement on the matter concentration at the star's center;

- While we see in (3.6) that for the non-relativistic limit M tends to zero at a pace of R^{-3} , it turns out that for the ultra-relativistic limit M is fixed at

$$2.02M_0 = 1.44M_\odot, \quad (3.8)$$

which is quite fascinating and calls for further inspection.

3.3 Solution to the General Case

Let's move on to solve (2.7)-(2.10) with R/R_0 varying over $(0, +\infty)^2$. One may divide that range into three intervals and apply (3.2)-(3.4) respectively to gain a complete solution. Nevertheless, here we better introduce another normalization on μ , first proposed by Chandrasekhar^[7]:

$$\theta = \frac{\gamma}{\gamma_c}, \quad (3.9)$$

where γ is the Lorentz factor, and γ_c is the value of γ at the center of the star, *i.e.*,

$$\gamma = 1 + \frac{\mu}{m_e c^2}, \quad \gamma_c = \gamma \Big|_{\xi=0}. \quad (3.10)$$

Then (2.7) with (2.8) becomes

$$\begin{cases} \frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d\theta}{d\xi} \right) + \left(\frac{\gamma_c R}{2R_0} \right)^2 \left(\theta^2 - \frac{1}{\gamma_c^2} \right)^{3/2} = 0 \\ \theta \Big|_{\xi=1} = \frac{1}{\gamma_c}, \quad \theta \Big|_{\xi=0} = 1, \quad \frac{d\theta}{d\xi} \Big|_{\xi=0} = 0 \end{cases}, \quad (3.11)$$

and we rewrite the density distribution and the total mass in

$$\begin{cases} \rho = \frac{1}{24} \rho_0 \cdot (\gamma_c^2 \theta^2 - 1)^{3/2} \\ M = -M_0 \left(\frac{\gamma_c R}{2R_0} \right) \cdot \frac{d\theta}{d\xi} \Big|_{\xi=1} \end{cases}. \quad (3.12)$$

Note that now our equation contains one more variable parameter γ_c , together with one more boundary condition $\theta|_{\xi=0} = 1$, we should expect that certain constraint between γ_c and R exists and can be solved spontaneously with $\theta(\xi)$.

The reason to consider Chandrasekhar's normalization consists in its numerical stability:

- θ is constantly restricted within $(0, 1)$;
- when $R/R_0 \rightarrow 0$, factor $\gamma_c R/2R_0$ converges to a finite number, namely $f_u(0) = 6.90$;
- when $R/R_0 \sim 1$, factor $1/\gamma_c^2$ is sufficiently small so that it would not eat up the θ^2 aside;
- when $R/R_0 \rightarrow +\infty$, numerical deformation arises as $\gamma_c R/2R_0$ explodes and $1/\gamma_c \rightarrow 1$, but luckily we are not particularly interested in such case.

It is proper to state that solution $\theta(\xi)$ would not experience much change as we adjust R/R_0 successively within the range of our interest, which is a favourable pattern for solving indefinite equation with variable parameters. Again we use MATLAB to solve these equations. The work is done by repeatedly calling "bvp4c" function, where at the k -th iteration we take the solution to the $(k-1)$ -th equation as

²To some extent we are not much interested in the solution to $R \rightarrow \infty$, for it is not a white dwarf anymore. Generally speaking $R \sim 10R_0$ should be helpful enough as the upper limit.

the initializer. To start with, one may make use of the ultra-relativistic solution $f_u(\xi)$ to initialize the first equation. In other words, it takes

$$\theta^{(0)} = \frac{1 + 2f_u}{1 + 2f_u(0)}, \quad \gamma_c^{(0)} = f_u(0), \quad R^{(0)} = 0, \quad (3.13)$$

and the whole iteration works as $R^{(k)}$ gradually increases each time.

Final results are plotted in Figure 3.1, where a spectrum of white dwarfs has been featured. We see that when a white dwarf gains mass, its radius decreases, its electrons render more relativistic features, and its matter redistributes under new equilibrium in a way that more proportion of mass concentrates to the center. The rising of relativistic effect also boosts the drop of R as M approaches $2.02M_0 = 1.44M_\odot$. Reaching such mass limit, a white dwarf becomes incredibly unstable as it collapses into a volumeless point at a pace of $dR/dM \rightarrow \infty$.

So far our result has been consistent with Chandrasekhar's work, and for its physical meaning $1.44M_\odot$ is finally verified as the Chandrasekhar limit.




Figure 3.1: The general solution and the derived results. In (b)(c)(d) former results at the non-relativistic limit and the ultra-relativistic limit are also added.

4 Conclusion

In this paper, we have discussed the equilibrium properties of non-rotating spherical helium white dwarfs. Our key equation of equilibrium was derived on the distribution of electron chemical potential, and through the paper we solved it step by step. We noticed that the radius R is variable in the equation, and different normalization should be applied to solve different cases. Solutions to the non-relativistic limit (for $R \rightarrow \infty$) and the ultra-relativistic limit (for $R \rightarrow 0$) were obtained at first as they are relatively simple in treatment, and the exact value of Chandrasekhar limit could be obtained at this moment though its physical meaning was not fully revealed. For the general case, certain numerical techniques are required to solve the equation, and we found that the normalization proposed by Chandrasekhar was very helpful for its stability within the range of our interest.

The relativistic nature of the electrons is proved to be crucial. It arised from the beginning as we tried to estimate the electron degeneracy pressure of a typical white dwarf. Later inspection shows that the more compressed a white dwarf is, the more relativistic its electrons could be, and it reacts on the density distribution to concentrate more mass around the center. Back to the most concerned mass-radius relation, our result shows that the relativistic effect casts a significant boost on the drop of radius as mass increases, and the Chandrasekhar limit is not only the upper mass limit but also an unstable point where the star collapses drastically at $dR/dM \rightarrow \infty$. All these results are consistent with Chandrasekhar's original work.

Further discussion may focus on the general relativistic correction to our model. Though we neglected the general relativistic effect estimating the escape velocity of a typical white dwarf, we cannot say for sure what would happen to those around the Chandrasekhar limit, extremely densed within an almost volumeless region. For such case, the Tolman-Oppenheimer-Volkoff (TOV) equation^[10] should be applied to the model. Other factors like rapid rotation, heavier elements, magnetic fields, etc, may also be taken into consideration.

5 Acknowledgement

This work is reformed from our course project of Mathematical Physics II of 2017 spring, instructed by Prof. Chunyuan Gao, Peking University School of Physics. Other contributor to the original project includes Chenkai Mao, Ruixiao Yao, Youran Sun, and Zizhao Wang. We gratefully thank them for the foundation and improvement of this work.

References

- [1] E. L. Schatzman. *White Dwarfs*. Amsterdam: North-Holland Pub. Co., 1958.
- [2] M. Richmond. Late stages of evolution for low-mass stars. *Lecture notes, Physics 230*, 2007.
- [3] R. K. Pathria and Paul D. Beale. *Statistical Mechanics*. Elsevier Ltd., 3rd edition, 2012.
- [4] D. R. Williams. Sun fact sheet. <https://nssdc.gsfc.nasa.gov/planetary/factsheet/sunfact.html>, NASA Goddard Space Flight Center, 2013.
- [5] J. Johnson. Extreme stars: White dwarfs & neutron stars. *Lecture Notes, Astronomy*, 162, 2007.
- [6] R. H. Fowler. On dense matter. *Monthly Notices of the Royal Astronomical Society*, 87:114–122, 1926.
- [7] S. Chandrasekhar. The highly collapsed configurations of a stellar mass (second paper). *Monthly Notices of the Royal Astronomical Society*, 95:207–225, 1935.
- [8] S. Chandrasekhar. The highly collapsed configurations of a stellar mass. *Monthly Notices of the Royal Astronomical Society*, 91:456–466, 1931.
- [9] B. Schultz. *A First Course in General Relativity*. Cambridge University Press, 2nd edition, 2009.
- [10] J. R. Oppenheimer and G. M. Volkoff. On massive neutron cores. *Physical Review*, 55:374–381, 1939.