

Introduction to EXP2-1



EXP2-1 Same or Different?

Semester B, Weeks 1-2

The **aim** of EXP2-1 is to develop tools and skills, in order to:

- Present and report data in the correct way
(units, significant figures, error, and others)
- Determine whether the data are statistically significant


Introduction to EXP2-1 Part 1



Part 1 Random variations

When we measure a value repeatedly, we do not get exactly the same result each time

Typically, we see random variation about some value



Repeat the measurement to reveal random variation

Random variation



Some definitions

Variables may be

Discrete or categorical: only have specific values

e.g. Women or Men

number of legs

polymer students or materials students

Continuous or numerical: can have any value

e.g. most things we measure: length,

temperature, mass, strength, stiffness etc.

Random variation



Some definitions


Data: individual measured values (observations) (e.g. Student's X height)

Sample: a number of similar measured values (e.g. QMES students' height)

Population: hypothetical set of all results (e.g. Chinese students' height)

A population of random variation follows the **“Normal Distribution”**.

The Normal Distribution




Expresses the probability of a value lying within a certain distance of the mean


The Normal Distribution




Properties of the Normal Distribution



Mean and dispersion




Same dispersion different mean



Same mean different dispersion

Measures of central tendency



- Mean (average)
- Mode
- Median

Measures of central tendency



Mean

- For the population: $\mu = \frac{\sum X_i}{N}$
- For the sample: $\bar{X} = \frac{\sum X_i}{n}$

Mode

- Most commonly occurring value in a set of numbers
- Usually used when data grouped together in classes
 - is then sensitive to class interval.
- Can have 1 value (unimodal) or >1 values (e.g. bimodal, trimodal)

Measures of central tendency



Median

- Defined by position $(n+1)/2$ along dataset

Example 1: 1 5 6 8 9

Example 2: 200 200 230 250 280 300 350 400 480

Example 3: 200 200 230 250 280 300 350 400 480 500

Median = $(280 + 300) / 2 = \underline{290}$

Characterising variation



Characterising location

Central tendency:

Sample

Population

Mean or average value

$$\bar{X} = \frac{\sum X}{n}$$

μ

n = number of events (or number of measurements)

Characterising variation



Characterising spread

Dispersion:

Sample

Population

Range

$$X_{\max} - X_{\min}$$

$+\infty$ to $-\infty$

Standard deviation,
(SD or St Dev)

$$S = \sqrt{\frac{\sum (X - \bar{X})^2}{n - 1}}$$

$$\sigma = \sqrt{\frac{\sum (X - \bar{X})^2}{N}}$$

Characterising variation



Standard deviation, SD

Within

1 SD = 68.26% of population


2 SD = 95.46% of population

3 SD = 99.73% of population

...

Out of 6 SD

$\approx 1/1\text{million}$ of population



Expresses the probability of a value lying within a certain distance of the mean.


Characterising variation



Confidence intervals, CI

Probability of $x\%$ of values occurring within certain limit of the mean

i.e. How confident are we that a value will be found in this range?



Example:

95% confidence interval = ± 1.96 SD of mean

99% confidence interval = ± 2.58 SD of mean

Characterising variation




Coefficient of variation, CV

$$C.V. = \frac{St\ Dev}{Mean}$$

Gives a measure of the width or sharpness of the distribution.

e.g., $CV(A) < CV(B)$



Usually expressed as a percentage $CV\% = \frac{S}{\bar{X}} * 100$

Reporting data




In science – data do not have a single value,
we also need to show *how well we know the value*

Location $\pm$ **spread** (n = number of measurements)


Mean $\pm$ St Dev, n	}	Use 1 of these
Mean $\pm$ $x\%$ confidence intervals, n		
Mean + CV, n		

Always state what it is that you have quoted
There are other ways to express variation


Other forms of distribution



Bimodal




Unimodal



Small Variability



Large Variability



Positively Skewed



Negatively Skewed

Other forms of distribution




Central tendency:

Mean = average value

Median = middle value

Mode = most frequently occurring value



Dispersion:

Histograms – divide the data into categories or ‘bins’

Determine the number of bins:

use $\sqrt{n}$ bins or *Sturges rule* $1 + 3.3 \log n$ bins


(where n = the total number of observation in the dataset)

Sampling distribution



- **Population** - the entire group
- **Sample** - the specific group
- Probability samples do not guarantee '**correct**' answers in terms of their estimate of the underlying **population** value
- **Each sample varies** and so each sample's estimate will be different (hopefully not by much)
- Still, one sample is just one of many **possible samples** from the population

Sample means



Samples: 1, 2, 3, ...

Sample means: $\bar{X}_1, \bar{X}_2, \bar{X}_3 \dots$


- **Sample means** are random variations
- Sample means centered on population mean, with enough samples, average of all sample means = true population mean

Standard Error of the Mean



Sampling distribution is the plot of sample means.

Standard error of mean = standard deviation of the sampling distribution



Standard error (SE) of mean

becomes important:

If we know what SE is then we can assess how **likely** it is that a given **sample** is **representative** of the '**true**' population value.

Standard Error of the Mean



If our sample is relatively large ($n > 30$),

then the **Central Limit Theorem** holds true.

This tells us that the sampling distribution will have:

- **Normal** distribution
- Sample mean equals the population mean $\bar{X} = \mu$
- Standard deviation of sampling distribution $SE = \frac{\sigma}{\sqrt{n}}$
- Known properties (e.g. **95%** of the sample means will lie within 1.96 standard errors of the population mean), 95% C.I. = $\pm 1.96SE$
- If $n > 30$, then this holds even if the target population is **not** normally distributed!

Standard Error of the Mean



So?

So we can use the equation $\frac{\sigma}{\sqrt{n}}$ to approximate the **standard error** of the sampling distribution (SE = standard deviation of the sampling distribution of the sample mean)

If we know SE then we can assess how **likely** it is that a given **sample** is **representative** of the '**true**' population value.

Standard Error of the Mean



Standard error (SE) of a sample estimate:

We can use **Central Limit Theorem** to estimate the **accuracy** of the sample estimate of the mean

SE (standard deviation of sampling distribution): $\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$

SE indicated the likely **spread** of the **sample means** around the **population mean**

- a measure of sampling error and an indication of how **likely** it is that our sample estimate is close to the **true** underlying **population** value.

And when $n > 30$ we can use the equation above to estimate it.

Standard Error of the Mean



Standard error of the sampling distribution:

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$

$$SD \text{ of population } \sigma = \sqrt{\frac{\sum(X - \bar{X})^2}{N}}$$

$$\sigma_{\bar{X}} = \frac{S}{\sqrt{n}}$$

$$SD \text{ of sample } S = \sqrt{\frac{\sum(X - \bar{X})^2}{n - 1}}$$

Implications:

- The **larger the sample size** (n), the **smaller the error** in the sample is likely to be.
- The **larger the standard deviation** in the sample (S), the **larger the error** in the sample is likely to be.

Standard Error of the Mean & C.I.



Standard error of estimate allows us to calculate a **confidence interval** around our sample estimate.

Confidence interval = a **range** within which we are reasonably sure the **real population value lies**.

What does the confidence interval mean?

For a 95% confidence interval, think that if you collected 100 samples then the true population mean would lie within the confidence intervals of 95 of those samples.

SE & Confidence Intervals



When $n > 30$ the CLT holds and confidence intervals for sample means:


$$\text{Confidence interval} = \bar{X} \pm Z \cdot \sigma_{\bar{X}}$$

sample estimate
of the mean

Standard error
(standard deviation of sample /
square root of n)

Z is a z-value taken from the normal curve.
Value of Z depends on how much error
we are willing to accept.

Confidence Intervals



Confidence interval	90%	95%	99%
z	1.65	1.96	2.58

Summary of random distribution



- **Random variation** follows the **normal distribution**.
- **St Dev** expresses the dispersion about the mean value.
- **SE** (Standard Error of the Mean) expresses how far a sample mean is from the “true” value (the population mean).
- **Confidence intervals** express the probability of any value being within a certain distance of the mean.
- **Coefficient of variation** expresses the width of the distribution.
- **Histograms** can be useful in showing distributed values.

Tasks



- Check your group assignments on QM+
- Review the 'Introduction' and follow the 'Instructions'
- Download the raw data files from QM+
- Follow 7-step plan, hold your GMM to solve the problem
- Bring your questions to be discussed in class

EXP2-1 Schedule – M Groups



Date:	Time:	Place:	Activity:
Monday 20 th	10:30-11:15	QMES Floor 1	Introduction to QXU5017 Introduction to EXP2-1 Part1
Tuesday 21 st	14:00-15:40		Group Meeting 1 (Correct data presenting)
Friday 24 th	10:30-12:10		Group Meeting 2 (Write Report)
Monday 27 th	10:30-12:10	QMES Floor 1	Introduction to EXP2-1 Part2
Tuesday 28 th	14:00-15:40		Group Meeting 3 (Different sample comparison)
Friday 3 rd	10:30-12:10		Group Meeting 4 (Complete report)
Sunday 5 th	17:00	QMPlus	Report submission

EXP2-1 Schedule – P Groups



Date:	Time:	Place:	Activity:
Monday 20 th	14:00-14:45	QMES Floor 1	Introduction to QXU5017 Introduction to EXP2-1 Part1
Tuesday 21 st	10:30-12:10		Group Meeting 1 (Correct data presenting)
Friday 24 th	14:00-15:40		Group Meeting 2 (Write Report)
Monday 27 th	14:00-15:40	QMES Floor 1	Introduction to EXP2-1 Part2
Tuesday 28 th	10:30-12:10		Group Meeting 3 (Different sample comparison)
Friday 3 rd	14:00-15:40		Group Meeting 4 (Complete report)
Sunday 5 th	17:00	QMPlus	Report submission