

Ex. 3

T115952129

07.10.2026

Question 1

```
# Data
rent <- c(12500, 13800, 11200, 15000, 12500, 26000,
          14300, 13100, 12500, 16700, 14900, 13600)
```

```
# 1. Mean
mean(rent)
```

```
## [1] 14675
```

```
# 2. Median
median(rent)
```

```
## [1] 13700
```

```
# 3. Mode
table(rent)
```

```
## rent
## 11200 12500 13100 13600 13800 14300 14900 15000 16700 26000
##      1      3      1      1      1      1      1      1      1      1
```

Question 2

```
rent2 <- rent[-6]
mean(rent2)
```

```
## [1] 13645.45
```

```
median(rent2)
```

```
## [1] 13600
```

```
table(rent2)
```

```
## rent2
## 11200 12500 13100 13600 13800 14300 14900 15000 16700
##      1      3      1      1      1      1      1      1      1
```

The median better describes the typical rent because it is less affected by the unusually high NT\$26,000 rent.

Question 3

```
# Grades
grades <- c(88, 72, 80)

# Weights
weights <- c(0.36, 0.32, 0.32)

# Calculate the weighted mean
final_grade <- sum(grades * weights)

# Display the final grade
final_grade
```

```
## [1] 80.32
```