



A-Stability

example: chemical reactions (Robertson 1966)

- ▶ catalytic reactions:



- ▶ A converts to B which converts to C which drives conversion of B to A
- ▶ kinetic rate equations:

$$\frac{du_1}{dt} = -0.04u_1 + 10^4 u_2 u_3$$

$$\frac{du_2}{dt} = 0.04u_1 - 3 \cdot 10^7 u_2^2 - 10^4 u_2 u_3$$

$$\frac{du_3}{dt} = 3 \cdot 10^7 u_2^2$$


u_1 , u_2 and u_3 are concentrations of A, B and C, respectively

numerical solution – short time T and small step h

```
def f(t,u): # Robertson reaction
    du = u.copy()
    du[0] = -0.04*u[0] + 10**4*u[1]*u[2]
    du[1] = 0.04*u[0] - 3*10**7*u[1]**2 - 10**4*u[1]*u[2]
    du[2] = 3*10**7*u[1]**2
    return(du)

phi = f # Euler
n = 1024; T=0.016; h = T/n
tk = np.linspace(0,T,n+1)
uk = np.zeros((n+1,3))
uk[0,0] = 1.0
for j in range(n):
    uk[j+1,:] = uk[j,:] + h*phi(j*h,uk[j,:])
```

```
plt.semilogy(tk,uk[:,0],label='A')  
plt.semilogy(tk,uk[:,1],label='B')  
plt.semilogy(tk,uk[:,2],label='C')  
plt.axis(ymax=10)  
plt.legend();
```




larger time T and step size h

- larger step size to reduce computational time

```
n = 128; T=0.16; h = T/n
tk = np.linspace(0,T,n+1)
uk = np.zeros((n+1,3))
uk[0,0] = 1.0
for j in range(n):
    uk[j+1,:] = uk[j,:] + h*phi(j*h,uk[j,:])
```

- ▶ larger step sizes introduce (numerical) fluctuations

```
plt.semilogy(tk,uk[:,0],label='A')  
plt.semilogy(tk,abs(uk[:,1]),label='B')  
plt.semilogy(tk,uk[:,2],label='C')  
plt.axis(ymin=10)  
plt.legend();
```



Stability of solutions of ODEs

► ODE

$$\frac{du}{dt} = f(t, u)$$

Definition (stable solution)

A solution $u(t)$ of the ODE is stable if for some $\epsilon > 0$ and all solutions $v(t)$ of the ODE with $\|v(0) - u(0)\| \leq \epsilon$ the difference $v(t) - u(t)$ is bounded.

Example For $t \in \mathbb{R}_+$ and the differential equation

$$\frac{du}{dt} = -u$$

any solution $u(t)$ is stable.

Definition (unstable solution)

A solution $u(t)$ of the ODE is unstable if for all $\epsilon > 0$ there exists a solution $v(t)$ of the ODE with $\|v(0) - u(0)\| \leq \epsilon$ such that the difference $v(t) - u(t)$ is unbounded.

Example For $t \in \mathbb{R}_+$ and the differential equation

$$\frac{du}{dt} = u$$

any solution $u(t)$ is unstable.

Definition (asymptotically stable solution) A solution $u(t)$ of the ODE is asymptotically stable if for some $\epsilon > 0$ and all solutions $v(t)$ of the ODE with $\|v(0) - u(0)\| \leq \epsilon$ one has $\lim_{t \rightarrow \infty} \|u(t) - v(t)\| = 0$.

perturbation analysis of stable solutions of ODEs

- ▶ consider a family of solutions $u_\epsilon(t) = u(t) + \epsilon v(t)$ which all satisfy the same ODE

$$\frac{du_\epsilon}{dt} = f(t, u_\epsilon)$$

where u is an asymptotically stable solution

- ▶ as $v(t)$ is bounded one has, for twice continuously differentiable f the Taylor expansion

$$f(t, u_\epsilon) = f(t, u) + \epsilon f_u(t, u)v + O(\epsilon^2)$$

- ▶ it follows that v satisfies approximately the linear ODE

$$\frac{dv}{dt} = f_u(t, u)v$$

and consequently, the solutions of this ODE with sufficiently small initial values have to be bounded if the solution $u(t)$ is stable

- ▶ if for $t \rightarrow \infty$ one has $f_u(t, u) \rightarrow \lambda$ then all solutions of

$$\frac{dv}{dt} = \lambda v$$

have to be asymptotically stable and thus $\lambda < 0$

- ▶ when considering multiple variables $f_u(t, u)$ is the Jacobi matrix then think of λ as a (complex) eigenvalue and $u(t) \in \mathbb{C}$
- ▶ this motivates the study of the numerical solution of

$$du/dt = \lambda u$$

and $u(t) \in \mathbb{C}$

amplification factor

- ▶ consider family of complex ODEs with $f(t, u) = \lambda u$

$$\frac{du}{dt} = \lambda u$$

- ▶ all one-step methods considered construct $\phi(t, u)$ through compositions of linear combinations of evaluations of f
- ▶ it follows that for the family of ODEs considered one has

$$u_{k+1} = \rho(\lambda h) u_k$$

where $\rho(z)$ is a polynomial with real coefficients and $\rho(0) = 1$ for the (explicit) one-step methods

- ▶ in the following we will also consider *implicit methods* for which $\rho(z)$ is a rational function
- ▶ $\rho(\lambda h)$ is the *amplification factor* and

$$u_k = \rho(\lambda h)^k u_0$$

A-stability

- ▶ the ODEs considered are asymptotically stable if $\operatorname{Re}(\lambda) < 0$ and in this case the exact solution $u(t)$ satisfies

$$u(t) \rightarrow 0, \quad t \rightarrow \infty$$

- ▶ we say that a one-step method is *A-stable* if

$$u_k \rightarrow 0, \quad k \rightarrow \infty$$

and this is the case

$$|\rho(\lambda h)| < 1$$

- ▶ the *region of A-stability* of a one-step method is

$$\Omega = \{z \in \mathbb{C} \mid |\rho(z)| < 1\}$$

A-stability of Euler's method

- ▶ Euler's method for the ODEs considered gives

$$u_{k+1} = (1 + \lambda h)u_k$$

- ▶ amplification factor for Euler

$$\rho(z) = 1 + z$$

- ▶ region of A-stability


$$\Omega = \{z \mid |1 + z| < 1\}$$

is a circle in the complex plane with radius 1 and centre -1

plotting the region of A-stability of Euler's method

```
xg = np.linspace(-2,2,100)
yg = np.linspace(-2,2,100)
X, Y = np.meshgrid(xg,yg)
Rho = np.sqrt((1+X)**2 + Y**2)
```

```
plt.contourf(X,Y,Rho,[0.0,1.0,]); plt.axis('square');
```




A-stability and choice of h for $du/dt = \lambda u$ with $\lambda < 0$

- convergence for small h and unstability for large h

```
f = lambda t, u, lam=-20.2 : lam*u
phi = f # Euler
n = 200; h=0.1; T = n*h;
tk = np.linspace(0,T,n+1)
uk = np.ones(n+1)
for k in range(0,n):
    uk[k+1] = uk[k] + h*phi(k,uk[k])
```


- ▶ $\lambda h = -2.02$ (just) outside region of A-stability

```
plt.plot(tk,uk,'.-');
```



backward Euler method

- ▶ also implicit Euler method as rhs depends on u_{k+1}

$$u_{k+1} = u_k + hf(t_{k+1}, u_{k+1})$$

- ▶ numerical solution, typically iterative method

$$u_{k+1}^0 = u_k$$

$$u_{k+1}^{j+1} = \alpha u_{k+1}^j + (1 - \alpha)(u_k + hf(t_{k+1}, u_{k+1}^j)), \quad j = 0, \dots, m$$

- ▶ for the iteration we choose a relaxed fixed point iteration (sometimes called corrector) with $\alpha \in [0, 1]$

amplification factor for backward Euler

- ▶ one-step for model problem

$$u_{k+1} = u_k + \lambda h u_{k+1}$$

- ▶ solve

$$u_{k+1} = \frac{1}{1 - \lambda h} u_k$$

- ▶ amplification factor

$$\rho(z) = \frac{1}{1 - z}$$


- ▶ the iterative solver gives a slightly different amplification factor which should be investigated ...

plotting the region of A-stability for backward Euler

- ▶ backward Euler is unconditionally (for all h) A-stable for all λ with $\operatorname{Re}(\lambda) < 0$

```
xg = np.linspace(-2,2,100)
yg = np.linspace(-2,2,100)
X, Y = np.meshgrid(xg,yg)
Rho = 1.0/np.sqrt((1-X)**2 + Y**2)
```

```
plt.contourf(X,Y,Rho,[0.0,1.0,]); plt.axis('square');
```




previous example with backward Euler

- convergence for small h and instability for large h

```
lam=-20.2
f = lambda t, u, lam = lam : lam*u
phi = f # Euler forward
n = 200; h=0.1; T = n*h; m=4; alpha=0.5
tk = np.linspace(0,T,n+1)
uk = np.ones(n+1)
for k in range(0,n):
    ukp1 = uk[k]
    for j in range(m): # corrector: relaxed fixpoint
        ukp1 = alpha*ukp1 + (1-alpha)*(uk[k] + h*phi(k,ukp1))
    uk[k+1] = ukp1
```

- ▶ solution decreasing (with error)

```
plt.semilogy(tk,uk, label='numerical')  
plt.semilogy(tk,np.exp(lam*tk), label='exact')  
plt.legend();
```



backward Euler for Robertson example


```
def f(t,u): # Robertson reaction
    du = u.copy()
    du[0] = -0.04*u[0] + 10**4*u[1]*u[2]
    du[1] = 0.04*u[0] - 3*10**7*u[1]**2 - 10**4*u[1]*u[2]
    du[2] = 3*10**7*u[1]**2
    return(du)

phi = f # Euler
s=128; n = s*32; T=s*0.16; h = T/n
m = 10; alpha = 0.1
tk = np.linspace(0,T,n+1)
uk = np.zeros((n+1,3))
uk[0,0] = 1.0
for j in range(n):
    ukp1 = uk[j,:]
    for i in range(m):
        ukp1 = (1-alpha)*ukp1 + alpha*(uk[j,:] + h*phi(j*h,ukp1))
    uk[j+1,:] = uk[j,:] + h*phi(j*h,ukp1)
```


- ▶ fluctuations in B disappear even for large T

```
def plotting():  
    plt.semilogy(tk, uk[:,0], label='A')  
    plt.semilogy(tk, abs(uk[:,1]), label='B')  
    plt.semilogy(tk, uk[:,2], label='C')  
    plt.axis(ymax=10)  
    plt.legend();
```

plotting()



amplification factor for Runge-Kutta

- ▶ recall: 2 stages of Runge-Kutta

- ▶ stage 1:

$$U_k^j = u_k + h \sum_{i=1}^m b_{ji} f(s_i, U_k^i), \quad j = 1, \dots, m$$

- ▶ stage 2:

$$u_{k+1} = u_k + h \sum_{j=1}^m c_j U_k^j$$

- ▶ stage 1: define $\rho_j(z)$ for every U_k^j :

$$\rho_j(z) = 1 + z \sum_{i=1}^m b_{ji} \rho_i(z)$$

- ▶ one needs to solve this system of equations for ρ_j

- ▶ stage 2:

$$\rho(z) = 1 + z \sum_{j=1}^m c_j \rho_j(z)$$

example 2: Heun's method

$$B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad c^T = \begin{bmatrix} 0.5 & 0.5 \end{bmatrix}$$

- ▶ first stage

- ▶ $\rho_1(z) = 1$ and $\rho_2(\lambda h) = 1 + z$

- ▶ second stage

$$\rho(z) = 1 + 0.5z + 0.5z(1 + z) = 1 + z + z^2/2$$

example 3: midpoint method

$$B = \begin{bmatrix} 0 & 0 \\ 0.5 & 0 \end{bmatrix}, \quad c^T = \begin{bmatrix} 0 & 1 \end{bmatrix}$$

- ▶ first stage

- ▶ $\rho_1(z) = 1$ and $\rho_2(z) = 1 + z/2$


- ▶ second stage

$$\rho(z) = 1 + z(1 + z/2) = 1 + z + z^2/2$$

region of A-stability for Heun and midpoint method

```
xg = np.linspace(-4,2,100)
yg = np.linspace(-3,3,100)
X, Y = np.meshgrid(xg,yg)
Rho = np.sqrt((1+X+(X*X-Y*Y)/2)**2 + (Y+X*Y)**2)
```

```
plt.contourf(X,Y,Rho,[0.0,1.0,]); plt.axis('square');
```



example 4: fourth order Runge Kutta method

$$B = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0.5 & 0 & 0 & 0 \\ 0 & 0.5 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}, \quad c^T = \frac{1}{6} \begin{bmatrix} 1 & 2 & 2 & 1 \end{bmatrix}$$

► first stage

$$\rho_1(z) = 1$$

$$\rho_2(z) = 1 + z/2$$

$$\rho_3(z) = 1 + z/2 + z^2/4$$

$$\rho_4(z) = 1 + z + z^2/2 + z^3/4$$


► second stage

$$\rho(z) = 1 + z + z^2/2 + z^3/6 + z^4/24$$

plot A-stability region for Runge-Kutta

```
xg = np.linspace(-4,2,100)
yg = np.linspace(-3,3,100)
X, Y = np.meshgrid(xg,yg)
Z = X + Y*1j
Rho = abs(1+Z+Z**2/2+Z**3/6+Z**4/24)
```

```
plt.contourf(X,Y,Rho,[0.0,1.0,]); plt.axis('square');
```



example 5: trapezoidal rule

- ▶ implicit method:

$$u_{k+1} = u_k + 0.5\lambda h(u_k + u_{k+1})$$

- ▶ solve

$$u_{k+1} = \frac{1 + \lambda h/2}{1 - \lambda h/2} u_k$$

- ▶ amplification factor

$$\rho(z) = \frac{1 + z/2}{1 - z/2}$$

- ▶ $|\rho(z)| < 1$ for all z with $\operatorname{Re}(z) < 0$ and larger than 1 else

plot A-stability region for (implicit) trapezoidal rule

```
xg = np.linspace(-4,2,100)
yg = np.linspace(-3,3,100)
X, Y = np.meshgrid(xg,yg)
Z = X + Y*1j
Rho = abs((1+Z/2)/(1-Z/2))
```

```
plt.contourf(X,Y,Rho,[0.0,1.0,]); plt.axis('square');
```

