


Romberg

$$\bar{I} = \int_a^b f(x) dx$$

- ▶ (composite) trapezoidal rule for quadrature

$$T(f, h) = \frac{h}{2} \left( f(a) + 2 \sum_{k=1}^{n-1} f(\underline{a + kh}) + f(b) \right)$$


where  $h = (b - a)/n$  for some integer  $n$

- ▶ the Romberg method is Richardson extrapolation applied to the trapezoidal rule
- ▶ first we need to establish an error formula for the trapezoidal rule
- ▶ while in the case of central differences one considers a sequence of increasingly coarser grids, for Romberg one consider a sequence of increasingly finer grids

## example

$$f(x) = e^{-x}$$

```
f = lambda x : np.exp(-x)
intf = 1.0 - 1/np.e
```

```
# trapezoidal rule
```

```
T = lambda f,n,a=0.0,b=1.0 : (b-a)/n*((f(a)+f(b))/2.0 \
    + np.sum(f(np.linspace(a+(b-a)/n,b-(b-a)/n,n-1))))
d = 20 # might have to set lower ...
q = np.zeros(d)
nv = 2**np.arange(d)*4
for i,n in enumerate(nv):
    q[i] = T(f,n)
h = 1.0/nv
```

```
def pl1(): # plotting commands
```


```
    plt.title('trapezoidal rule')
```

```
    plt.gca().invert_xaxis();plt.loglog(h,h*h,label="$h^2$")
```

```
    plt.xlabel('h');plt.ylabel('error')
```

```
    plt.loglog(h,abs((q - intf)/intf),'. ',label="$Q(f)-I(f)$")
```

p11()



# Euler-Maclaurin

## Theorem

Let  $f \in C^{2k+1}[a, b]$  and  $h = (b - a)/n$ . Then

$$T(f, h) - \int_a^b f(x) dx = \sum_{k=1}^{m-1} h^{2k} \frac{B_{2k}}{(2k)!} \left( f^{(2k-1)}(b) - f^{(2k-1)}(a) \right) + \underline{h^{2m}} \frac{B_{2m}}{(2m)!} (b - a) f^{(2m)}(\xi)$$

where  $B_{2k}$  are the Bernoulli numbers.

- ▶ proof by induction and integration by parts
- ▶ first few Bernoulli numbers

$$B_0 = 1, B_2 = 1/6, B_4 = -1/30, B_6 = 1/42, B_8 = -1/30, \dots$$

- ▶ improve quadrature using derivatives
- ▶ case of  $m = 1$  see earlier discussion of trapezoidal rule
- ▶  $f$  periodic and analytic: error  $o(h^m)$  for all  $m$

- ▶ in this case: trapezoidal rule better than Gauss quadrature!

(see Wikipedia)

## Recursive computation of $T(2^{-k}h)$

- compute sequence  $T(f, 2^{-k}(b-a))$ , for  $k = 0, \dots, m$  by

$$T(f, b-a) = \frac{(b-a)(f(a) + f(b))}{2}$$

$$T(f, (b-a)/2) = \frac{T(f, b-a)}{2} + \frac{(b-a)f(a + (b-a)/2)}{2},$$

$$T(f, (b-a)/4) = \frac{T(f, (b-a)/2)}{2} + \frac{(b-a)(f(a + (b-a)/4) + f(a + 3(b-a)/4))}{4}$$

$\vdots$

$$\underline{T(f, (b-a)/2^m)} = \frac{T(f, (b-a)/2^{m-1})}{2} + \frac{(b-a) \sum_{i=1}^{2^{m-1}} f(a + (2i-1)(b-a)/2^m)}{2^m}$$

# Romberg integration = Richardson extrapolation for trapezoidal rule

- ▶ we use

$$T(f, h) = a_1 h^2 + a_2 h^4 + a_3 h^6 + \dots$$

and eliminate  $a_1$ , then  $a_2$ , etc


- ▶ establish lower triangular tableau  $R[j, k]$  with
  - ▶ leftmost column

$$R[j, 0] = T(2^{-j}h)$$

- ▶ elements

$$R[j, k] = \frac{4^k R[j, k-1] - R[j-1, k-1]}{4^k - 1}$$

- ▶ use the efficient method to update  $T(2^{-j}h)$
- ▶ error approximation  $e_h \approx R[j, j] - R[j+1, j+1]$



example  $f(x) = \exp(-x)$

$h$

$2h$

```
R = lambda f,n: (4*T(f,n) - T(f,n/2))/3
```

```
qr = np.zeros(d)
```

```
for i,n in enumerate(nv[1:]):
```

```
    qr[i] = R(f,n)
```

```
def pl2(): # plotting commands
```

```
    plt.title('trapezoidal rule')
```

```
    plt.gca().invert_xaxis();plt.loglog(h,h*h,label="$h^2$")
```

```
    plt.xlabel('h');plt.ylabel('error')
```


```
    plt.loglog(h,abs((q - intf)/intf),'. ',label="trapezoidal")
```

```
    plt.loglog(h,abs((qr - intf)/intf),'. ',label="extrapolated")
```

```
    plt.legend();
```




p12()



## Romberg tableau

- all the  $R(i, j)$  required for  $R(0, k)$ :



- follows from

$$R[j, k] = \frac{4^k R[j, k-1] - R[j-1, k-1]}{4^k - 1}$$

- use  $R(j, k)$  for error estimate on diagonal

$$e_j = R(j, j) - \phi(0) \approx R(j, j) - R(j+1, j+1)$$

## example

```
mr = 5; h = 0.1
R = np.zeros((mr,mr))
for n in range(mr):
    R[n,0] = T(f,2**n)
for k in range(1,mr):
    for j in range(k,mr):
        R[j,k] = (4**k*R[j,k-1] - R[j-1,k-1])/(4**k-1)
print(R)
print("exact value: {}".format(intf))

[[ 0.68393972  0.          0.          0.          0.
   0.64523519  0.63233368  0.          0.          0.
   0.63540943  0.63213418  0.63212088  0.          0.
   0.63294342  0.63212141  0.63212056  0.63212056  0.
   0.63232631  0.63212061  0.63212056  0.63212056  0.63212056]
exact value: 0.6321205588285577
```

## errors

```
for j in range(mr):  
    for k in range(j+1):  
        print("{:1.2e}".format(R[j,k]-intf),end='    ')  
    print("\n")
```

5.18e-02

1.31e-02      2.13e-04

3.29e-03      1.36e-05      3.16e-07

8.23e-04      8.56e-07      5.06e-09      1.23e-10

2.06e-04      5.36e-08      7.96e-11      4.94e-13      1.21e-14



## error estimate on diagonal

```
# error approximation for diagonal
```

```
print("approx",end='    ')
```

```
for k in range(mr-1):
```

```
    print("{:1.4e}".format(R[k,k]-R[k+1,k+1]),end='    ')
```

```
print("\nexact",end='    ')
```

```
for k in range(mr-1):
```

```
    print("{:1.4e}".format(R[k,k]-intf),end='    ')
```

```
approx    5.1606e-02  2.1280e-04  3.1606e-07  1.2341e-10
```

```
exact     5.1819e-02  2.1312e-04  3.1618e-07  1.2342e-10
```

## comments

- ▶ practical implementation will require stopping criteria based on error estimate
- ▶ the columns converge according to the Euler-Maclaurin formula
- ▶ the diagonal may not converge!
- ▶ in contrast to differentiation, rounding errors for integration are usually not a problem