

Instructor(s): Dr. Kenneth Duru  
First Semester 2019  
Mathematical Sciences Institute  
Australian National University

## Assignment 2

This assignment must be submitted by **5th April 5pm**. Late Submissions will incur a 5% penalty per working day. Assignment submissions will close on the **12th April 5pm**. Submissions after this time will be invalid.

### Question 1 (Gaussian Elimination)

- 1a. Compute a vector  $\mathbf{u}$  such that  $E = I - \mathbf{u}e_2^T$  satisfies

$$Ex = E(x_1, x_2, x_3, x_4)^T = (x_1, x_2, 0, 0)^T.$$

$$\mathbf{e}_2 = (0, 1, 0, 0)^T$$

- 1b. Show that

$$M = (I - \alpha e_3 e_2^T)(I - \beta e_4 e_2^T)$$

is an elementary matrix.

- 1c. Find a vector  $x$  for which  $Mx = (0, 4, 0, 0)^T$ .

### Question 2 (Linear Iterative Methods)

- 2a. Given a linear system of equations  $Ax = b$  with a symmetric positive definite matrix  $A \in \mathbb{R}^{4 \times 4}$  which has eigenvalues  $1, 1/4, 1/9, 1/16$ . Consider the iterative method defined by

$$x^{(k+1)} = x^{(k)} - \omega(Ax^{(k)} - b).$$

Can you choose  $\omega$  such that method is convergent? If so, what is the best possible  $\omega$ ?

- 2b. Discuss the convergence of the Jacobi method for  $Ax = b$  with the tridiagonal matrix

$$A = \begin{bmatrix} 4 & -1 & & \\ -1 & 5 & -1 & \\ & -1 & 6 & -1 \\ & & -1 & 3 \end{bmatrix}.$$

- Does the Jacobi method converge for this matrix?
- What is the convergence rate?

### Question 3 (Gradient Descent and CG)

3a. Show that for the steepest descent method the direction vectors  $d^{(k)}$  and  $d^{(k+1)}$  are orthogonal.

3b. Let  $A = \begin{bmatrix} 2 & -1 & & \\ -1 & 5 & -1 & \\ & -1 & 6 & -1 \\ & & -1 & 3 \end{bmatrix}$ . Show that  $A$  is positive definite. Discuss

the convergence of the conjugate gradient method for  $Ax = b$  with this matrix  $A$  and arbitrary vectors  $b$ .

### Question 4 (Nonlinear Equations and Bisection)

4a. Show that  $f(x) = x^4 - 8x - 2$  has a zero in the interval  $[0, 4]$ . 4b. How many iteration steps would you require with bisection to get an error  $|e^{(k)}| < 0.01$ ?

### Question 5 (Fixed Point Methods)

Consider  $f(x) = x^4 - 8x - 2$ . How would you choose  $\alpha$  such that

$$x^{(k+1)} = F(x^{(k)})$$

is convergent where  $F(x) = x - \alpha f(x)$  for  $x \in [0, 4]$ .

### Question 6 (Newton's Method)

Does the Newton's method for  $f(x) = x^4 - 8x - 2$  converge for any point.

Find the point for which the Newton's method does not converge for this function