1 Wave particle duality

$$u_v dv = \frac{8\pi v^2}{c^3} \cdot \frac{hv}{e^{\frac{hv}{K_b T}} - 1}$$
 (2)

$$E = hv (3)$$

$$\frac{1}{2}m_e v_k^2 = E_k = hv - W (5)$$

$$\frac{1}{\lambda} = R_H \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \tag{6}$$

$$R_H = 1.097373 \times 10^7 m^{-1}$$
 $n_1 = 1 \Rightarrow Lyman (7)$

$$n_1 = 2 \Rightarrow Balmer$$
 (8)

$$n_1 = 3 \Rightarrow Paschen$$
 (9)

$$n_1 = 4 \Rightarrow Brackett$$
 (10)

$$n_1 = 15 \Rightarrow Pfund \tag{11}$$

$$n_2 > n_1 \tag{12}$$

De Broglie Relationship: $E^2 = p^2c^2 + m^2c^4 \ (13)$

for zero rest mass:
$$E^2 = p^2 c^2 \to E = pc$$
 (14)

$$hv = h\frac{c}{\lambda} = pc \tag{15}$$

$$\therefore p = \frac{h}{\lambda} \tag{16}$$

$$E_K = eV - \frac{p^2}{2m} \tag{18}$$

$$p = \sqrt{2meV} \tag{19}$$

$$\lambda = \frac{h}{\sqrt{2meV}} \tag{20}$$

$$\lambda = 2d\sin(\theta) \tag{21}$$

$$\sin^2(\theta) = \frac{C}{V}$$
, Where $C = \frac{h^2}{8med^2}$ (22)

Bohr Model (23)

$$F_{centripedal} = F_{electric} \tag{24}$$

$$m\frac{v^2}{r} = \frac{e^2}{4\pi\epsilon_0 r^2} \tag{25}$$

$$\therefore v^2 = \frac{e^2}{4\pi\epsilon_0 mr} \tag{26}$$

$$I_{orbit} = n\lambda = 2\pi r$$
 Where n is the quantisation condition (27)

$$\therefore v = \frac{hn}{2\pi mr} \tag{28}$$

$$L = mvr = n\frac{h}{2\pi} = n\hbar \tag{29}$$

As such, allowed radii are described by the expression: (30)

$$r_n = \frac{h^2 \epsilon_0 n^2}{\pi m e^2} = a_{Bohr} n^2 \tag{31}$$

$$r_1 = a_{Bohr} = \frac{h^2 \epsilon_0}{\pi m e^2} = 5.3 \times 10^{-11} m \tag{32}$$

$$E_k = \frac{1}{2}mv^2 = \frac{e^2}{8\pi\epsilon_0 r} \tag{33}$$

$$E_T = E_k + E_c = \frac{e^2}{8\pi\epsilon_0 r} - \frac{e^2}{4\pi\epsilon_0 r^2} = -\frac{e^2}{8\pi\epsilon_0 r}$$
(34)

Substiting in
$$r_n$$
 gives: (35)

$$E_T = -\left(\frac{me^4}{8\epsilon_0^2 h^2}\right) \cdot \frac{1}{n^2} = \frac{-13.6eV}{n^2}, \ 1 \ eV \approx 1.602 \times 10^{-19} J \tag{36}$$

$$\Delta E = E_i - E_f = \frac{me^4}{8\epsilon_0^2 h^2} \left(\frac{1}{n_f^2} - \frac{1}{n_i^2}\right)$$
 (37)

$$textsince: E = hv = h\frac{c}{\lambda} \tag{38}$$

$$\frac{1}{\lambda} = \frac{me^4}{8\epsilon_0^2 h^3} \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right) = R_H \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$$
(39)