



Logistic Regression

测验, 5 个问题

1
point

1.

Suppose that you have trained a logistic regression classifier, and it outputs on a new example x a prediction $h_{\theta}(x) = 0.7$. This means (check all that apply):


- ☐ Our estimate for $P(y = 1|x; \theta)$ is 0.7.
- ☐ Our estimate for $P(y = 1|x; \theta)$ is 0.3.
- ☐ Our estimate for $P(y = 0|x; \theta)$ is 0.7.
- ☐ Our estimate for $P(y = 0|x; \theta)$ is 0.3.

1
point

2.

Suppose you have the following training set, and fit a logistic regression classifier $h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2)$.

x_1	x_2	y
1	0.5	0
1	1.5	0
2	1	1
3	1	0



Which of the following are true? Check all that apply.

- ☐ Adding polynomial features (e.g., instead using $h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_1^2 + \theta_4 x_1 x_2 + \theta_5 x_2^2)$) could increase how well we can fit the training data.
- ☐ At the optimal value of θ (e.g., found by fminunc), we will have $J(\theta) \geq 0$.
- ☐ Adding polynomial features (e.g., instead using $h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_1^2 + \theta_4 x_1 x_2 + \theta_5 x_2^2)$) would increase $J(\theta)$ because we are now summing over more terms.
- ☐ If we train gradient descent for enough iterations, for some examples $x^{(i)}$ in the training set it is possible to obtain $h_{\theta}(x^{(i)}) > 1$.

1
point

3.

For logistic regression, the gradient is given by $\frac{\partial}{\partial \theta_j} J(\theta) = \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$. Which of these is a correct gradient descent update for logistic regression with a learning rate of α ? Check all that apply.

Logistic Regression

测验, 5 个问题

- ☐ $\theta := \theta - \alpha \frac{1}{m} \sum_{i=1}^m (\theta^T x - y^{(i)}) x^{(i)}.$
- ☐ $\theta := \theta - \alpha \frac{1}{m} \sum_{i=1}^m \left(\frac{1}{1 + e^{-\theta^T x^{(i)}}} - y^{(i)} \right) x^{(i)}.$
- ☐ $\theta_j := \theta_j - \alpha \frac{1}{m} \sum_{i=1}^m (\theta^T x - y^{(i)}) x_j^{(i)}$ (simultaneously update for all j).
- ☐ $\theta := \theta - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x^{(i)}.$

1
point

4.

Which of the following statements are true? Check all that apply.


- ☐ The one-vs-all technique allows you to use logistic regression for problems in which each $y^{(i)}$ comes from a fixed, discrete set of values.
- ☐ The cost function $J(\theta)$ for logistic regression trained with $m \geq 1$ examples is always greater than or equal to zero.
- ☐ For logistic regression, sometimes gradient descent will converge to a local minimum (and fail to find the global minimum). This is the reason we prefer more advanced optimization algorithms such as fminunc (conjugate gradient/BFGS/L-BFGS/etc).
- ☐ Since we train one classifier when there are two classes, we train two classifiers when there are three classes (and we do one-vs-all classification).

1
point


5.

Suppose you train a logistic classifier $h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2)$. Suppose $\theta_0 = -6, \theta_1 = 0, \theta_2 = 1$. Which of the following figures represents the decision boundary found by your classifier?


☐ Figure:




☐ Figure:



☐ Figure:





☒ 我（志超 杨）了解提交不是我自己完成的作业 将永远不会通过此课程或导致我的 Coursera 帐号被关闭。了解荣誉准则的更多信息

Submit Quiz

