# On Sequences Associated to the Invariant Theory of Rank Two Lie Algebras

#### Yi Zhang

Department of Mathematical Sciences The University of Texas at Dallas, USA

Joint work with Alin Bostan, Jordan Tirrell and Bruce Westbury



# The On-Line Encyclopedia of Integer Sequences (OEIS)



OEIS is an online database of integer sequences, such as Fibonacci numbers (A000045), Catalan numbers (A000108).

Yi Zhang, UTD 2/2

## Two families of sequences in OEIS

a (OEIS tag)	0	1	2	3	4	5
A059710	1	0	1	1	4	10
A108307	1	1	2	5	15	51
A108304	1	2	5	15	52	202

The first family of sequences (octant sequences)

a (OEIS tag)	0	1	2	3	4	5
A151366	1	0	2	2	12	30
A236408	1	1	3	9	33	131
A001181	1	2	6	22	92	422
A216947	1	3	11	49	221	1113

The second family of sequences (quadrant sequences)

- ▶ Those sequences are associated to the invariant theory of the exceptional simple Lie algebra  $G_2$  of rank 2.
- The quadrant sequences are related to the octant sequences by the branching rules for SL(3) of  $G_2$ .

Yi Zhang, UTD 3/25

## Octant sequences

The first family of sequences can be interpreted as lattice walks restricted to the octant. We call them octant sequences.

- ▶ A059710: enumerates the multiplicities of the trivial representation in the tensor powers of V, which is the 7-D fundamental representation of  $G_2$ .
- ▶ A108307: enumerates enhanced 3-noncrossing set partitions.
- ▶ A108304: enumerates 3-noncrossing set partitions.

(Lin, 2018; Gil and Tirrell, 2019): A108307 and A108304 are related by the binomial transform.

Yi Zhang, UTD 4/2

#### Motivation and Contribution

(Bostan, Tirrell, Westbury and Z., 2019): A059710 and A108307 are also related by the binomial transform.

Mihailovs' conjecture: Let  $T_3(n)$  be the *n*-th term of A059710. Then  $T_3$  is determined by  $T_3(0) = 1$ ,  $T_3(1) = 0$ ,  $T_3(2) = 1$  and

$$14(n+1)(n+2) T_3(n) + (n+2)(19n+75) T_3(n+1) +2(n+2)(2n+11) T_3(n+2) - (n+8)(n+9) T_3(n+3) = 0.$$

(Bostan, Tirrell, Westbury and Z., 2019): Three independent proofs of Mihailovs' conjecture.

- ► Two proofs are based on binomial relation between A059710 and A108307, together with a result by Bousquet-Mélou and Xin.
- ▶ The third one is a direct proof by the method of algebraic residues, which leads to closed formulae for the generating function of *T*<sub>3</sub> in terms of hypergeometric functions.

Yi Zhang, UTD 5/25

## Quadrant sequences

The second family of sequences can be interpreted as lattice walks restricted to the quadrant. We call them quadrant sequences.

- ▶ A151366: enumerates nonpositive bipartite trivalent graphs.
- ▶ A236408: enumerates pasting diagrams.
- ▶ A001181: enumerates Baxter permutations.
- ▶ A216947: enumerates 2-coloured noncrossing set partitions.

Question: What are relations between quadrant sequences?

Yi Zhang, UTD 6/2

#### Motivation and Contribution

(Marberg, 2013): a combinatorial proof that A151366, A001181, and A216947 are related by binomial transforms.

(Bostan, Tirrell, Westbury and Z., 2019): Derive a uniform recurrence equation for quadrant sequences and show that they are related by binomial transform by the representation theory of simple Lie algebras.

Yi Zhang, UTD 7/2

## Outline

 binomial relation between the first and second octant sequences

▶ Three independent proofs of Mihailovs' conjecture

▶ The quadrant sequences are related by binomial transforms

Yi Zhang, UTD 8/25

#### **Preliminaries**

Definition 1 Let G be a reductive complex algebraic group and let V be a representation of G. The sequence associated to (G,V), denoted  $\mathbf{a}_V$ , is the sequence whose n-th term is the multiplicity of the trivial representation in the tensor power  $\otimes^n V$ .

Example 1 Let V be the 7-D fundamental representation of  $G_2$ . Then A059710 is the sequence associated with  $(G_2, V)$ .

Let **a** be a sequence with n-th term a(n), the binomial transform of **a** is the sequence, denoted  $\mathcal{B}\mathbf{a}$ , whose n-th term is

$$\sum_{i=0}^{n} \binom{n}{i} a(i).$$

Yi Zhang, UTD 9/2

#### **Preliminaries**

Lemma 1 Assume  $\mathbf{a}_V$  is the sequence associated to (G,V) as specified in Definition 1. Then  $\mathbf{a}_{V\oplus\mathbb{C}}=\mathcal{B}\mathbf{a}_V$ .

Lemma 2 Assume **a** enumerates walks in a lattice, confined to a domain D, using a set of steps S. Then  $\mathcal{B}\mathbf{a}$  also enumerates walks in a lattice restricted to D with steps  $S \coprod \{0\}$ .

Lemma 3 Let G(t) be the generating function of **a**. For  $k \in \mathbb{Z}$ , denote the generating function of  $\mathcal{B}^k$ **a** by  $\mathcal{B}^k G$ . Then

$$(\mathcal{B}^k G)(t) = \frac{1}{1-k \ t} G\left(\frac{t}{1-k \ t}\right).$$

Yi Zhang, UTD 10/25


Let V be the 7-D fundamental representation of  $G_2$ . Then

A059710 is the sequence associated to  $(G_2, V)$ . Let  $T_3(n)$  be its n-th term.


A108307 enumerates enhanced 3-noncrossing set partitions. Let  $E_3(n)$  be its n-th term.

Yi Zhang, UTD 11/25

In terms of lattice walks, we can interpret  $T_3$  and  $E_3$  as follows:




Steps in weight lattice of  $G_2$ 



Steps in octant related to  $E_3(n)$ 

Yi Zhang, UTD 12/25

In terms of lattice walks, we can interpret  $T_3$  and  $E_3$  as follows:



If we make a linear transformation  $(x, y) \rightarrow (x + y, y)$ , then it identifies the six non-zero steps, as well as the two domains.

Yi Zhang, UTD 12/25

Recall: Lemma 2 Assume **a** enumerates walks in a lattice, confined to a domain D, using a set of steps S. Then  $\mathcal{B}\mathbf{a}$  also enumerates walks in a lattice restricted to D with steps  $S \mid \{0\}$ .

Yi Zhang, UTD 13/25

Recall: Lemma 2 Assume **a** enumerates walks in a lattice, confined to a domain D, using a set of steps S. Then  $\mathcal{B}\mathbf{a}$  also enumerates walks in a lattice restricted to D with steps  $S \coprod \{0\}$ .

By Lemma 2 and the previous figures, we conclude that  $E_3$  is the binomial transform of  $T_3$ .

Yi Zhang, UTD 13/25

Recall: Lemma 2 Assume **a** enumerates walks in a lattice, confined to a domain D, using a set of steps S. Then  $\mathcal{B}\mathbf{a}$  also enumerates walks in a lattice restricted to D with steps  $S \mid \{0\}$ .

By Lemma 2 and the previous figures, we conclude that  $E_3$  is the binomial transform of  $T_3$ .

(Lin, 2018; Gil and Tirrell, 2019): A108307 and A108304 are related by the binomial transform.

Recall: Lemma 1 Assume  $\mathbf{a}_V$  is the sequence associated to (G, V) as specified in Definition 1. Then  $\mathbf{a}_{V \oplus \mathbb{C}} = \mathcal{B} \mathbf{a}_V$ .

Thus, the octant sequences are sequences associated to

$$(G_2, V)$$
,  $(G_2, V \oplus \mathbb{C})$ ,  $(G_2, V \oplus 2\mathbb{C})$ .

Yi Zhang, UTD 13/2

# First proof of Mihailovs' conjecture

Mihailovs' conjecture: Let  $T_3(n)$  be the *n*-th term of A059710. Then  $T_3$  is determined by  $T_3(0) = 1$ ,  $T_3(1) = 0$ ,  $T_3(2) = 1$  and

$$14(n+1)(n+2) T_3(n) + (n+2)(19n+75) T_3(n+1) +2(n+2)(2n+11) T_3(n+2) - (n+8)(n+9) T_3(n+3) = 0.$$

(Bousquet-Mélou and Xin, 2005): Let  $E_3(n)$  be the *n*-th term of A108307. Then  $E_3$  is given by  $E_3(0) = E_3(1) = 1$ , and

$$8(n+3)(n+1)E_3(n) + (7n^2 + 53n + 88)E_3(n+1) - (n+8)(n+7)E_3(n+2) = 0.$$

Yi Zhang, UTD 14/25

# First proof of Mihailovs' conjecture

Recall: We prove that  $E_3$  is the binomial transform of  $T_3$ . Thus,

$$T_3(n) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} E_3(k).$$

Set 
$$f(n,k) = (-1)^{n-k} \binom{n}{k} E_3(k)$$
.

- **b** By Bousquet-Mélou and Xin's result, f(n, k) is holonomic function, which satisfies ordinary difference equations for n and k, respectively.
- ▶ Idea: Using creative telescoping method (Zeilberger, 1990), which is an algorithmic approach to compute a differential/difference equation for the integration/sum of holonomic functions, to drive a recurrence equation for T<sub>3</sub>.

Yi Zhang, UTD 15/

# First proof of Mihailovs' conjecture

Using the Koutschan's Mathematica package HolonomicFunctions.m that implements Chyzak's algorithm for creative telescoping, we find exactly the recurrence equation in Mihailovs' conjecture.

Yi Zhang, UTD 16/25

# Second proof of Mihailovs' conjecture

Recall: We prove that  $E_3$  is the binomial transform of  $T_3$ . Let  $\mathcal{T}(t) = \sum_{n \geq 0} T_3(n)t^n$  and  $\mathcal{E}(t) = \sum_{n \geq 0} E_3(n)t^n$ . Then

$$\mathcal{T}(t) = rac{1}{1+t} \cdot \mathcal{E}\left(rac{t}{1+t}
ight).$$

- **)** By Bousquet-Mélou and Xin's result, we can derive an ODE for  $\mathcal{E}(t)$ .
- ▶ Using the closure properties of holonomic function (the sum, product and algebraic substitution of holonomic functions is still holonomic), we can derive an ODE for  $\mathcal{T}(t)$  and convert it into a linear recurrence for  $T_3(n)$ , which is exactly the recurrence equation in Mihailovs' conjecture.

Yi Zhang, UTD 17/2

# Third proof of Mihailovs' conjecture

Idea: In terms of lattice walks, we can interpret  $T_3(n)$  to be the constant term of  $W K^n$ , where

$$K = (1 + x + y + xy + x^{-1} + y^{-1} + (xy)^{-1}),$$

and

$$W = x^{-2}y^{-3}(x^2y^3 - xy^3 + x^{-1}y^2 - x^{-2}y + x^{-3}y^{-1} - x^{-3}y^{-2} + x^{-2}y^{-3} - x^{-1}y^{-3} + xy^{-2} - x^2y^{-1} + x^3y - x^3y^2).$$

Let  $\mathcal{T}(t) = \sum_{n\geq 0} T_3(n)t^n$ . Then  $\mathcal{T}(t)$  is the constant coefficient  $[x^0y^0]$  of W/(1-tK). In other words,  $\mathcal{T}(t)$  is equal to the algebraic residue of W/(xy-txyK), which is proportional to the contour integral of W/(xy-txyK) over a cycle.

Yi Zhang, UTD 18/25

# Third proof of Mihailovs' conjecture

Using creative telescoping method, we can compute a 6-th order ODE for  $\mathcal{T}(t)$ . Moreover, by using factorization of differential operators, we can show that  $L_3(\mathcal{T}(t))=0$ , where  $\partial=\frac{d}{dt}$  and

$$L_3 = t^2 (2 t + 1) (7 t - 1) (t + 1) \partial^3 + 2 t (t + 1) (63 t^2 + 22 t - 7) \partial^2 + (252 t^3 + 338 t^2 + 36 t - 42) \partial + 28 t (3 t + 4).$$

Converting it into a linear recurrence for  $T_3(n)$ , we get exactly the recurrence equation in Mihailovs' conjecture.

Yi Zhang, UTD 19/25

## Closed formulae

By factorization of the operator  $L_3$  and algorithms for solving 2-nd order ODEs, we derive the following closed formula for  $\mathcal{T}(t)$ :

$$\mathcal{T}(t) = \frac{1}{30 t^5} \left[ R_1 \cdot {}_2F_1 \left( \frac{1}{3} \, {}_2^{\frac{2}{3}} ; \phi \right) + R_2 \cdot {}_2F_1 \left( \frac{2}{3} \, {}_3^{\frac{4}{3}} ; \phi \right) + 5 P \right],$$

where

$$R_1 = \frac{(t+1)^2 (214 t^3 + 45 t^2 + 60 t + 5)}{t-1},$$

$$R_2 = 6 \frac{t^2 (t+1)^2 (101 t^2 + 74 t + 5)}{(t-1)^2},$$

and

$$\phi = \frac{27(t+1)t^2}{(1-t)^3}, \qquad P = 28t^4 + 66t^3 + 46t^2 + 15t + 1.$$

Yi Zhang, UTD

#### Closed formulae

By elliptic curve theory, we derive an alternative formula for  $\mathcal{T}(t)$ :

$$\frac{P}{6 t^5} + \frac{(7 t - 1) (2 t + 1) (t + 1)}{360 t^5} \Big( (155 t^2 + 182 t + 59) (11 t + 1) H(t) + (341 t^3 + 507 t^2 + 231 t + 1) (5 t + 1) H'(t) \Big),$$

where

$$H(t) = \frac{1}{g_2^{1/4}} \cdot {}_2F_1\left(\frac{1}{12}, \frac{5}{12}; \frac{1728}{J}\right),$$

$$J = \frac{(t-1)^3 (25 t^3 + 21 t^2 + 3 t - 1)^3}{t^6 (1-7 t) (2 t + 1)^2 (t+1)^3},$$

and

$$g_2 = (t-1)(25t^3 + 21t^2 + 3t - 1).$$

Yi Zhang, UTD 21/25

## Binomial relations between quadrant sequences

Definition 2 Let  $\tilde{V}$  be the defining representation of SL(3) and denote the dual by  $\tilde{V}^*$ . For  $k \ge 0$ , we define  $S_k$  to be the sequence associated to  $(SL(3), \tilde{V} \oplus \tilde{V}^* \oplus k \mathbb{C})$ .

Remark: SL(3) is the maximal subgroup of  $G_2$ . Let V be the 7-D fundamental representation of  $G_2$ . Then  $S_k$  is the the sequence associated to  $(SL(3), (V \oplus k\mathbb{C}) \downarrow_{SL(3)})$ .

Lemma 4 Let  $C_k$  be the generating function of  $S_k$ , where  $k \ge 0$ . Then  $C_k$  is the constant coefficient of  $[x^0y^0]$  of W/(1-tK), where

$$K = k + x + y + x^{-1} + y^{-1} + \frac{x}{y} + \frac{y}{x}$$

and

$$W = 1 - \frac{x^2}{y} + x^3 - x^2y^2 + y^3 - \frac{y^2}{x}.$$

Yi Zhang, UTD 22/

## Binomial relations between quadrant sequences

By Lemma 4,  $S_3$  is identical to the quadrant sequence A216947.

(Marberg, 2013): The *n*-th term  $C_2(n)$  of  $S_3$  is given by  $C_2(0) = 1$ ,  $C_2(1) = 3$  and

$$(n+5)(n+6) \cdot C_2(n+2) - 2(5n^2 + 36n + 61) \cdot C_2(n+1) + 9(n+1)(n+4) \cdot C_2(n) = 0.$$

By Lemma 1,  $S_k$ 's are related by binomial transforms. Thus, by Lemma 3, the generating function of  $S_k$  is

$$\mathcal{A}(t) := rac{1}{1-kt} \cdot \mathcal{C}\left(rac{t}{1-kt}
ight)$$

where C(t) is the generating function of  $S_3$ .

Yi Zhang, UTD 23/25

## Binomial relations between quadrant sequences

Using closure properties of holonomic functions, we derive a uniform 4-th order recurrence equation for  $S_k$  with k as a parameter.

By comparing the recurrence equations between  $S_k$ 's and quadrant sequences, and then checking initial terms, we show that

Theorem: The sequences  $S_0$ ,  $S_1$ ,  $S_2$ ,  $S_3$  are identical to quadrant sequences. In particular, quadrant sequences are related by binomial transforms.

Yi Zhang, UTD 24/25

## Summary

- A combinatorial proof of binomial relation between the first and second octant sequences
- Three independent proofs of Mihailovs' conjecture
  - ▶ Two proofs are based on binomial relation between the first and second octant sequences
  - A direct proof by the method of algebraic residues, which leads to closed formulae for the generating function of the first octant sequence
- The quadrant sequences are related by binomial transforms

Yi Zhang, UTD 25/25

## Summary

- A combinatorial proof of binomial relation between the first and second octant sequences
- Three independent proofs of Mihailovs' conjecture
  - ▶ Two proofs are based on binomial relation between the first and second octant sequences
  - A direct proof by the method of algebraic residues, which leads to closed formulae for the generating function of the first octant sequence
- ▶ The quadrant sequences are related by binomial transforms

Thanks!

Yi Zhang, UTD 25/25