

Lecture 10: Conditioning and Stability

Amos Ron

University of Wisconsin - Madison

February 22, 2021

Outline

1 Conditioning

- Introduction to the notion of conditioning
- The condition number of a matrix

2 Stability

- Stability of factorization algorithms
- Stability of QR -factorization

Outline

1 Conditioning

- Introduction to the notion of conditioning
- The condition number of a matrix

2 Stability

- Stability of factorization algorithms
- Stability of QR -factorization

Blank page

Solving a linear system in a finite precision environment

We are given a square invertible $m \times m$ A , and two equations:

- An equation

$$Ax = b_0, \quad x_0 := A^{-1}b_0.$$

- An equation

$$Ax = b_1, \quad x_1 := A^{-1}b_1,$$

with $\|x_0\|$ and $\|x_1\|$ of about the same order.

Solving a linear system in a finite precision environment

We are given a square invertible $m \times m$ A , and two equations:

- An equation

$$Ax = b_0, \quad x_0 := A^{-1}b_0.$$

- An equation

$$Ax = b_1, \quad x_1 := A^{-1}b_1,$$

with $\|x_0\|$ and $\|x_1\|$ of about the same order.

Question:

How large/small can the ratio

$$\frac{\|b_0\|}{\|b_1\|}$$

be?

Solving a linear system in a finite precision environment

Why should we care about such question?

Assume, for example, that

$$\frac{\|b_0\|}{\|b_1\|} = 10^{12}$$

Choose $b_2 = 10^{-4}b_1$. Then

- Denote $x_2 := A^{-1}b_2$, then $x_2 = 10^{-4}x_1$, therefore, more or less,

$$\|x_2\| \approx 10^{-4}\|x_0\|.$$

- But:

$$\|b_2\| = 10^{-4}\|b_1\| = 10^{-16}\|b_0\|.$$

Solving a linear system in a finite precision environment

Summary of the current discussion

- We assumed that, for the given A , there exist x_0, x_1 , such that

$$||x_0|| \approx ||x_1||, \quad \text{and} \quad ||Ax_0|| \approx 10^{12}||Ax_1||.$$

- We concluded that there exist b_0, b_2 , such that

$$||b_0|| \approx 10^{16}||b_2|| \quad \text{and} \quad ||A^{-1}b_0|| \approx 10^4||A^{-1}b_2||.$$

Solving a linear system in a finite precision environment

Summary of the current discussion

- We assumed that, for the given A , there exist x_0, x_1 , such that

$$\|x_0\| \approx \|x_1\|, \quad \text{and} \quad \|Ax_0\| \approx 10^{12} \|Ax_1\|.$$

- We concluded that there exist b_0, b_2 , such that

$$\|b_0\| \approx 10^{16} \|b_2\| \quad \text{and} \quad \|A^{-1}b_0\| \approx 10^4 \|A^{-1}b_2\|.$$

This is a disaster!

Consider the two equations $Ax = b_0$, and $Ax = b_0 + b_2$. In Matlab, the two systems are the same. But the solutions of the two systems are very different!

The definition of condition number

Definition: Condition Number

A is any matrix (not necessarily square). $\|\cdot\|$ is some norm.

$$\text{cond}(A) := \frac{\max\{\|Av\| : \|v\| = 1\}}{\min\{\|Av\| : \|v\| = 1\}}.$$

The definition of condition number

Definition: Condition Number

A is any matrix (not necessarily square). $\|\cdot\|$ is some norm.

$$\text{cond}(A) := \frac{\max\{\|Av\| : \|v\| = 1\}}{\min\{\|Av\| : \|v\| = 1\}}.$$

Note: If A is square and invertible, then

$$\text{cond}(A) = \|A\| \|A^{-1}\|.$$

The 2-condition number

Example: A is 2×2

$$\text{cond}(A) = s(1)/s(2).$$

The 2-condition number

Example: Q is orthogonal

$$\text{cond}(Q) = 1.$$

Outline

1 Conditioning

- Introduction to the notion of conditioning
- The condition number of a matrix

2 Stability

- Stability of factorization algorithms
- Stability of QR -factorization

Condition numbers in the context of factorization algorithms

When solving $Ax = b$, we factored $A = BC$. We need now to solve

- $By = b$.
- $Cx = y$.

So, the condition numbers that matter are of B and of C !

Condition numbers in the context of factorization algorithms

When solving $Ax = b$, we factored $A = BC$. We need now to solve

- $By = b$.
- $Cx = y$.

So, the condition numbers that matter are of B and of C !
Not hard to show:

$$\text{cond}(A) \leq \text{cond}(B)\text{cond}(C).$$

The problem is that it is possible that

$$\text{cond}(A) \ll \text{cond}(B)\text{cond}(C).$$

In this case **the factorization algorithm is unstable**.

Stability of QR -factorization

Theorem: QR -factorization is as stable as it gets!

Assume $A = QR$, then:

- $\text{cond}(A) = \text{cond}(R)$.
- $\text{cond}(Q) = 1$.