

American Psychiatric Association (APA)

United States

As the leading psychiatric organization in the world, APA has more than 40,000 members involved in psychiatric practice, research, and academia representing the diversity of the patients for whom they care. APA Clinical Practice Guidelines provide evidence-based recommendations for the assessment and treatment of psychiatric disorders and are intended to assist in clinical decision making by presenting systematically developed patient care strategies in a standardized format.

<https://www.psychiatry.org/>

Link to recommendations: <https://www.psychiatry.org/psychiatrists/practice/clinical-practice-guidelines/legacy-practice-guidelines-for-major-depressive-di>

Table

Recommended treatments by APA according to their empirical support

Strong	Cognitive-behavioural therapy ^{1, 3, 6, 7, 7+, 8+} , Interpersonal psychotherapy ^{1, 6, 7, 7+, 8}
Moderate	Marital therapy, Family therapy, Psychodynamic therapy ^{1, 6, 7, 7+, 8+}
Weak/Controversial	Group cognitive-behavioural therapy ^{1, 3, 6, 7, 7+, 8+} , Group interpersonal psychotherapy ^{1, 6, 7, 7+, 8} , Group problem-solving therapy ^{1, 6, 7, 7+, 8+} , Group psychodynamic therapy ^{1, 6, 7, 7+, 8+} , Problem-solving therapy ^{1, 6, 7, 7+, 8+}
Insufficient	-
No evidence assesment	-
First line	Cognitive-behavioural therapy ⁺ , Interpersonal psychotherapy ⁺