

Assignment 1

Name: 陈子蔚 SID: 12332440

1

As for $\mathbf{0}$, $\forall i, d_i = 0 \wedge s = 0 \wedge E = 0$. Only 1 case.

As for signed floating numbers, $0 < d_0 \leq \beta - 1 \wedge 1 \leq 0 \leq \beta - 1, 0 < i \leq p - 1 \wedge L \leq E \leq U$.

$(\beta - 1)\beta^{p-1}(U - L + 1)$ cases.

Totally, $(\beta - 1)\beta^{p-1}(U - L + 1) + 1$ cases.

The smallest fraction part is 1. The smallest exponent part is L . The smallest positive floating number is $1 \times \beta^L = \beta^L$.

The largest fraction part is $(\beta - \beta^{-(p-1)})$. The largest exponent part is U . The largest positive floating number is $(\beta - \beta^{-(p-1)}) \times \beta^U = \beta^{U+1}(1 - \beta^{-p})$.

2

a

$$\epsilon = \sin(x + h) - \sin(x) \quad (1)$$

b

$$\epsilon_r = \frac{\sin(x + h) - \sin(x)}{\sin(x)} \quad (2)$$

c

$$\kappa = \frac{\left| \frac{\sin(x+h) - \sin(x)}{\sin(x)} \right|}{\left| \frac{h}{x} \right|} \quad (3)$$

$$= \frac{\left| \frac{\sin(x+h) - \sin(x)}{\sin(x)} \right|}{\left| \frac{h}{x} \right|} \quad (4)$$

$$= \left| \frac{\sin(x + h) - \sin(x)}{h} \cdot \frac{x}{\sin(x)} \right| \quad (5)$$

$$\simeq \left| \frac{x \cos(x)}{\sin(x)} \right| \quad (6)$$


$$= \left| \frac{x}{\tan(x)} \right| \quad (7)$$

d


The problem is highly sensitive where the condition number is extremely large, which typically occurs when $\tan(x)$ is close to 0.

3

a



b



4

b

A tolerance such that it is less than the machine epsilon was used, which is a reasonable estimate for when to stop the series summation.

Output

```
1 # x_list: [1, -1, 5, -5, 10, -10, 15, -15, 20, -20]
2 19
3 19
4 37
5 37
6 53
7 53
8 68
9 68
10 83
11 83
12 # max order
13 [(1, 2.7182818284590455, 2.718281828459045, 4.440892098500626e-16),
14 (-1, 0.36787944117144245, 0.36787944117144233, 1.1102230246251565e-
15 16),
16 (5, 148.4131591025766, 148.4131591025766, 0.0),
17 (-5, 0.006737946999084642, 0.006737946999085467, 8.248610128269718e-
18 16),
19 (10, 22026.465794806714, 22026.465794806718, 3.637978807091713e-12),
20 (-10, 4.5399929670419935e-05, 4.5399929762484854e-05,
21 9.206491905638936e-14),
22 (15, 3269017.3724721107, 3269017.3724721107, 0.0),
23 (-15, 3.059100025508472e-07, 3.059023205018258e-07,
24 7.682049021416746e-12),
25 (20, 485165195.40979046, 485165195.4097903, 1.7881393432617188e-07),
26 (-20, 6.147561848704381e-09, 2.061153622438558e-09,
27 4.086408226265824e-09)]
28 # [(x, my_exp, builtin_exp, error)]
```

5

$$\|x\|_1 = \sum_{1 \leq i \leq n} |x_i| \quad (8)$$

$$\|x\|_2 = \sqrt{\sum_{1 \leq i \leq n} |x_i|^2} \quad (9)$$

$$\|x\|_\infty = \max_{1 \leq i \leq n} |x_i| \quad (10)$$

Notice that,

$$\forall i, |x_i| \leq \sum_{1 \leq i \leq n} |x_i| = \|x\|_1 \quad (11)$$

As for the 1st inequality,

$$\|x\|_2^2 = \sum_{1 \leq i \leq n} |x_i|^2 = \sum_{1 \leq i \leq n} |x_i| \cdot |x_i| \quad (12)$$

$$\leq \sum_{1 \leq i \leq n} |x_i| \cdot \|x\|_1 = \|x\|_1 \cdot \sum_{1 \leq i \leq n} |x_i| = \|x\|_1^2 \quad (13)$$

Since $\|x\|_2 \geq 0 \wedge \|x\|_1 \geq 0$,

$$\|x\|_2 \leq \|x\|_1 \quad (14)$$

Then,

$$\|x\|_1 = \sum_{1 \leq i \leq n} |x_i| = x^\top [1, \dots, 1]^\top \leq \|x\|_2 \cdot \|[1, \dots, 1]^\top\|_2 = \sqrt{n} \|x\|_2 \quad (15)$$

Thus,

$$\|x\|_2 \leq \|x\|_1 \leq \sqrt{n} \|x\|_2 \quad (16)$$

As for the 2nd inequality, suppose $j = \arg \max_{0 \leq i \leq n} |x_i|$,

$$\|x\|_\infty^2 = x_j^2 \leq \sum_{1 \leq i \leq n, i \neq j} x_i^2 + x_j^2 = \sum_{1 \leq i \leq n} x_i^2 = \|x\|_2^2 \quad (17)$$

That is,

$$\|x\|_\infty \leq \|x\|_2 \quad (18)$$

Notice that,

$$\forall i, |x_i| \leq |x_j| \quad (19)$$

Then,

$$\|x\|_2 = \sqrt{\sum_{1 \leq i \leq n} |x_i|^2} \leq \sqrt{\sum_{1 \leq i \leq n} |x_j|^2} = \sqrt{n |x_j|^2} = \sqrt{n} |x_j| = \sqrt{n} \|x\|_\infty \quad (20)$$

6

a&b

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 2 \\ 1 & 2 & 1 & 4 \\ 1 & 3 & 2 & 6 \end{array} \right] \xrightarrow{r_2 - r_1, r_3 - r_1} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 2 \\ 0 & 1 & 1 & 2 \\ 0 & 2 & 2 & 4 \end{array} \right] \xrightarrow{r_2 - \frac{1}{2} r_3} \left[\begin{array}{ccc|c} 1 & 1 & 0 & 2 \\ 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{array} \right] \quad (21)$$

There are only 2 pivots in A and $b \in c(A)$ since $x = [1, 1, 1]^\top$ is a possible solution.

Thus, A is singular and there are infinite solutions to $Ax = b$.

c

$$\|A\|_1 = \max_{1 \leq j \leq n} \|A_{(:,j)}\|_1 = 6 \quad (22)$$

$$\|A\|_\infty = \max_{1 \leq i \leq n} \|A_{(i,:)}\|_1 = 6 \quad (23)$$

d

Since A is singular,

$$\text{cond}(A) = \infty \quad (24)$$

7

a

Note that L here is the L^{-1} in the slide.

$$U = L_m P_m \cdots L_1 P_1 A = LPA \quad (25)$$

$$P = P_m \cdots P_1 \quad (26)$$

$$L = L_m \cdots L_1 P_1 P^\top \quad (27)$$

b

$$Ax = b \Leftrightarrow LPAx = Ux = LPb \quad (28)$$

$$A[x_1, \dots, x_n] = [b_1, \dots, b_n] \Leftrightarrow I = [e_1, \dots, e_n] = AA^{-1} \quad (29)$$

c

$$\text{cond}(A) = \|A\| \|A^{-1}\| \quad (30)$$

Output

```
1 # P
2 [[0. 1. 0.]
3  [0. 0. 1.]
4  [1. 0. 0.]]
5 # L
6 [[ 1.    0.    0. ]
7  [-1.    1.    0. ]
8  [ 0.25 -0.5   1.  ]]
9 # U
10 [[4.  4.  2. ]
11  [0.  2.  2. ]
12  [0.  0.  0.5]]
13 # A_inv
14 [[ 1.   1.  -1. ]
15  [-2.  -1.  1.5]
16  [ 2.   0.5 -1. ]]
17 # A_inv @ A
18 [[1. 0. 0.]
19  [0. 1. 0.]
```

```

20 | [0. 0. 1.]]
21 | # cond_1
22 | 60.0
23 | # cond_inf
24 | 63.0

```

8

The code of `LPU_factorization` and `Ux_equal_b_solution` in [7](#) is also used here. Moreover, the package `scipy` is used to check the correctness.

Output

```

1 | [-28.28427125  20.           10.           -30.           14.14213562
2 |    20.           0.          -30.           7.07106781  25.
3 |    20.          -35.35533906  25.           ]
4 | [ 0.00000000e+00  0.00000000e+00 -3.55271368e-15 -1.77635684e-15
5 |    0.00000000e+00  0.00000000e+00  8.88178420e-16  0.00000000e+00
6 |    0.00000000e+00  0.00000000e+00  8.88178420e-16  4.44089210e-15
7 |    7.10542736e-15]
8 | # solutions by the manual program
9 | [-28.28427125  20.           10.           -30.           14.14213562
10 |    20.           0.          -30.           7.07106781  25.
11 |    20.          -35.35533906  25.           ]
12 | [ 0.00000000e+00  0.00000000e+00 -3.55271368e-15 -1.77635684e-15
13 |    0.00000000e+00  0.00000000e+00  8.88178420e-16  0.00000000e+00
14 |    0.00000000e+00  0.00000000e+00  8.88178420e-16  4.44089210e-15
15 |    0.00000000e+00]
16 | # solutions by scipy

```