

$$(i) \quad \nabla \times (\nabla \times \underline{u}) = \nabla(\nabla \cdot \underline{u}) - \nabla^2 \underline{u}$$

$$\begin{aligned} \nabla \times (\nabla \times \underline{u}) &= \epsilon_{ijk} \partial_j (\nabla \times \underline{u})_k = \\ &= \epsilon_{ijk} \partial_j (\epsilon_{klm} \partial_l u_m) = \epsilon_{ijk} \epsilon_{klm} \partial_j \partial_l u_m = \\ &= \epsilon_{kij} \epsilon_{klm} \partial_j \partial_l u_m = (\delta_{il} \delta_{jm} - \delta_{jl} \delta_{im}) \partial_j \partial_l u_m = \\ &= \partial_j \partial_i u_j - \partial_l \partial_l u_i = \nabla(\nabla \cdot \underline{u}) - \nabla^2 \underline{u} \quad \text{AS REQUIRED.} \end{aligned}$$

✓

$$\nabla \times (\underline{u} \times \underline{v}) = \underline{u}(\nabla \cdot \underline{v}) - \underline{v}(\nabla \cdot \underline{u}) + (\underline{v} \cdot \nabla) \underline{u} - (\underline{u} \cdot \nabla) \underline{v}$$

$$\begin{aligned} \nabla \times (\underline{u} \times \underline{v}) &= \epsilon_{ijk} \partial_j (\underline{u} \times \underline{v})_k = \epsilon_{ijk} \partial_j (\epsilon_{klm} u_l v_m) = \\ &= \epsilon_{kij} \epsilon_{klm} \partial_j (u_l v_m) = (\delta_{il} \delta_{jm} - \delta_{im} \delta_{jl}) (v_m \partial_j u_l + \\ &\quad + u_l \partial_j v_m) = \\ &= v_j \partial_j u_i - v_i \partial_l u_l + u_l \partial_l v_m - u_j \partial_j v_i = \end{aligned}$$

✓

$$= (\underline{v} \cdot \nabla) \underline{u} - \underline{v}(\nabla \cdot \underline{u}) + \underline{u}(\nabla \cdot \underline{v}) - (\underline{u} \cdot \nabla) \underline{v} =$$

REARRANGE:

$$= \underline{u}(\nabla \cdot \underline{v}) - \underline{v}(\nabla \cdot \underline{u}) + (\underline{v} \cdot \nabla) \underline{u} - (\underline{u} \cdot \nabla) \underline{v} \quad \text{AS DESIRED.}$$

(ii) DIVERGENCE THEOREM:

$$\int_V (\nabla \cdot \underline{F}) dV = \oint_S \underline{F} \cdot d\underline{S}$$

WHERE \underline{F} IS
A VECTOR FIELD,
 V IS A VOLUME
BOUNDED BY
CLOSED SURFACES,
 $d\underline{S}$ IS OUTWARD
POINTING NORMAL
ON SURFACE (i.e. $\underline{n} dS$)

$$\int_V \nabla \cdot (\underline{F} \times \underline{G}) dV = \oint_S (\underline{F} \times \underline{G}) \cdot \underline{ds} \quad \text{BY DIVERGENCE THM.}$$

$$\nabla \cdot (\underline{F} \times \underline{G}) = \partial_i (\epsilon_{ijk} F_j G_k) =$$

$$= \epsilon_{ijk} [(\partial_i F_j) G_k + F_j (\partial_i G_k)] =$$

$$= G_k \epsilon_{kij} \partial_i F_j + \epsilon_{jki} F_j \partial_i G_k =$$

$$= \underline{G} \cdot (\nabla \times \underline{F}) - \underline{F} \cdot (\nabla \times \underline{G})$$


say which properties of ϵ are used.
CYCLIC PROPERTY

PLUG BACK IN:

$$\int_V [\underline{G} \cdot (\nabla \times \underline{F}) - \underline{F} \cdot (\nabla \times \underline{G})] dV = \oint_S (\underline{F} \times \underline{G}) \cdot \underline{\hat{n}} dS$$

(iii)

$$\underline{F} = (xz \sin(yz) + x^3, \cos(yz), 3zy^2 - e^{x^2+y^2})$$



$$\nabla \cdot \underline{F} = \left(\frac{\partial}{\partial x} (xz \sin(yz) + x^3) + \frac{\partial}{\partial y} (\cos(yz)) + \frac{\partial}{\partial z} (3zy^2 - e^{x^2+y^2}) \right) =$$


$$= z \sin(yz) + 3x^2 - z \sin(yz) + 3y^2 =$$

$$= 3x^2 + 3y^2 = 3(x^2 + y^2) \quad \checkmark$$

$$\oint_S \underline{F} \cdot \underline{\hat{n}} dS = \int_V (\nabla \cdot \underline{F}) dV = \int_V 3(x^2 + y^2) dV =$$

2014P1Q1(III)

$$\int_{z=0}^4 \int_{r=0}^{\sqrt{4-z}} \int_{\theta=0}^{2\pi} 3r^2 r d\theta dr dz =$$



$$z = 4 - x^2 - y^2$$

$$4 - z = x^2 + y^2 = r^2$$

$$r = \sqrt{4 - z}$$

$$= 2\pi \int_{z=0}^4 \int_{r=0}^{\sqrt{4-z}} 3r^3 dr dz =$$

$$= 2\pi \int_{z=0}^4 3 \left[\frac{r^4}{4} \right]_0^{\sqrt{4-z}} dz = 2\pi \int_{z=0}^4 3 \frac{(4-z)^2}{4} dz = \frac{3}{2}\pi \int_{z=0}^4 (4-z)^2 dz =$$

$$= \frac{3}{2}\pi \int_{z=0}^4 16 - 8z + z^2 dz = \frac{3}{2}\pi \left([16z]_0^4 - 8 \left[\frac{z^2}{2} \right]_0^4 + \left[\frac{z^3}{3} \right]_0^4 \right) =$$

$$= \frac{3}{2}\pi (64 - 64 + \frac{164}{3}) = \underline{\underline{32\pi}} \quad \checkmark$$