

2013P2Q4(D)(i)

CAUCHY'S RESIDUE THEOREM:

$$\oint_C f(z) dz = 2\pi i \sum \text{RESIDUES OF } f \text{ INSIDE } C$$

$$f(z) = f(z_0) \cancel{(z-z_0)^0} + f'(z_0)(z-z_0)^1 + \frac{1}{2!} f''(z_0)(z-z_0)^2 + \dots$$

$$\frac{f(z)}{z-z_0} = f(z_0)(z-z_0)^{-1} + f'(z_0) + \frac{1}{2!} f''(z_0)(z-z_0)^1 + \frac{1}{3!} f'''(z_0)(z-z_0)^2 + \dots$$

RESIDUE OF

$$\frac{f(z)}{z-z_0} \text{ IS THIS}$$

SO BY CRT:

$$\oint_C \frac{f(z)}{z-z_0} dz = 2\pi i f(z_0)$$

(THIS IS JUST CAUCHY'S INTEGRAL FORMULA)

$$(ii) \frac{e^{iz}}{a^2+z^2} = \frac{e^{iz}}{(z-ia)(z+ia)}$$

SO THE SINGULARITIES

OF ~~THE~~ THEINTEGRAND: $z = \pm ia$

THESE ARE FIRST ORDER POLES.


FROM

RESIDUE THEOREM

$$\oint_C \frac{f(z)}{z-z_0} dz = 2\pi i f(z_0)$$

WE COULD:

$$\oint_C \frac{e^{iz}}{(z-ia)(z+ia)} dz = \oint_C \frac{\frac{e^{iz}}{z+ia}}{z-ia} dz = 2\pi i \frac{e^{i(ia)}}{(ia)-(-ia)} = 2\pi i \frac{e^{-a}}{2ia} = \frac{\pi}{a} e^{-a}$$



$C = C_{sc} + C_{re}$
 PATH ALONG SEMICIRCLE ABOVE REAL AXIS, RADIUS = R

PATH ALONG REAL LINE FROM -R TO R

$$\oint_C \frac{e^{iz}}{a^2+z^2} dz = \underbrace{\oint_{C_{sc}} \frac{e^{iz}}{a^2+z^2} dz}_{I_{sc}} + \underbrace{\int_{C_{re}} \frac{e^{iz}}{a^2+z^2} dz}_{I_{re}}$$

AS $R \rightarrow \infty$, $I_{sc} \rightarrow 0$, BECAUSE LENGTH OF C_{sc} IS INCREASING PROPORTIONAL TO π , WHILE THE VALUE OF INTEGRAND $\left(\frac{e^{iz}}{a^2+z^2}\right)$ GOES TO 0 WITH A RATE $\frac{1}{r^2}$ AS R (AND z) $\rightarrow \infty$; $O(\pi)$ INCREASE MULTIPLIED BY $O\left(\frac{1}{r^2}\right)$ DECREASE RESULTS IN $O\left(\frac{1}{r}\right)$ DECREASE, SO $I_{sc} \rightarrow 0$.

$$\lim_{R \rightarrow \infty} I_{re} = \int_{-\infty}^{\infty} \frac{e^{ix}}{a^2+x^2} dx = \oint_C \frac{e^{iz}}{a^2+z^2} dz = \frac{\pi}{a} e^{-a}$$

2013 PZQ 4(II)

$$\frac{\pi}{a} e^{-a} = \int_{-\infty}^{\infty} \frac{e^{ix}}{a^2 + x^2} dx = \int_{-\infty}^{\infty} \frac{\cos x}{a^2 + x^2} dx + i \int_{-\infty}^{\infty} \frac{\sin x}{a^2 + x^2} dx$$

$\int_{-\infty}^{\infty} \frac{\sin x}{a^2 + x^2} dx = 0$ (ODD)

$$\Rightarrow \int_{-\infty}^{\infty} \frac{\cos x}{a^2 + x^2} dx = \frac{\pi}{a} e^{-a}$$

(iii) $I = \int_0^{2\pi} \frac{1}{1 + \epsilon \cos \theta} d\theta \quad -1 < \epsilon < 1$

$z = e^{i\theta} \rightarrow \cos \theta = \frac{z + \frac{1}{z}}{2}$

$\Rightarrow dz = i e^{i\theta} d\theta \Rightarrow d\theta = \frac{1}{iz} dz$

$$I = \oint_{|z|=1} \frac{1}{1 + \epsilon \frac{z + \frac{1}{z}}{2}} \frac{1}{iz} dz = \oint_{|z|=1} \frac{1(-i)}{z + \epsilon(\frac{z^2 + 1}{2})} dz = \oint_{|z|=1} \frac{-2i}{2z + \epsilon z^2 + \epsilon} dz$$

CLOSED CONTOUR INTEGRAL.

POLES OF INTEGRAND WHERE DENOMINATOR = 0

INTEGRATION PATH:
UNIT CIRCLE
CENTERED AT
ORIGIN, CCW.

$$2z + \epsilon z^2 + \epsilon = 0$$


$$\epsilon z^2 + 2z + \epsilon = 0$$

$$z_{\pm} = \frac{-2 \pm \sqrt{4 - 4\epsilon^2}}{2\epsilon} = \frac{-2 \pm 2\sqrt{1 - \epsilon^2}}{2\epsilon} = \frac{-1 \pm \sqrt{1 - \epsilon^2}}{\epsilon}$$

REWRITE INTEGRAND:

$$\frac{-2i}{2z + \epsilon z^2 + \epsilon} = \frac{-2i}{\epsilon(z - z_+)(z - z_-)}$$

BOTH TERMS INVOLVING z ARE
TO THE POWER 1 \Rightarrow THE
POLES ARE SIMPLE.



BY VIETA'S: $z_+ z_- = \frac{\epsilon}{\epsilon} = 1 \Rightarrow |z_+| \text{ OR } |z_-| \text{ IS ALWAYS LESS OR EQUAL TO } 1$

$$z_+ z_- = 1 \text{ IF } \epsilon = \pm 1$$

\Rightarrow THIS CANNOT BE
(NOT ALLOWED
BY PROBLEM),

SO:

$$|z_+| |z_-| < 1$$


\Rightarrow ONE POLE IS ALWAYS
INSIDE UNIT CIRCLE.

2013 P2Q4(III)

$$\oint_{|z|=1} \frac{-zi}{\epsilon(z-z_+)(z-z_-)} dz = I$$

$$|z|=1$$

$$\text{IF } |z_+| < 1:$$



$$I = \oint_{|z|=1} \frac{-zi}{\epsilon(z-z_+)(z-z_-)} dz = f(z_+) \cdot 2\pi i = 2\pi i \cdot \frac{-zi}{\epsilon(z_+-z_-)} =$$

$$= \frac{4\pi}{\epsilon(z_+-z_-)} = \frac{4\pi}{\epsilon \left(\frac{-1+\sqrt{1-\epsilon^2}}{\epsilon} - \frac{-1-\sqrt{1-\epsilon^2}}{\epsilon} \right)} =$$

$$= \frac{4\pi}{2\sqrt{1-\epsilon^2}} = \frac{2\pi}{\sqrt{1-\epsilon^2}}$$

$$\text{IF } |z_-| < 1:$$

$$I = \oint_{|z|=1} \frac{-zi}{\epsilon(z-z_+)(z-z_-)} dz = 2\pi i \cdot f(z_-) = 2\pi i \cdot \frac{-zi}{\epsilon(z_--z_+)} =$$

$$= \frac{4\pi}{\epsilon(z_--z_+)} = \frac{4\pi}{\epsilon \left(\frac{-1-\sqrt{1-\epsilon^2}}{\epsilon} - \frac{-1+\sqrt{1-\epsilon^2}}{\epsilon} \right)} =$$

$$= \frac{2\pi}{-\sqrt{1-\epsilon^2}}$$

THE RESULTS DISAGREE, SO WE SHOULD FIND IF $|z_+| < 1$ OR $|z_-| < 1$.

$$z_- = \frac{-1-\sqrt{1-\epsilon^2}}{\epsilon} \quad \left[\begin{array}{l} -1-\sqrt{1-\epsilon^2} < -1 \\ 1 > \epsilon > -1 \end{array} \right] \Rightarrow |z_-| > 1 \Rightarrow |z_+| < 1, \text{ so:}$$

$$\oint_{|z|=1} \frac{-zi}{\epsilon(z-z_+)(z-z_-)} dz = \int_0^{2\pi} \frac{1}{1+\epsilon \cos \theta} d\theta = \frac{2\pi}{\sqrt{1-\epsilon^2}}$$