

2014P2Q6(I)

(i) TRANSFORMATION LAW FOR A TENSOR OF ORDER n :

$$T'_{i_1 \dots i_n} = L_{i_1 j_1} \dots L_{i_n j_n} T_{j_1 \dots j_n}$$

WHERE $L_{ij} = \underline{e}_i' \cdot \underline{e}_j$ ✓

ISOTROPIC TENSORS
ARE TENSORS WITH
THE SAME COMPONENTS
IN ALL FRAMES: ✓

$$T'_{ij\epsilon \dots} = T_{ij\epsilon}$$

WHERE \underline{e}_i' IS THE i -TH
BASIS VECTOR OF THE
FRAME WE ARE
TRANSFORMING TO

& \underline{e}_j IS THE j -TH

✓ BASIS VECTOR OF THE
FRAME WE ARE TRANS-
FORMING FROM.
v bad grammar!

(ii) CONSIDER: $\frac{\partial x_j}{\partial x_i'} = \frac{\partial (L_{\epsilon j} x_{\epsilon}')}{\partial x_i'} = L_{\epsilon j} \frac{\partial x_{\epsilon}'}{\partial x_i'} =$

$$x_i' = L_{ij} x_j \quad | \cdot L_{ji}$$

$$L_{ji} x_i' = L_{ji} L_{ij} x_j \quad | \text{to.}$$

$$L_{ji} x_i' = x_j \quad \text{SINCE } L \text{ IS ORTHOGONAL.}$$

$$= L_{\epsilon j} \delta_{\epsilon i} = L_{ij}$$

BY CHAIN RULE:

$$\frac{\partial}{\partial x_i'} = \frac{\partial x_j}{\partial x_i'} \frac{\partial}{\partial x_j} = L_{ij} \frac{\partial}{\partial x_j}$$

TRANSFORMATION LAW
SATISFIED SO $\frac{\partial}{\partial x_i}$ IS

A TENSOR (OF ORDER 1).

~~what is your
def of a
tensor?~~ ✓
Is all this
needed for
4 marks?

$$\left(\frac{\partial u_i}{\partial x_j} \right)' = \left(\frac{\partial}{\partial x_j} \right)' (u_i)' = L_{jq} \frac{\partial}{\partial x_q} L_{ip} u_p = L_{jq} L_{ip} \frac{\partial u_p}{\partial x_q}$$

SO $\frac{\partial u_p}{\partial x_q}$ TRANSFORMS LIKE A TENSOR, SO IT IS A TENSOR.
(WE HAVE 2 INDICES, SO ITS AN ORDER 2 ONE.)

2014P2QC(II)

SAT 14:30 KETRY
REVISION
SUPO
BACKINGHAM(iii) TRANSFORMATION
LAW FOR AN
AXIAL VECTOR:

$$a'_i = \det(L) L_{ij} a_j$$

is this the
question?OBEYS THE SAME
TRANSFORMATION
LAW AS A VECTOR IF:

$$\det(L) = 1$$

$$(\nabla \times u)_i = (\epsilon_{ij\ell})' (\partial_j x_\ell) =$$

 ϵ IS A PSEUDOTENSOR,
THAT'S WHY IT TRANSFORMS LIKE THIS.

$$= \det(L) L_{ip} L_{jq} L_{\ell r} \epsilon_{pqr} L_{j\alpha} L_{\ell\beta} \partial_\alpha x_\beta =$$

$$= \det(L) L_{ip} L_{jq} L_{j\alpha} L_{\ell r} L_{\ell\beta} \epsilon_{pqr} \partial_\alpha x_\beta =$$

$$= \det(L) L_{ip} \delta_{q\alpha} \delta_{r\beta} \epsilon_{pqr} \partial_\alpha x_\beta =$$

$$= \det(L) L_{ip} \epsilon_{pqr} \partial_q x_r =$$

$$= \det(L) L_{ip} (\nabla \times u)_p$$

 $\nabla \times u$ TRANSFORMS AS AN AXIAL-VECTOR,SO IT CONSTITUTES AN AXIAL-VECTOR
(FIELD).

$$(iv) \frac{\partial u_i}{\partial x_j} = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right) + \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} - \frac{\partial u_j}{\partial x_i} \right)$$

SYMMETRIC
SECOND ORDER
TENSORANTISYMMETRIC
SECOND ORDER
TENSOR

$$\epsilon_{ijk} u_k = \begin{pmatrix} 0 & u_3 & -u_2 \\ -u_3 & 0 & u_1 \\ u_2 & -u_1 & 0 \end{pmatrix}$$

THIS IS THE GENERAL
FORM OF ANTISYMMETRIC
SECOND ORDER TENSORS,
SO ANY OF SUCH CAN BE EQUAL TO IT.

2016 P2 Q6 (III)

$$\epsilon_{ijk} \omega_k = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} - \frac{\partial u_j}{\partial x_i} \right)$$

$$\frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right) = \hat{S}_{ij}$$

SYMMETRIC ORDER 2 TENSOR.

$$\hat{S}_{ij} = p \delta_{ij} + \tau_{ij}$$

~~$p = +p$~~

$$p = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_i} + \frac{\partial u_j}{\partial x_j} \right) = \frac{\partial u_i}{\partial x_i}$$

we want τ_{ij}

$$\tau_{ij} = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right) - \frac{\partial u_i}{\partial x_i}$$

$$\omega_k = \frac{1}{\epsilon_{ijk}} \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} - \frac{\partial u_j}{\partial x_i} \right)$$

(v) $u_i = (dx_2, bx_1, 0)$

$$p = \frac{\partial u_i}{\partial x_i} = \frac{\partial}{\partial x_1} dx_2 + \frac{\partial}{\partial x_2} bx_1 + \frac{\partial}{\partial x_3} 0 = 0$$

$$\omega_k = \frac{1}{\epsilon_{ijk}} \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} - \frac{\partial u_j}{\partial x_i} \right)$$

IF $k=1$:

$$\omega_1 = \frac{1}{\epsilon_{231}} \frac{1}{2} \left(\frac{\partial u_2}{\partial x_3} - \frac{\partial u_3}{\partial x_2} \right) = + \frac{1}{\epsilon_{321}} \frac{1}{2} \left(\frac{\partial u_3}{\partial x_2} - \frac{\partial u_2}{\partial x_3} \right) =$$

$$= \frac{1}{2} \left(\frac{\partial}{\partial x_3} (bx_1) - \frac{\partial}{\partial x_2} 0 \right) = 0$$

not the full method

2014 P2 Q6 (IV)

For $q=2$:

$$W_2 = \frac{1}{\epsilon_{132}} \frac{1}{2} \left(\frac{\partial \mu_1}{\partial x_3} - \frac{\partial \mu_3}{\partial x_1} \right) = + \frac{1}{\epsilon_{312}} \frac{1}{2} \left(\frac{\partial \mu_3}{\partial x_1} - \frac{\partial \mu_1}{\partial x_3} \right) =$$

$$= + \frac{1}{2} \left(\frac{\partial}{\partial x_1} 0 - \frac{\partial}{\partial x_3} a x_2 \right) = 0$$

For $q=3$

$$W_3 = \frac{1}{\epsilon_{123}} \frac{1}{2} \left(\frac{\partial \mu_1}{\partial x_2} - \frac{\partial \mu_2}{\partial x_1} \right) = \frac{1}{2} \left(\frac{\partial}{\partial x_2} (a x_2) - \frac{\partial}{\partial x_1} (b x_1) \right) =$$

$$= \frac{1}{2} (a - b)$$

SO WE HAVE: $W_q = \delta_{q3} \frac{1}{2} (a - b)$

$$\frac{\partial \mu_i}{\partial x_j} = \begin{pmatrix} \frac{\partial a x_2}{\partial x_1} & \frac{\partial a x_2}{\partial x_2} & \frac{\partial a x_2}{\partial x_3} \\ \frac{\partial b x_1}{\partial x_1} & \frac{\partial b x_1}{\partial x_2} & \frac{\partial b x_1}{\partial x_3} \\ \frac{\partial 0}{\partial x_1} & \frac{\partial 0}{\partial x_2} & \frac{\partial 0}{\partial x_3} \end{pmatrix} = \begin{pmatrix} 0 & a & 0 \\ b & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \text{well yes. directly.}$$

$$\hat{S}_{ij} = \frac{1}{2} \left(\frac{\partial \mu_i}{\partial x_j} + \frac{\partial \mu_j}{\partial x_i} \right) = \begin{pmatrix} 0 & \frac{a+b}{2} & 0 \\ \frac{a+b}{2} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \begin{array}{l} \text{THIS IS ALREADY} \\ \text{✓ TRACELESS, SO:} \\ \hat{S}_{ij} = \hat{S}_{ji} \end{array}$$

$$\begin{aligned} \text{Tr } \hat{S} = 0 &\Rightarrow \sum_{i=1}^3 \lambda_i = 0 \\ \det \hat{S} = 0 &\Rightarrow \prod_{i=1}^3 \lambda_i = 0 \end{aligned} \quad \left. \vphantom{\begin{aligned} \text{Tr } \hat{S} = 0 \\ \det \hat{S} = 0 \end{aligned}} \right\} \rightarrow \lambda_1 = 0, \quad \lambda_2 = -\lambda_3 \quad \text{good.}$$

~~$$\begin{pmatrix} \frac{a+b}{2} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \frac{a+b}{2} \begin{pmatrix} y \\ x \\ 0 \end{pmatrix} \Rightarrow x = y = \frac{1}{\sqrt{2}} \sqrt{\frac{2}{a+b}}$$~~

2016 P2 Q6 (V)

$$\begin{pmatrix} 0 & \frac{a+b}{2} & 0 \\ \frac{a+b}{2} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \lambda_2 \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \frac{a+b}{2} y \\ \frac{a+b}{2} x \\ 0 \end{pmatrix}$$

too sparse

$$\left. \begin{aligned} \lambda_2 x &= \frac{a+b}{2} y \\ \lambda_2 y &= \frac{a+b}{2} x \\ x^2 + y^2 &= 1 \end{aligned} \right\} \begin{aligned} &\cancel{x \neq y} \\ &\lambda_2 x = \left(\frac{a+b}{2} \right)^2 \frac{x}{\lambda_2} \end{aligned}$$

$$\lambda_2^2 = \left(\frac{a+b}{2} \right)^2$$

$$e_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \leftarrow \lambda_2 = \frac{a+b}{2}$$

$$e_3 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$$

$$\lambda_3 = -\frac{a+b}{2}$$

$$e_1 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

PRINCIPAL
VALUES

PRINCIPAL
AXES

Very good ideas
but too inefficient.