

New plasmids for low-noise gene expression in *Escherichia coli*

This manuscript ([permalink](#)) was automatically generated from [zach-hensel/low-noise-manuscript@ee6757d](#) on August 31, 2018.

Authors

- **Zach Hensel**

 [0000-0002-4348-6229](#) ·  [zach-hensel](#) ·  [alchemytoday](#)

Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, Oeiras, Portugal · Funded by Project LISBOA-01-0145-FEDER-007660 (Microbiologia Molecular, Estrutural e Celular)

- **João Silva**

 [XXXX-XXXX-XXXX-XXXX](#)

Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, Oeiras, Portugal

Abstract

Some microbiology experiments and biotechnology applications can be improved if it is possible to tune the expression of two different genes at the same time with cell-to-cell variation at or below the level of genes constitutively expressed from the chromosome (the “extrinsic noise limit”). This was recently achieved for a single gene by exploiting negative autoregulation by the tetracycline repressor, and bicistronic gene expression to reduce gene expression noise. We report several new plasmids that use the same principles to achieve simultaneous, low-noise expression for two genes. This system was moved to a compatible plasmid backbone, and a system based on the lactose repressor was found to also exhibit gene expression noise below the extrinsic noise limit. We further investigated the effects of increasing repressor binding strength and the addition of a weak constitutive promoter on protein induction and expression noise.

Introduction

We recently reported the development of a plasmid-based gene-expression system in which a gene of interest is expressed bicistronically with the tetracycline repressor (TetR) [1]. Using this gene expression system, cell-to-cell is below the “extrinsic noise limit” observed for genes expressed from the chromosome [2]. When TetR and GFP are expressed bicistronically, GFP induction and gene expression noise is similar to that observed for a TetR-GFP fusion protein with autoregulation [3]. Compared to induction of gene expression under the control of a constitutively expressed transcriptional repressor, the inducer dose-response is relatively linearized, and gene expression noise is much lower at intermediate induction levels.

Methods

Strain construction

All plasmids were constructed using isothermal assembly [4] of fragments generated by PCR or double stranded DNA synthesis (IDT) and transformed Top10 E. coli cells (IBA #5-1600-020). Transformants were screened by colony PCR and verified by sequencing. Sequence maps are included as supplementary files and plasmids will be made available at AddGene after deposit and screening [5]. Purified plasmids were transformed into E. coli strain MG1655 by growing 2 mL of culture in SOB media at 37 C to OD600=0.4, washing twice with 1 mL ice-cold water, resuspending in 40 µL water, electroporation of 1–10 ng plasmid with the EC1 setting of a Micropulser (BioRad), and recovering for 1 hour at 37 C in SOC media.

Plasmid pZH509 [1] was used as a template to amplify the bicistronic regulatory construct including the $P_{\text{LtetO-1}}$ promoter, GFPmut2 [6], tn10 TetR [7] and rrnB T1 transcription terminator [8]. This was assembled by isothermal assembly with the backbone from pGB2 [9] with the pSC101 origin of replication and spectinomycin resistance to generate plasmid pJS101. Plasmids with the pSC101 and p15a origins of replication have been co-transformed in previous synthetic biology experiments [10].

Cell Growth

test

Flow Cytometry

blah

Data analysis

ok

Microscopy

123

References

1. A plasmid-based Escherichia coli gene expression system with cell-to-cell variation below the extrinsic noise limit

Zach Hensel

PLOS ONE (2017-10-30) <https://doi.org/10.1371/journal.pone.0187259>

2. Quantifying E. coli Proteome and Transcriptome with Single-Molecule Sensitivity in Single Cells

Y. Taniguchi, P. J. Choi, G.-W. Li, H. Chen, M. Babu, J. Hearn, A. Emili, X. S. Xie

Science (2010-07-29) <https://doi.org/10.1126/science.1188308>

3. Noise in transcription negative feedback loops: simulation and experimental analysis

Yann Dublanche, Konstantinos Michalodimitrakis, Nico Kümmerer, Mathilde Foglierini, Luis Serrano

Molecular Systems Biology (2006-08-01) <https://doi.org/10.1038/msb4100081>

4. Enzymatic assembly of DNA molecules up to several hundred kilobases

Daniel G Gibson, Lei Young, Ray-Yuan Chuang, J Craig Venter, Clyde A Hutchison, Hamilton O Smith

Nature Methods (2009-04-12) <https://doi.org/10.1038/nmeth.1318>

5. Addgene: Zach Hensel Lab Plasmidshttps://www.addgene.org/Zach_Hensel/

6. FACS-optimized mutants of the green fluorescent protein (GFP).

BP Cormack, RH Valdivia, S Falkow

Gene (1996) <https://www.ncbi.nlm.nih.gov/pubmed/8707053>

7. Nucleotide sequence of the repressor gene of the TN10 tetracycline resistance determinant

Kathleen Postle, Toai T. Nguyen, Kevin P. Bertrand

Nucleic Acids Research (1984) <https://doi.org/10.1093/nar/12.12.4849>

8. Intrinsic termination of T7 RNA polymerase mediated by either RNA or DNA.

L Hartvig, J Christiansen

The EMBO journal (1996-09-02) <https://www.ncbi.nlm.nih.gov/pubmed/8887568>

9. A pSC101-derived plasmid which shows no sequence homology to other commonly used cloning vectors.

G Churchward, D Belin, Y Nagamine

Gene (1984-11) <https://www.ncbi.nlm.nih.gov/pubmed/6098521>

10. BglBrick vectors and datasheets: A synthetic biology platform for gene expression

Taek Lee, Rachel A Krupa, Fuzhong Zhang, Meghdad Hajimorad, William J Holtz, Nilu Prasad, Sung Lee, Jay D Keasling

Journal of Biological Engineering (2011) <https://doi.org/10.1186/1754-1611-5-12>