

# Factoring Structural Attributes in Infrastructures' Deterioration Analysis

Zachary Hamida Doctorant  
James-A. Goulet Professeur



Polytechnique Montréal, Canada  
Département des génies civil, géologique et des mines

September 19, 2019

*Transports,  
Mobilité durable  
et Électrification  
des transports*

Québec 

Partenaire

# Outline

---

## **Contexte & Objectifs**

## **Deterioration Model**

## **Regression Framework**

## **Synthetic Data Analyses**

## **Next Steps**

---

# Database of Visual Inspections


Year: 2017

Structure: #002



# Database of Visual Inspections

structure



Year: 2017  
Structure: #002



# Database of Visual Inspections

structure

002

2008

2010

2012

2014

2016


2018

année

Year: 2017  
Structure: #002



# Database of Visual Inspections




Year: 2017

Structure: #002



# Database of Visual Inspections




Year: 2017  
Structure: #002



$\mathbf{b}_{002}$ : {type, material, DJMA, Location, ... }

# Example: Series of Inspections on Structural Element



# Objectives

- **Model the deterioration** behaviour based on the data from network of bridges

# 📍 Plan de la section


---

## Deterioration Model


- 2.1 Proposed Deterioration Model
  - 2.2 Model Performance
-

## Proposed Deterioration Model


## Deterioration Model Flowchart




# Deterioration Model Performance: Condition Estimate




# Deterioration Model Performance: Condition Estimate




# Deterioration Model Performance: Condition Estimate



# Deterioration Model Performance: Condition Estimate



# Deterioration Model Performance: Speed Estimate



# Limitations & Opportunities:

## Limitations:

# Limitations & Opportunities:

## Limitations:

- The estimate of the deterioration speed is of limited capacity.

# Limitations & Opportunities:

## Limitations:

- The estimate of the deterioration speed is of limited capacity.
- The model is based on deterioration data only.

# Limitations & Opportunities:

## Limitations:

- The estimate of the deterioration speed is of limited capacity.
- The model is based on deterioration data only.

# Limitations & Opportunities:

## Limitations:

- The estimate of the deterioration speed is of limited capacity.
- The model is based on deterioration data only.

## Opportunities:

# Limitations & Opportunities:

## Limitations:

- The estimate of the deterioration speed is of limited capacity.
- The model is based on deterioration data only.

## Opportunities:

- There are similarities across the structures/structural elements.

# Limitations & Opportunities:

## Limitations:

- The estimate of the deterioration speed is of limited capacity.
- The model is based on deterioration data only.

## Opportunities:

- There are similarities across the structures/structural elements.

**Use regression to incorporate the similarity information across the structures.**

# 📍 Plan de la section

---

## Regression Framework

- 3.1 Regression Formulation
  - 3.2 Kernel Regression: Derivation
-

# Concept & Notations

For structural attributes:  $\mathcal{Z} = \{z_1, z_2, \dots, z_B\}$

# Concept & Notations

For structural attributes:  $\mathcal{Z} = \{z_1, z_2, \dots, z_B\}$

Define a similarity measure across structures,

$$k(z_i, z_j) = \exp \left( -\frac{1}{2} \sum_{p=1} \left( \frac{z_{i,p} - z_{j,p}}{l_p} \right)^2 \right).$$

# Concept & Notations

For structural attributes:  $\mathcal{Z} = \{z_1, z_2, \dots, z_B\}$

Define a similarity measure across structures,

$$k(z_i, z_j) = \exp \left( -\frac{1}{2} \sum_{p=1} \left( \frac{z_{i,p} - z_{j,p}}{l_p} \right)^2 \right).$$

$$k(z_i, z_j) \in [0, 1]$$

**$z_i$  is the same for all structural elements in bridge  $b_i$**

# Kernel Regression

$$h(z) = \frac{\sum_{i=1}^N k(z, z_i)x_{z,i}}{\sum_{j=1}^N k(z, z_j)}$$

- $N$ : number of control point (unknown).

# Kernel Regression

$$h(z) = \frac{\sum_{i=1}^N k(z, z_i)x_{z,i}}{\sum_{j=1}^N k(z, z_j)}$$

- $N$ : number of control point (unknown).
- $k(., .)$ : the kernel function.

# Kernel Regression

$$h(z) = \frac{\sum_{i=1}^N k(z, z_i)x_{z,i}}{\sum_{j=1}^N k(z, z_j)}$$

- $N$ : number of control point (unknown).
- $k(., .)$ : the kernel function.
- $z_i$ : the  $i$ -th control point (a fixed point within the range of values for the structural attribute).

# Kernel Regression

$$h(z) = \frac{\sum_{i=1}^N k(z, z_i)x_{z,i}}{\sum_{j=1}^N k(z, z_j)}$$

- $N$ : number of control point (unknown).
- $k(., .)$ : the kernel function.
- $z_i$ : the  $i$ -th control point (a fixed point within the range of values for the structural attribute).
- $x_{z,i}$ : the  $i$ -th hidden control point (unknown).

## Regression Formulation

## Kernel Regression

$$h(z) = \frac{k(z, z_1)}{\sum_{j=1}^N k(z, z_j)} x_{z,1} + \frac{k(z, z_2)}{\sum_{j=1}^N k(z, z_j)} x_{z,2} + \cdots + \frac{k(z, z_N)}{\sum_{j=1}^N k(z, z_j)} x_{z,N},$$

## Regression Formulation

## Kernel Regression

$$h(z) = \frac{k(z, z_1)}{\sum_{j=1}^N k(z, z_j)} x_{z,1} + \frac{k(z, z_2)}{\sum_{j=1}^N k(z, z_j)} x_{z,2} + \cdots + \frac{k(z, z_N)}{\sum_{j=1}^N k(z, z_j)} x_{z,N},$$

$$\lambda_i = \frac{k(z, z_i)}{\sum_{j=1}^N k(z, z_j)},$$

## Regression Formulation

## Kernel Regression

$$h(z) = \frac{k(z, z_1)}{\sum_{j=1}^N k(z, z_j)} x_{z,1} + \frac{k(z, z_2)}{\sum_{j=1}^N k(z, z_j)} x_{z,2} + \cdots + \frac{k(z, z_N)}{\sum_{j=1}^N k(z, z_j)} x_{z,N},$$

$$\lambda_i = \frac{k(z, z_i)}{\sum_{j=1}^N k(z, z_j)},$$

$$h(z) = [ \begin{array}{cccc} \lambda_1 & \lambda_2 & \dots & \lambda_N \end{array} ] \times \begin{bmatrix} x_{z,1} \\ x_{z,2} \\ \dots \\ x_{z,N} \end{bmatrix}$$

## Regression Formulation

## Kernel Regression

$$h(z) = \frac{k(z, z_1)}{\sum_{j=1}^N k(z, z_j)} x_{z,1} + \frac{k(z, z_2)}{\sum_{j=1}^N k(z, z_j)} x_{z,2} + \cdots + \frac{k(z, z_N)}{\sum_{j=1}^N k(z, z_j)} x_{z,N},$$

$$\lambda_i = \frac{k(z, z_i)}{\sum_{j=1}^N k(z, z_j)},$$

$$\dot{x}_0 = h(z) = [ \lambda_1 \quad \lambda_2 \quad \dots \quad \lambda_N ] \times \begin{bmatrix} x_{z,1} \\ x_{z,2} \\ \dots \\ x_{z,N} \end{bmatrix}$$

## Regression Formulation

## Kernel Regression

$$h(z) = \frac{k(z, z_1)}{\sum_{j=1}^N k(z, z_j)} x_{z,1} + \frac{k(z, z_2)}{\sum_{j=1}^N k(z, z_j)} x_{z,2} + \cdots + \frac{k(z, z_N)}{\sum_{j=1}^N k(z, z_j)} x_{z,N},$$

$$\lambda_i = \frac{k(z, z_i)}{\sum_{j=1}^N k(z, z_j)},$$

$$\dot{x}_0 = \lambda \times x_z$$

## Regression Formulation

## Kernel Regression

$$h(z) = \frac{k(z, z_1)}{\sum_{j=1}^N k(z, z_j)} x_{z,1} + \frac{k(z, z_2)}{\sum_{j=1}^N k(z, z_j)} x_{z,2} + \cdots + \frac{k(z, z_N)}{\sum_{j=1}^N k(z, z_j)} x_{z,N},$$

$$\lambda_i = \frac{k(z, z_i)}{\sum_{j=1}^N k(z, z_j)},$$

$$\dot{x}_0 = \underbrace{\lambda}_{\text{Known}} \times x_z$$

## Regression Formulation


## Kernel Regression

$$h(z) = \frac{k(z, z_1)}{\sum_{j=1}^N k(z, z_j)} x_{z,1} + \frac{k(z, z_2)}{\sum_{j=1}^N k(z, z_j)} x_{z,2} + \cdots + \frac{k(z, z_N)}{\sum_{j=1}^N k(z, z_j)} x_{z,N},$$


$$\lambda_i = \frac{k(z, z_i)}{\sum_{j=1}^N k(z, z_j)},$$

$$\dot{x}_0 = \underbrace{\lambda}_{\text{Known}} \times \overbrace{x_z}^{\text{Unknown}}$$

## Kernel Regression: Derivation


Estimating  $x_z$ : Hypotheses

## Kernel Regression: Derivation

Estimating  $x_z$ : Hypotheses

- Regression is performed for the initial time step only.

## Estimating $x_z$ : Hypotheses



- Regression is performed for the initial time step only.
- No correlation between  $x_z$  and the components of  $x^{Ki}$ .

## Estimating $x_z$ : Sequential Method

The framework start with structural element  $e_1^1$  with:

## Estimating $\mathbf{x}_z$ : Sequential Method

The framework start with structural element  $e_1^1$  with:

$$\mathbf{x}_{t,k}^j = \left[ x_{t,k}^j \dot{x}_{t,k}^j \ddot{x}_{t,k}^j \mathbf{x}_z \right]^T, \quad \mathbf{x}_z = \mathbf{0}.$$

## Estimating $x_z$ : Sequential Method

The framework starts with structural element  $e_1^1$  with:

$$\mathbf{x}_{t,k}^j = \left[ x_{t,k}^j \dot{x}_{t,k}^j \ddot{x}_{t,k}^j \mathbf{x}_z \right]^T, \quad \mathbf{x}_z = \mathbf{0}.$$

The prediction for the speed at time ( $t = 0$ ),

## Estimating $\mathbf{x}_z$ : Sequential Method

The framework starts with structural element  $e_1^1$  with:

$$\mathbf{x}_{t,k}^j = \begin{bmatrix} x_{t,k}^j & \dot{x}_{t,k}^j & \ddot{x}_{t,k}^j & \mathbf{x}_z \end{bmatrix}^\top, \quad \mathbf{x}_z = \mathbf{0}.$$

The prediction for the speed at time ( $t = 0$ ),

$$\mathbf{x}_{0|z,k}^j = \mathbf{A}^{\text{Kr}} \mathbf{x}_{t,k}^j + \mathbf{w}^{\text{Kr}} : \mathbf{W}^{\text{Kr}} \sim \mathcal{N}(\mathbf{w}^{\text{Kr}}; \mathbf{0}, \mathbf{Q}^{\text{Kr}}),$$

## Estimating $\mathbf{x}_z$ : Sequential Method

The framework starts with structural element  $e_1^1$  with:

$$\mathbf{x}_{t,k}^j = \begin{bmatrix} x_{t,k}^j & \dot{x}_{t,k}^j & \ddot{x}_{t,k}^j & \mathbf{x}_z \end{bmatrix}^\top, \quad \mathbf{x}_z = \mathbf{0}.$$

The prediction for the speed at time ( $t = 0$ ),

$$\mathbf{x}_{0|z,k}^j = \mathbf{A}^{\text{Kr}} \mathbf{x}_{t,k}^j + \mathbf{w}^{\text{Kr}} : \mathbf{W}^{\text{Kr}} \sim \mathcal{N}(\mathbf{w}^{\text{Kr}}; \mathbf{0}, \mathbf{Q}^{\text{Kr}}),$$

where:

## Estimating $\mathbf{x}_z$ : Sequential Method

The framework starts with structural element  $e_1^1$  with:

$$\mathbf{x}_{t,k}^j = \begin{bmatrix} x_{t,k}^j & \dot{x}_{t,k}^j & \ddot{x}_{t,k}^j & \mathbf{x}_z \end{bmatrix}^\top, \quad \mathbf{x}_z = \mathbf{0}.$$

The prediction for the speed at time ( $t = 0$ ),

$$\mathbf{x}_{0|z,k}^j = \mathbf{A}^{\text{Kr}} \mathbf{x}_{t,k}^j + \mathbf{w}^{\text{Kr}} : \mathbf{W}^{\text{Kr}} \sim \mathcal{N}(\mathbf{w}^{\text{Kr}}; \mathbf{0}, \mathbf{Q}^{\text{Kr}}),$$

where:

$$\mathbf{A}^{\text{Kr}} = \begin{bmatrix} 1 & 0 & 0 & \mathbf{0}_{1 \times N} \\ 0 & 0 & 0 & \lambda_{1 \times N} \\ 0 & 0 & 1 & \mathbf{0}_{1 \times N} \\ 0 & 0 & 0 & \mathbf{I}_{N \times N} \end{bmatrix}, \quad \mathbf{Q}^{\text{Kr}} = \text{Block diag} (\mathbf{0}, (\sigma_{W,0}^1)^2 \times \mathbf{I}_{N \times N})$$

The transition model for time steps  $t = 0$  to  $T$ ,

$$\mathbf{x}_{t,z|(t-1,z),k}^j = \mathbf{A}^{\text{Se}} \mathbf{x}_{(t-1,z)|((t-1,z),k)}^j + \mathbf{w} : \mathbf{W} \sim \mathcal{N}(\mathbf{w}; \mathbf{0}, \mathbf{Q}).$$

where,

$$\mathbf{A}^{\text{Se}} = \text{Block diag} (\mathbf{A}^{\text{Ki}}, \mathbf{I}_{\mathbb{N} \times \mathbb{N}}),$$

$$\mathbf{Q} = \text{Block diag} (\mathbf{Q}^{\text{Ki}}, \mathbf{0}_{\mathbb{N} \times \mathbb{N}}),$$

$\mathbf{A}^{\text{Ki}}$  is defined by,

$$\mathbf{A}^{\text{Ki}} = \begin{bmatrix} 1 & \Delta t & \frac{\Delta t^2}{2} \\ 0 & 1 & \Delta t \\ 0 & 0 & 1 \end{bmatrix}.$$

## Kernel Regression: Derivation

The Kalman filter and Kalman smoother are employed.

## Kernel Regression: Derivation

The Kalman filter and Kalman smoother are employed. The Kalman smoother is utilized to update  $x_z$ :

The Kalman filter and Kalman smoother are employed. The Kalman smoother is utilized to update  $\mathbf{x}_z$ :

$$\mathbf{J}_{\mathbf{z},k}^j = \Sigma_{\mathbf{z},k}^j (\mathbf{A}^{\text{Kr}})^T \left( \Sigma_{0|\mathbf{z},k}^j \right)^{-1},$$

## Kernel Regression: Derivation

The Kalman filter and Kalman smoother are employed. The Kalman smoother is utilized to update  $\mathbf{x}_z$ :

$$\begin{aligned}\mathbf{J}_{\mathbf{z},k}^j &= \boldsymbol{\Sigma}_{\mathbf{z},k}^j (\mathbf{A}^{\text{Kr}})^T \left( \boldsymbol{\Sigma}_{0|\mathbf{z},k}^j \right)^{-1}, \\ \boldsymbol{\mu}_{\mathbf{z}|T,k}^j &= \boldsymbol{\mu}_{\mathbf{z},k}^j + \mathbf{J}_{\mathbf{z},k}^j \left( \boldsymbol{\mu}_{0|T,k}^j - \boldsymbol{\mu}_{0|\mathbf{z},k}^j \right),\end{aligned}$$

## Kernel Regression: Derivation

The Kalman filter and Kalman smoother are employed. The Kalman smoother is utilized to update  $\mathbf{x}_z$ :

$$\begin{aligned}\mathbf{J}_{\mathbf{z},k}^j &= \Sigma_{\mathbf{z},k}^j (\mathbf{A}^{K_r})^\top \left( \Sigma_{0|\mathbf{z},k}^j \right)^{-1}, \\ \boldsymbol{\mu}_{\mathbf{z}|T,k}^j &= \boldsymbol{\mu}_{\mathbf{z},k}^j + \mathbf{J}_{\mathbf{z},k}^j \left( \boldsymbol{\mu}_{0|T,k}^j - \boldsymbol{\mu}_{0|\mathbf{z},k}^j \right), \\ \Sigma_{\mathbf{z}|T,k}^j &= \Sigma_{\mathbf{z},k}^j + \mathbf{J}_{\mathbf{z},k}^j \left( \Sigma_{0|T,k}^j - \Sigma_{0|\mathbf{z},k}^j \right) \left( \mathbf{J}_{\mathbf{z},k}^j \right)^\top.\end{aligned}$$

The Kalman filter and Kalman smoother are employed. The Kalman smoother is utilized to update  $\mathbf{x}_z$ :

$$\begin{aligned}\mathbf{J}_{\mathbf{z},k}^j &= \Sigma_{\mathbf{z},k}^j (\mathbf{A}^{K_r})^\top \left( \Sigma_{0|\mathbf{z},k}^j \right)^{-1}, \\ \boldsymbol{\mu}_{\mathbf{z}|T,k}^j &= \boldsymbol{\mu}_{\mathbf{z},k}^j + \mathbf{J}_{\mathbf{z},k}^j \left( \boldsymbol{\mu}_{0|T,k}^j - \boldsymbol{\mu}_{0|\mathbf{z},k}^j \right), \\ \Sigma_{\mathbf{z}|T,k}^j &= \Sigma_{\mathbf{z},k}^j + \mathbf{J}_{\mathbf{z},k}^j \left( \Sigma_{0|T,k}^j - \Sigma_{0|\mathbf{z},k}^j \right) \left( \mathbf{J}_{\mathbf{z},k}^j \right)^\top.\end{aligned}$$

The obtained  $\mathbf{x}_z$  is then utilized as an initial  $\mathbf{x}_z$  for the next structural element.

The Kalman filter and Kalman smoother are employed. The Kalman smoother is utilized to update  $\mathbf{x}_z$ :

$$\begin{aligned}\mathbf{J}_{\mathbf{z},k}^j &= \Sigma_{\mathbf{z},k}^j (\mathbf{A}^{K_r})^\top \left( \Sigma_{0|\mathbf{z},k}^j \right)^{-1}, \\ \boldsymbol{\mu}_{\mathbf{z}|T,k}^j &= \boldsymbol{\mu}_{\mathbf{z},k}^j + \mathbf{J}_{\mathbf{z},k}^j \left( \boldsymbol{\mu}_{0|T,k}^j - \boldsymbol{\mu}_{0|\mathbf{z},k}^j \right), \\ \Sigma_{\mathbf{z}|T,k}^j &= \Sigma_{\mathbf{z},k}^j + \mathbf{J}_{\mathbf{z},k}^j \left( \Sigma_{0|T,k}^j - \Sigma_{0|\mathbf{z},k}^j \right) \left( \mathbf{J}_{\mathbf{z},k}^j \right)^\top.\end{aligned}$$

The obtained  $\mathbf{x}_z$  is then utilized as an initial  $\mathbf{x}_z$  for the next structural element.

**Problem:** it takes ver long time.

The Kalman filter and Kalman smoother are employed. The Kalman smoother is utilized to update  $\mathbf{x}_z$ :

$$\begin{aligned}\mathbf{J}_{\mathbf{z},k}^j &= \Sigma_{\mathbf{z},k}^j (\mathbf{A}^{K_r})^\top \left( \Sigma_{0|\mathbf{z},k}^j \right)^{-1}, \\ \boldsymbol{\mu}_{\mathbf{z}|T,k}^j &= \boldsymbol{\mu}_{\mathbf{z},k}^j + \mathbf{J}_{\mathbf{z},k}^j \left( \boldsymbol{\mu}_{0|T,k}^j - \boldsymbol{\mu}_{0|\mathbf{z},k}^j \right), \\ \Sigma_{\mathbf{z}|T,k}^j &= \Sigma_{\mathbf{z},k}^j + \mathbf{J}_{\mathbf{z},k}^j \left( \Sigma_{0|T,k}^j - \Sigma_{0|\mathbf{z},k}^j \right) \left( \mathbf{J}_{\mathbf{z},k}^j \right)^\top.\end{aligned}$$

The obtained  $\mathbf{x}_z$  is then utilized as an initial  $\mathbf{x}_z$  for the next structural element.

**Problem:** it takes ver long time.  
**(too many structural elements).**

# Estimating $x_z$ : Batch Method

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_0|_z = \lambda_k \times x_z$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_0|z = \lambda_k \times x_z$$

$$\text{KalmanFilter}(\mu_{t-1|(t-1,z)}, \Sigma_{t-1|(t-1,z)}, y_t, A_t, Q_t, C_t, R_t),$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_0|z = \lambda_k \times x_z$$

KalmanFilter( $\mu_{t-1|(t-1,z)}$ ,  $\Sigma_{t-1|(t-1,z)}$ ,  $y_t$ ,  $A_t$ ,  $Q_t$ ,  $C_t$ ,  $R_t$ ),  
KalmanSmoothen( $\mu_{T|(T-1,z)}$ ,  $\Sigma_{T|(T-1,z)}$ ,  $A_t$ ),

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

KalmanFilter( $\mu_{t-1|(t-1,z)}$ ,  $\Sigma_{t-1|(t-1,z)}$ ,  $y_t$ ,  $A_t$ ,  $Q_t$ ,  $C_t$ ,  $R_t$ ),

KalmanSmoothen( $\mu_{T|(T-1,z)}$ ,  $\Sigma_{T|(T-1,z)}$ ,  $A_t$ ),

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1},$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

KalmanFilter( $\mu_{t-1|(t-1,z)}$ ,  $\Sigma_{t-1|(t-1,z)}$ ,  $y_t$ ,  $A_t$ ,  $Q_t$ ,  $C_t$ ,  $R_t$ ),

KalmanSmoothen( $\mu_{T|(T-1,z)}$ ,  $\Sigma_{T|(T-1,z)}$ ,  $A_t$ ),

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1}, \dot{\Sigma}_{0|T,z} = \text{Block diag} \left( (\dot{\sigma}_{0|z,1}^1)^2, \dots, (\dot{\sigma}_{0|z,E}^B)^2 \right),$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

KalmanFilter( $\mu_{t-1|(t-1,z)}$ ,  $\Sigma_{t-1|(t-1,z)}$ ,  $y_t$ ,  $A_t$ ,  $Q_t$ ,  $C_t$ ,  $R_t$ ),

KalmanSmoothen( $\mu_{T|(T-1,z)}$ ,  $\Sigma_{T|(T-1,z)}$ ,  $A_t$ ),

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1}, \dot{\Sigma}_{0|T,z} = \text{Block diag} \left( (\dot{\sigma}_{0|z,1}^1)^2, \dots, (\dot{\sigma}_{0|z,E}^B)^2 \right),$$
$$A^{Ba} = [\lambda_1 \dots \lambda_E]^T \in \mathbb{R}^{E \times N},$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

KalmanFilter( $\mu_{t-1|(t-1,z)}$ ,  $\Sigma_{t-1|(t-1,z)}$ ,  $y_t$ ,  $A_t$ ,  $Q_t$ ,  $C_t$ ,  $R_t$ ),

KalmanSmoothen( $\mu_{T|(T-1,z)}$ ,  $\Sigma_{T|(T-1,z)}$ ,  $A_t$ ),

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1}, \dot{\Sigma}_{0|T,z} = \text{Block diag} \left( (\dot{\sigma}_{0|z,1}^1)^2, \dots, (\dot{\sigma}_{0|z,E}^B)^2 \right),$$

$$A^{Ba} = [\lambda_1 \dots \lambda_E]^T \in \mathbb{R}^{E \times N},$$

$$\dot{\mu}_{0|z} = A^{Ba} \times \mu_z,$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

KalmanFilter( $\mu_{t-1|(t-1,z)}, \Sigma_{t-1|(t-1,z)}, y_t, A_t, Q_t, C_t, R_t$ ),

KalmanSmoothen( $\mu_{T|(T-1,z)}, \Sigma_{T|(T-1,z)}, A_t$ ),

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1}, \dot{\Sigma}_{0|T,z} = \text{Block diag} \left( (\dot{\sigma}_{0|z,1}^1)^2, \dots, (\dot{\sigma}_{0|z,E}^B)^2 \right),$$

$$A^{Ba} = [\lambda_1 \dots \lambda_E]^T \in \mathbb{R}^{E \times N},$$

$$\dot{\mu}_{0|z} = A^{Ba} \times \mu_z, \dot{\Sigma}_{0|z} = A^{Ba} \times \Sigma_z \times (A^{Ba})^T$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

KalmanFilter( $\mu_{t-1|(t-1,z)}$ ,  $\Sigma_{t-1|(t-1,z)}$ ,  $y_t$ ,  $A_t$ ,  $Q_t$ ,  $C_t$ ,  $R_t$ ),

KalmanSmoothen( $\mu_{T|(T-1,z)}$ ,  $\Sigma_{T|(T-1,z)}$ ,  $A_t$ ),

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1}, \dot{\Sigma}_{0|T,z} = \text{Block diag} \left( (\dot{\sigma}_{0|z,1}^1)^2, \dots, (\dot{\sigma}_{0|z,E}^B)^2 \right),$$

$$A^{Ba} = [\lambda_1 \dots \lambda_E]^T \in \mathbb{R}^{E \times N},$$

$$\dot{\mu}_{0|z} = A^{Ba} \times \mu_z, \dot{\Sigma}_{0|z} = A^{Ba} \times \Sigma_z \times (A^{Ba})^T + 10^{-6} \times I$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

$\text{KalmanFilter}(\mu_{t-1|(t-1,z)}, \Sigma_{t-1|(t-1,z)}, y_t, A_t, Q_t, C_t, R_t)$ ,

$\text{KalmanSmoothen}(\mu_{T|(T-1,z)}, \Sigma_{T|(T-1,z)}, A_t)$ ,

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1}, \dot{\Sigma}_{0|T,z} = \text{Block diag} \left( (\dot{\sigma}_{0|z,1}^1)^2, \dots, (\dot{\sigma}_{0|z,E}^B)^2 \right),$$

$$A^{Ba} = [\lambda_1 \dots \lambda_E]^T \in \mathbb{R}^{E \times N},$$

$$\dot{\mu}_{0|z} = A^{Ba} \times \mu_z, \dot{\Sigma}_{0|z} = A^{Ba} \times \Sigma_z \times (A^{Ba})^T + 10^{-6} \times I$$

$$J_z = \Sigma_z (A^{Ba})^T \dot{\Sigma}_{0|z}^{-1},$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

$\text{KalmanFilter}(\mu_{t-1|(t-1,z)}, \Sigma_{t-1|(t-1,z)}, y_t, A_t, Q_t, C_t, R_t),$

$\text{KalmanSmoothen}(\mu_{T|(T-1,z)}, \Sigma_{T|(T-1,z)}, A_t),$

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1}, \dot{\Sigma}_{0|T,z} = \text{Block diag} \left( (\dot{\sigma}_{0|z,1}^1)^2, \dots, (\dot{\sigma}_{0|z,E}^B)^2 \right),$$

$$A^{Ba} = [\lambda_1 \dots \lambda_E]^T \in \mathbb{R}^{E \times N},$$

$$\dot{\mu}_{0|z} = A^{Ba} \times \mu_z, \dot{\Sigma}_{0|z} = A^{Ba} \times \Sigma_z \times (A^{Ba})^T + 10^{-6} \times I$$

$$J_z = \Sigma_z (A^{Ba})^T \dot{\Sigma}_{0|z}^{-1},$$

$$\mu_{z|T} = \mu_z + J_z (\dot{\mu}_{0|T,z} - \dot{\mu}_{0|z}),$$

# Estimating $x_z$ : Batch Method

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

$\text{KalmanFilter}(\mu_{t-1|(t-1,z)}, \Sigma_{t-1|(t-1,z)}, y_t, A_t, Q_t, C_t, R_t),$   
 $\text{KalmanSmoothen}(\mu_{T|(T-1,z)}, \Sigma_{T|(T-1,z)}, A_t),$

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1}, \dot{\Sigma}_{0|T,z} = \text{Block diag} \left( (\dot{\sigma}_{0|z,1}^1)^2, \dots, (\dot{\sigma}_{0|z,E}^B)^2 \right),$$

$$A^{Ba} = [\lambda_1 \dots \lambda_E]^T \in \mathbb{R}^{E \times N},$$

$$\dot{\mu}_{0|z} = A^{Ba} \times \mu_z, \dot{\Sigma}_{0|z} = A^{Ba} \times \Sigma_z \times (A^{Ba})^T + 10^{-6} \times I$$

$$J_z = \Sigma_z (A^{Ba})^T \dot{\Sigma}_{0|z}^{-1},$$

$$\mu_{z|T} = \mu_z + J_z (\dot{\mu}_{0|T,z} - \dot{\mu}_{0|z}),$$

$$\Sigma_{z|T} = \Sigma_z + J_z (\dot{\Sigma}_{0|T,z} - \dot{\Sigma}_{0|z}) J_z^T.$$

# Estimating $x_z$ : Batch Method

## Loop

For each element  $e_k \in E$ :

$$\dot{x}_{0|z} = \lambda_k \times x_z$$

$\text{KalmanFilter}(\mu_{t-1|(t-1,z)}, \Sigma_{t-1|(t-1,z)}, y_t, A_t, Q_t, C_t, R_t),$

$\text{KalmanSmoothen}(\mu_{T|(T-1,z)}, \Sigma_{T|(T-1,z)}, A_t),$

$$\dot{\mu}_{0|T,z} \in \mathbb{R}^{E \times 1}, \dot{\Sigma}_{0|T,z} = \text{Block diag} \left( (\dot{\sigma}_{0|z,1}^1)^2, \dots, (\dot{\sigma}_{0|z,E}^B)^2 \right),$$

$$A^{Ba} = [\lambda_1 \dots \lambda_E]^T \in \mathbb{R}^{E \times N},$$

$$\dot{\mu}_{0|z} = A^{Ba} \times \mu_z, \dot{\Sigma}_{0|z} = A^{Ba} \times \Sigma_z \times (A^{Ba})^T + 10^{-6} \times I$$

$$J_z = \Sigma_z (A^{Ba})^T \dot{\Sigma}_{0|z}^{-1},$$

$$\mu_{z|T} = \mu_z + J_z (\dot{\mu}_{0|T,z} - \dot{\mu}_{0|z}),$$

$$\Sigma_{z|T} = \Sigma_z + J_z (\dot{\Sigma}_{0|T,z} - \dot{\Sigma}_{0|z}) J_z^T.$$

# Computational Challenges

- $\dot{\Sigma}_{0|T,z}$  is diagonal matrix → assumes no correlations.

# Computational Challenges

- $\dot{\Sigma}_{0|T,z}$  is diagonal matrix  $\rightarrow$  assumes no correlations.
- The response  $\dot{\mu}_{0|T,z}$  estimate is not good initially, ( $I \rightarrow$  large).

# Computational Challenges

- $\dot{\Sigma}_{0|T,z}$  is diagonal matrix → assumes no correlations.
- The response  $\dot{\mu}_{0|T,z}$  estimate is not good initially, ( $I \rightarrow \text{large}$ ).
- If  $I$  is large

# Computational Challenges

- $\dot{\Sigma}_{0|T,z}$  is diagonal matrix  $\rightarrow$  assumes no correlations.
- The response  $\dot{\mu}_{0|T,z}$  estimate is not good initially, ( $I \rightarrow$  large).
- If  $I$  is large  $\rightarrow \lambda_1 \approx \lambda_2 \dots \approx \lambda_E$

# Computational Challenges

- $\dot{\Sigma}_{0|T,z}$  is diagonal matrix  $\rightarrow$  assumes no correlations.
- The response  $\dot{\mu}_{0|T,z}$  estimate is not good initially, ( $I \rightarrow$  large).
- If  $I$  is large  $\rightarrow \lambda_1 \approx \lambda_2 \dots \approx \lambda_E$   
 $\rightarrow \dot{\Sigma}_{0|z}$  has high  $\rho$

# Computational Challenges

- $\dot{\Sigma}_{0|T,z}$  is diagonal matrix  $\rightarrow$  assumes no correlations.
- The response  $\dot{\mu}_{0|T,z}$  estimate is not good initially, ( $I \rightarrow$  large).
- If  $I$  is large  $\rightarrow \lambda_1 \approx \lambda_2 \cdots \approx \lambda_E$   
 $\rightarrow \dot{\Sigma}_{0|z}$  has high  $\rho \rightarrow \dot{\Sigma}_{0|z}$  is almost singular

# Computational Challenges

- $\dot{\Sigma}_{0|T,z}$  is diagonal matrix  $\rightarrow$  assumes no correlations.
- The response  $\dot{\mu}_{0|T,z}$  estimate is not good initially, ( $I \rightarrow$  large).
- If  $I$  is large  $\rightarrow \lambda_1 \approx \lambda_2 \cdots \approx \lambda_E$   
 $\rightarrow \dot{\Sigma}_{0|z}$  has high  $\rho \rightarrow \dot{\Sigma}_{0|z}$  is almost singular  
 $\rightarrow \dot{\Sigma}_{0|z}^{-1}$  high values

# Computational Challenges

- $\dot{\Sigma}_{0|T,z}$  is diagonal matrix  $\rightarrow$  assumes no correlations.
- The response  $\dot{\mu}_{0|T,z}$  estimate is not good initially, ( $I \rightarrow$  large).
- If  $I$  is large  $\rightarrow \lambda_1 \approx \lambda_2 \cdots \approx \lambda_E$   
 $\rightarrow \dot{\Sigma}_{0|z}$  has high  $\rho \rightarrow \dot{\Sigma}_{0|z}$  is almost singular  
 $\rightarrow \dot{\Sigma}_{0|z}^{-1}$  high values  $\rightarrow J_z$  high values

# Computational Challenges

- $\dot{\Sigma}_{0|T,z}$  is diagonal matrix  $\rightarrow$  assumes no correlations.
- The response  $\dot{\mu}_{0|T,z}$  estimate is not good initially, ( $I \rightarrow$  large).
- If  $I$  is large  $\rightarrow \lambda_1 \approx \lambda_2 \cdots \approx \lambda_E$ 
  - $\rightarrow \dot{\Sigma}_{0|z}$  has high  $\rho \rightarrow \dot{\Sigma}_{0|z}$  is almost singular
  - $\rightarrow \dot{\Sigma}_{0|z}^{-1}$  high values  $\rightarrow J_z$  high values
  - $\rightarrow \mu_{z|T}$  high values.

Verification with Synthetic Data

# Synthetic Structural Attribute


# Synthetic Structural Attribute

$$z^1 = \exp(\dot{x})$$

## Verification with Synthetic Data

# Synthetic Structural Attribute



$$z^1 = \exp(\dot{x})$$



## Verification with Synthetic Data

# Synthetic Structural Attribute

$$z^1 = \exp(\dot{x})$$





## Verification with Synthetic Data

## Synthetic Structural Attribute

$$z^1 = \exp(\dot{x})$$


$$z^2 = \log(|\dot{x}|)$$





## Verification with Synthetic Data

## Synthetic Structural Attribute

$$z^1 = \exp(\dot{x})$$





$$z^2 = \log(|\dot{x}|)$$





## Verification with Synthetic Data

## Synthetic Structural Attribute

$$z^1 = \exp(\dot{x})$$




$$z^2 = \log(|\dot{x}|)$$





## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^1$


## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^1$ 



## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^1$ 

## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^1$ 


## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^1$ 



## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^2$


## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^2$ 



## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^2$ 

## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^2$ 

## Verification with Synthetic Data

Estimating  $\dot{\mu}_0$  from  $z^2$ 

Next Steps

## Next:

1. Assess the performance of Kernel Regression with noisy functions.

Next Steps

## Next:

1. Assess the performance of Kernel Regression with noisy functions.
2. Full integration within the original framework.

Next Steps

## Next:

1. Assess the performance of Kernel Regression with noisy functions.
2. Full integration within the original framework.
3. Perform analyses on Real Data.