

Lecture 27: Intro to Differential Equations and IVPs

Goal: Be able to solve (very basic) Initial Value Problems through integration.

For $y=f(x)$, a differential equation (diff.eq) is an equation involving x, y , and the derivatives of y .

E.g., $\frac{d^2 y}{dx^2} = -9.81$ [Gravity]

$$\frac{dy}{dx} = y \quad [\text{Exponential Growth}]$$

$$\frac{d^2 y}{dx^2} + y = \cos x \quad [\text{Damped Harmonic Motion}]$$

Ex1/ Verify $y = 3e^{2x}$ is a solution to the diff eq.

$$\frac{dy}{dx} = 2y$$

$$\frac{d}{dx}(3e^{2x}) = 2 \cdot 3e^{2x} = 6e^{2x}$$

$$2(3e^{2x}) = 6e^{2x} //$$

Ex2/ Solve the diff. eq. $y' = \frac{1}{x}$

$$y' = \frac{1}{x}$$


$$\int y' dx = \int \frac{1}{x} dx$$

$$y = \ln|x| + C$$

General Sol.
to
 $y' = \frac{1}{x}$

A function describing all possible solutions is called the general solution to the diff eq

Ex3/ Find the general sol. to $y' = 3x^2$



Q: Is there a way to have a unique (only 1) Solution?

A: Say y also needs to pass through a point (e.g. $y(0) = 0$). This is called an initial condition.

★ Usually we specify when $x=0$, but that's not always.

IVPs

An initial value problem (IVP) is a diff eq combined with initial conditions

Ex 5/ The system

$$\begin{cases} y'(x) = \sqrt{x} \\ y(0) = 1 \end{cases}$$

is an IVP. How can we solve?

① Find the general solution

$$y'(x) = \sqrt{x}$$

$$\int y'(x) dx = \int \sqrt{x} dx$$

$$y(x) = \int x^{\frac{1}{2}} dx$$

$$y(x) = \frac{x^{\frac{1}{2}+1}}{\frac{1}{2}+1} + C$$

$$y(x) = \frac{2}{3} x^{\frac{3}{2}} + C$$

② Plug in IC, solve for C $y(0) = 1$

$$1 = y(0) = \frac{2}{3} (0)^{\frac{3}{2}} + C$$

$$1 = 0 + C$$

$$C = 1$$

Conclusion: $y(x) = \frac{2}{3} x^{\frac{3}{2}} + 1$ is the sol. to the IVP.

Def We say y is proportional to x if there is a constant k s.t. $y = kx$

Exb The size of a population is governed by $\frac{dP}{dt} = \sqrt[3]{t} \leftarrow \frac{dP}{dt}$ is proportional to $\sqrt[3]{t}$ where $k=1$

Where P is the population after t days. If the initial population is 1000, find the pop. size after 8 days.

I.e., solve

$$\begin{cases} \frac{dP}{dt} = \sqrt[3]{t} \\ P(0) = 1000 \end{cases}$$

$$\frac{dP}{dt} = \sqrt[3]{t}$$

$$\int \frac{dP}{dt} dt = \int \sqrt[3]{t} dt$$

$$P(t) = \int t^{\frac{1}{3}} dt$$

$$P(t) = \frac{t^{\frac{1}{3}+1}}{\frac{1}{3}+1} + C$$

$$P(0) = 1000$$

$$P(t) = \frac{3}{4} t^{\frac{4}{3}} + C$$

$$1000 = P(0) = 0 + C$$

$$C = 1000$$

$$P(t) = \frac{3}{4} t^{\frac{4}{3}} + 1000$$

Thus,

$$P(8) = \frac{3}{4} 8^{\frac{4}{3}} + 1000 = \frac{3}{4} (\sqrt[3]{8})^4 + 1000$$

$$= 12 + 1000 = 1012$$

Conclusion: The population size is 1012 after 8 days.

Multiple ICs

Fact: To have a unique solution, you need N ICs where N is the highest order of derivatives in the diff eq.

Ex7/ Solve the IVP

$$\begin{cases} y''(t) = t-3 \\ y'(1) = \frac{1}{2} \\ y(0) = 1 \end{cases}$$



$$y''(t) = t-3$$

$$\int y''(t) dt = \int (t-3) dt$$

$$y'(t) = \frac{t^{1+1}}{1+1} - 3t + C$$

$$y'(t) = \frac{1}{2}t^2 - 3t + C$$

$$\frac{1}{2} = y'(1) = \frac{1}{2} - 3 + C$$

$$\frac{1}{2} = \frac{1}{2} - 3 + C$$

$$0 = -3 + C$$

$$\rightarrow C = 3$$

$$y'(t) = \frac{1}{2}t^2 - 3t + 3$$

$$\int y'(t) dt = \int \left(\frac{1}{2}t^2 - 3t + 3 \right) dt$$

$$y(t) = \frac{1}{2} \cdot \frac{t^{2+1}}{2+1} - 3 \cdot \frac{t^{1+1}}{1+1} + 3t + D$$

$$y(t) = \frac{1}{6}t^3 - \frac{3}{2}t^2 + 3t + D$$

$$1 = y(0) = 0 + D$$

$$D = 1$$

Thus, the solution is $y(t) = \frac{1}{6}t^3 - \frac{3}{2}t^2 + 3t + 1$

NOTE: You can always check your answers by differentiating

$$y(0) = 0 - 0 + 0 + 1 = 1 \quad \checkmark$$

$$y'(t) = \frac{1}{2}t^2 - 3t + 3$$

$$y'(1) = \frac{1}{2} - 3 + 3 = \frac{1}{2} \quad \checkmark$$

$$y''(t) = t - 3 \quad \checkmark$$

Ex 8/ A ball is kicked from the ground with an initial velocity of $20 \frac{m}{s}$. Describe the vertical height of the ball as a function of time (assume gravity is the only force acting on the ball)

Solve the system

$$\begin{array}{l} s'' \rightarrow a(t) = -9.81 \\ s' \rightarrow v(0) = 20 \\ s \rightarrow s(0) = 0 \end{array}$$

Accel. \xrightarrow{s} velocity $\xrightarrow{\int}$ pos.

$$a(t) = -9.81$$

$$\int a(t) dt = \int (-9.81) dt$$

$$v(t) = -9.81t + C$$

$$v(0) = 20$$

$$20 = v(0) = -9.81(0) + C$$

$$C = 20$$

$$v(t) = -9.81t + 20$$

$$\int v(t) dt = \int (-9.81t + 20) dt$$

$$s(t) = -9.81\left(\frac{1}{2}t^2\right) + 20t + D$$

$$s(t) = -4.905t^2 + 20t + D$$

$$\underline{s(0) = 0}$$

$$0 = s(0) = D$$

$$D = 0$$

Thus,

$$s(t) = -4.905t^2 + 20t$$

Ex 9 (Hw27, Q4) Solve the IVP

$$\begin{cases} y' = 3\cos x + 5 \end{cases}$$

$$\begin{cases} y\left(\frac{3\pi}{2}\right) = 4 \end{cases}$$

$$y' = 3\cos x + 5$$

$$\int y' dx = \int (3\cos x + 5) dx$$

$$y(x) = 3\sin x + 5x + C$$

$$y\left(\frac{3\pi}{2}\right) = 4$$



$$4 = y\left(\frac{3\pi}{2}\right) = 3\sin\left(\frac{3\pi}{2}\right) + 5\left(\frac{3\pi}{2}\right) + C$$

$$4 = 3(-1) + \frac{15\pi}{2} + C$$

$$4 = -3 + \frac{15\pi}{2} + C$$

$$7 = \frac{15\pi}{2} + C$$

$$C = 7 - \frac{15\pi}{2}$$

Thus, $y(x) = 3\sin x + 5x + 7 - \frac{15\pi}{2}$