

Problems for Day 2 (Lectures 16-28): 1st/2nd Derivative Test, Optimization, Riemann Sums

1. Let $f(x) = \frac{x^2+3}{x-1}$

(a) Locate all (if any) asymptotes.

$$\frac{x^2+3}{x-1}$$

VA's: $x-1 \stackrel{\text{set}}{=} 0$
 $\boxed{x=1}$

HA's: Occur when

deg top \leq deg bottom

2 ~~2~~ 1

$$\begin{array}{r} \textcircled{x+1} \\ x-1 \overline{) x^2+3} \\ \underline{-x^2-x} \\ x+3 \\ \underline{-x-1} \\ 4 \end{array}$$

Slant:

$$\boxed{y=x+1}$$

(b) Find all critical values.

$$f(x) = \frac{x^2+3}{x-1}$$

$$f'(x) = \frac{2x(x-1) - (x^2+3)(1)}{(x-1)^2} = \frac{x^2-2x+3}{(x-1)^2}$$

$$= \frac{(x+1)(x-3)}{(x-1)^2} \stackrel{\text{set}}{=} 0$$

$$(x+1)(x-3) = 0$$

$$x = -1, 3$$

$$\frac{(x+1)(x-3)}{(x-1)^2}$$

(c) When is f increasing? When is f decreasing?

(c) Which is y increasing? Which is y decreasing?

	\leftarrow					
		-1		1	3	
Test Point	-100		0		2	100
Sign of $(x-1)^2$	+		+		+	+
Sign of $x+1$	-		+		+	+
Sign of $x-3$	-		-		-	+
Sign of f'	+		-		-	+
Results of 1st Derivative Test	Inc	\nearrow	\searrow Dec	\searrow Dec	\searrow Dec	\nearrow Inc
Inc: $(-\infty, -1) \cup (3, \infty)$			Dec: $(-1, 1) \cup (1, 3)$			

(d) Use the 1st Derivative Test to determine the locations of any local maximums/minimums.



Local Max at $x = -1$

Local Min at $x = 3$

(e) Verify your answer using the 2nd derivative test.

$$f'(x) = \frac{x^2 - 2x + 3}{(x-1)^2}$$

$$f''(x) = \frac{(2x-2)(x-1)^2 - (x^2-2x+3)2(x-1)(1)}{(x-1)^4}$$

$$f''(x) = \frac{8(x-1)}{(x-1)^4} = \frac{8}{(x-1)^3}$$




$$f''(-1) = \frac{8}{(-1-1)^3} = -1 < 0 \Rightarrow \text{Local Max at } x = -1$$

$$f''(3) = \frac{8}{(2-1)^3} = 1 > 0 \Rightarrow \text{Local Min at } x = 3$$

(f) Find all inflection points (if any).

$$f''(x) = \frac{8}{(x-1)^3} \stackrel{\text{set}}{=} 0$$

$$\Rightarrow 8 = 0 \quad \text{No Solution!}$$



Still could change concavity at asymptote


(g) When is f concave up? When is f concave down?

$\frac{8}{(x-1)^3}$		
Test Point	0	100
Sign of f''	-	+
Results of CTest	CD	CU

$\begin{pmatrix} ++ \\ -- \end{pmatrix}$

CD on $(-\infty, 1)$
 CU on $(1, \infty)$

(h) Draw a graph of f .




2. Compute $\lim_{x \rightarrow \infty} \frac{2x^2 - 4}{5x^2 - 4x + 6}$

$$\lim_{x \rightarrow \infty} \frac{2x^2 - 4}{5x^2 - 4x + 6} = \lim_{x \rightarrow \infty} \frac{x^2 \left(2 - \frac{4}{x^2}\right)}{x^2 \left(5 - \frac{4}{x} + \frac{6}{x^2}\right)}$$

$$= \lim_{x \rightarrow \infty} \frac{2 - \frac{4}{x^2}}{5 - \frac{4}{x} + \frac{6}{x^2}} = \frac{2 - 0}{5 - 0 + 0} = \boxed{\frac{2}{5}}$$

3. (Exam 3, Problem 8) A family wants to fence a rectangular play area alongside the wall of their house. The wall of their house bounds one side of the play area. If they want the play area to be exactly 2500 ft², what is the least amount of fencing needed? Round your answer to the nearest tenth place.



Obj Minimize $P(x,y) = 2x + y$
 Given $xy = 2500$

$$y = \frac{2500}{x}$$

$$P(x) = 2x + \frac{2500}{x}$$

$$P'(x) = 2 - \frac{2500}{x^2} \stackrel{\text{Set}}{=} 0$$

$$2 = \frac{2500}{x^2}$$

$$\frac{x^2}{2500} = \frac{1}{2}$$

$$x^2 = 1250 ; x > 0$$

Potential loc for a min $\rightarrow x = \sqrt{1250} = \sqrt{5^4 \cdot 2} = 25\sqrt{2}$



$$y = \frac{2500}{x} \Rightarrow y = \frac{2500}{25\sqrt{2}} = \frac{100}{\sqrt{2}} = 50\sqrt{2}$$

$$P(25\sqrt{2}, 50\sqrt{2}) = 2(25\sqrt{2}) + 50\sqrt{2} = 100\sqrt{2} \approx 141.4 \text{ ft}$$

4. (Exam 3, Problem 9) A company's marketing department has determined that if their product is sold at the price of p dollars per unit, they can sell $q = 1200 - 100p$ units. Each unit costs 6 dollars to make. What price, p , should the company charge to maximize their profit?

$$\text{Profit} = \text{Revenue} - \text{Costs} = \left(\frac{\text{Price}}{\text{Item}} \right) \left(\frac{\# \text{ items}}{\text{Sold}} \right) - \left(\frac{\text{Cost}}{\text{item}} \right) \left(\frac{\# \text{ of items sold}}{\text{}} \right)$$

$$P_{(p,q)} = pq - 6q$$


$$\text{Obj: Max } P_{(p,q)} = pq - 6q$$

$$\text{Given } q = 1200 - 100p$$

$$\begin{aligned} P_{(p)} &= p(1200 - 100p) - 6(1200 - 100p) \\ &= 1200p - 100p^2 - 7200 + 600p \end{aligned}$$

$$P_{(p)} = -100p^2 + 1800p - 7200$$

$$\begin{aligned} \frac{dP}{dp} &= -200p + 1800 \stackrel{\text{set}}{=} 0 \\ -200p &= -1800 \\ p &= 9 \end{aligned}$$



The company should sell their products at \$9.

5. Solve the IVP:

$$\begin{cases} f'(t) = \sin t + 2t \\ f(0) = 5 \end{cases}$$

$$\frac{2t^{1+1}}{1+1}$$

$$f'(t) = \sin t + 2t$$

$$\int f'(t) dt = \int (\sin t + 2t) dt$$


$$f(t) = -\cos t + t^2 + C$$

$$\underline{f(0)=5:}$$

$$f(0) = -1 + 0 + C \stackrel{\text{set}}{=} 5$$

$$C = 6$$

6. Write the left and right Riemann Sums to estimate $\int_1^4 \sqrt{4x+1} \, dx$ when $N = 6$.



$$\Delta x = \frac{b-a}{N} = \frac{4-1}{6} = \frac{1}{2}$$

$$L_N = \sum_{i=0}^{N-1} f(a + i\Delta x) \Delta x$$

$$L_6 = \sum_{i=0}^5 \sqrt{4(1 + i(\frac{1}{2})) + 1} \left(\frac{1}{2}\right) = \frac{1}{2} \sum_{i=0}^5 \sqrt{4(1 + \frac{i}{2}) + 1}$$

$$= \frac{1}{2} \left[\underbrace{\sqrt{4(1)} + 1}_{i=0} + \underbrace{\sqrt{4(\frac{3}{2})} + 1}_{i=1} + \underbrace{\sqrt{4(2)} + 1}_{i=2} + \underbrace{\sqrt{4(\frac{5}{2})} + 1}_{i=3} + \underbrace{\sqrt{4(3)} + 1}_{i=4} + \underbrace{\sqrt{4(\frac{7}{2})} + 1}_{i=5} \right]$$

$$\approx 9.3385$$

$$R_6 = \frac{1}{2} \sum_{i=1}^6 \sqrt{4(1 + \frac{i}{2}) + 1}$$

$$= \frac{1}{2} \left[\underbrace{\sqrt{4(\frac{3}{2})} + 1}_{i=1} + \underbrace{\sqrt{4(2)} + 1}_{i=2} + \underbrace{\sqrt{4(\frac{5}{2})} + 1}_{i=3} + \underbrace{\sqrt{4(3)} + 1}_{i=4} + \underbrace{\sqrt{4(\frac{7}{2})} + 1}_{i=5} + \underbrace{\sqrt{4(4)} + 1}_{i=6} \right]$$

$$\approx 10.2820$$