

MA 16200: Plane Analytic Geometry and Calculus II

Lecture 4: Calculating Volumes & Solids of Revolution

Zachariah Pence

Purdue University

Sections Covered: 6.3

Derivation

General Slicing Method

Definition 1

Suppose a solid object extends from $x = a$ to $x = b$, and the cross-sectional area at a point x is given by a function $A(x)$ that can be integrated on $[a, b]$. Then the volume of the solid is:


$$V = \int_a^b A(x) \, dx$$

Example

Problem 2

Consider a solid whose base is the region in the first quadrant bounded by the curve $y = \sqrt{5-x}$ and the line $x = 3$, and whose cross sections through the solid perpendicular to the x -axis are squares.

- 1 Find an expression for the cross-sectional area $A(x)$ at a point $x \in [0, 3]$.
- 2 Find the volume of the solid.




Extra Space

Extra Space

Examples

Problem 3


Use the general slicing method to find the volume of the solid whose base is the triangle with vertices $(0,0)$, $(3,0)$, and $(0,2)$, and whose cross sections perpendicular to the base and parallel to the y-axis are semicircles



Extra Space

Extra Space

Solids of Revolution



Disk Method (about the x-axis)

Definition 4

Let $f \geq 0$ be continuous on $[a, b]$. If the region R bounded by the graph of f , the x -axis, and the lines $x = a$ and $x = b$ is revolved about the x -axis, the volume of the resulting solid of revolution is:

$$V = \int_a^b \pi [f(x)]^2 dx$$

Examples

Problem 5

Find the volume of the solid obtained by rotating the region bounded by the given curves about the specified line. Sketch the region and a typical disk.

1 $y = e^x; x = 1; y = 0$; about the x -axis.

2 $y = \frac{1}{x}; x = 1; x = 2; y = 0$; about the x -axis.

Extra Space

Extra Space

Extra Space

Extra Space

Washer Method (about the x-axis)

Definition 6

Let f and g be continuous function with $f(x) \geq g(x) \geq 0$ on $[a, b]$. Let R be the region bounded by $y = f(x)$, $y = g(x)$, $x = a$, and $x = b$. When R is revolved around the x-axis, the volume of the resulting solid of revolution is:

$$V = \int_a^b \pi[f(x)]^2 dx - \int_a^b \pi[g(x)]^2 dx = \int_a^b \pi([f(x)]^2 - [g(x)]^2) dx$$

Examples

Problem 7

Find the volume of the solid obtained by rotating the region bounded by the given curves about the specified line. Sketch the region and a typical washer.

- 1 $y = \sec x$; $y = 1$; $x = -1$; $x = 1$; *about the x -axis.*
- 2 $y = x^2$; $y^2 = x$; *about the x -axis.*

Extra Space

Extra Space

Extra Space

Extra Space

Rotating about the y-axis

Definition 8

Let p and q be continuous functions with $p(y) \geq q(y) \geq 0$ on $[c, d]$. Let R be the region bounded by $x = p(y)$, $x = q(y)$, and the lines $y = c$ and $y = d$. When R is revolved about the y-axis, the volume of the resulting solid of revolution is given by

$$V = \int_c^d \pi[p(y)^2 - q(y)^2] dy$$

If $q(y) = 0$, the disk method results:

$$V = \int_c^d \pi[p(y)]^2 dy$$

Examples

Problem 9

Find the volume of the solid obtained by rotating the region bounded by the given curves about the specified line. Sketch the region and a typical disk/washer.

- 1 $x = y - y^2$; $x = 0$; about the y -axis.
- 2 $y^2 = x$; $y = \frac{1}{2}x$; about the y -axis.

Extra Space

Extra Space

Extra Space

Extra Space

Examples

Problem 10

Find the volume of the solid obtained by rotating the region bounded by the given curves about the specified line. Sketch the region and a typical disk/washer.

- 1 $y = x; y = \sqrt{x}$ about $y = 1$.
- 2 $y = \ln x; x = 0$; on the interval $0 \leq y \leq 1$; about $x = -1$.

Extra Space

Extra Space

Extra Space


Extra Space

Volume of a Sphere

Problem 11

Show the volume of a sphere of radius R is:

$$V = \frac{4}{3}\pi R^3$$



Rotate $\sqrt{R^2 - x^2}$ about the x-axis, by the disk method:


$$\begin{aligned} V &= \int_{-R}^R \pi (\sqrt{R^2 - x^2})^2 dx = 2\pi \int_0^R (R^2 - x^2) dx \\ &= 2\pi \left[R^2 x - \frac{1}{3} x^3 \right]_0^R = 2\pi \left[\frac{2}{3} R^3 \right] = \frac{4}{3} \pi R^3 \end{aligned}$$

Volume of a Cone

Problem 12

Show the volume of a cone of base radius R and height h is:

$$V = \frac{h}{3}\pi R^2$$



Rotate $x = \frac{R}{h}y$ about the y -axis, by the disk method:

$$\begin{aligned} V &= \int_0^h \pi \left(\frac{R}{h}y \right)^2 dy = \pi \frac{R^2}{h^2} \left[\frac{1}{3}y^3 \right]_0^h \\ &= \pi \frac{R^2}{h^2} \left[\frac{1}{3}h^3 \right] = \frac{h}{3}\pi R^2 \end{aligned}$$

Volume of a Wine Barrel

Problem 13

The sides of a wine barrel can be approximated by the parabola:

$$y = R - cx^2; \quad -\frac{h}{2} \leq x \leq \frac{h}{2}$$


where R is the maximum radius, h is the height, and $c > 0$ a constant. Show that the volume is:

$$V = \frac{\pi h}{3} \left[2R^2 + r^2 - \frac{2}{5}(R - r)^2 \right]$$

where r is the minimum radius.

Finding the Volume

Rotate $y = R - cx^2$ about the x-axis. By the Disk Method:



$$\begin{aligned}
 V &= \int_{-h/2}^{h/2} \pi(R - cx^2)^2 dx = 2\pi \int_0^{h/2} (R - cx^2)^2 dx \\
 &= 2\pi \int_0^{h/2} (R^2 - 2Rcx^2 + c^2x^4) dx \\
 &= 2\pi \left[R^2x - \frac{2Rc}{3}x^3 + \frac{c^2}{5}x^5 \right] \\
 &= 2\pi \left[R^2(h/2) - \frac{2Rc}{3}(h/2)^3 + \frac{c^2}{5}(h/2)^5 \right] \\
 &= \pi R^2h - \frac{Rc\pi}{6}h^3 + \frac{c^2\pi}{80}h^5
 \end{aligned}$$

cont.

$$V = \pi h \left(R^2 - \frac{Rc}{6}h^2 + \frac{c^2}{80}h^4 \right) = \pi h \left(R^2 - \frac{2R}{3} \left(\frac{ch^2}{4} \right) + \frac{1}{5} \left(\frac{ch^2}{4} \right)^2 \right)$$

$$\text{Let } d = \frac{ch^2}{4} = R - r:$$

$$\begin{aligned} V &= \pi h \left(R^2 - \frac{2R}{3}d + \frac{1}{5}d^2 \right) = \frac{\pi h}{3} \left(3R^2 - 2Rd + \frac{3}{5}d^2 \right) \\ &= \frac{\pi h}{3} \left(2R^2 + R^2 + d^2 - \frac{2}{5}d^2 - 2Rd \right) \\ &= \frac{\pi h}{3} \left(2R^2 - \frac{2}{5}d^2 + R^2 - 2Rd + d^2 \right) = \frac{\pi h}{3} \left(2R^2 - \frac{2}{5}d^2 + (R - d)^2 \right) \\ &= \frac{\pi h}{3} \left(2R^2 + r^2 - \frac{2}{5}d^2 \right) = \frac{\pi h}{3} \left(2R^2 + r^2 - \frac{2}{5}(R - r)^2 \right) \end{aligned}$$