

MA 16200: Plane Analytic Geometry and Calculus II

Lecture 9: Intro to Trigonometric Integrals

Zachariah Pence

Purdue University

Sections Covered: 8.3

Basics

$$-\ln|\cos x| = \ln|(\cos x)^{-1}|$$

- $\int \sin x = -\cos x + C$
- $\int \cos x = \sin x + C$
- $\int \sec^2 x \, dx = \tan x + C$
- $\int \csc x \cot x \, dx = -\csc x + C$
- $\int \sec x \tan x \, dx = \sec x + C$
- $\int \csc^2 x \, dx = -\cot x + C$
- $\int \tan x = \ln|\sec x| + C$
- $\int \csc x = -\ln|\csc x + \cot x| + C$
- $\int \sec x = \ln|\sec x + \tan x| + C$
- $\int \cot x = \ln|\sin x| + C$

$$\begin{aligned} \int \frac{1}{u} &= \\ \int \tan x \, dx &= (-1) \int \frac{\sin x}{\cos x} \, dx \\ &\quad \underbrace{\qquad\qquad\qquad}_{u=\cos x} \\ &= -\ln|\cos x| + C_0 \\ &= \ln|\sec x| + C \\ \hline \int \cot x \, dx &= \int \frac{\cos x}{\sin x} \, dx \\ &\quad \underbrace{\qquad\qquad\qquad}_{u=\sin x} \\ &= \ln|\sin x| + C \end{aligned}$$

$$\begin{aligned}
 \int \csc x \, dx &= \int \csc x \left(\frac{\csc x + \cot x}{\csc x + \cot x} \right) dx \\
 &= (-1) \int \frac{(\csc^2 x + \cot x \csc x)}{\csc x + \cot x} dx \\
 &= -\ln |\csc x + \cot x| + C
 \end{aligned}$$

$$\begin{aligned}
 \int \sec x \, dx &= \int \sec x \left(\frac{\sec x + \tan x}{\sec x + \tan x} \right) dx \\
 &= \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} dx \\
 &= \ln |\sec x + \tan x| + C
 \end{aligned}$$

Basic Idea

We want to integrate functions like:

- $\int \sin^3 x \, dx$
- $\int \sin^5 x \cos^3 x \, dx$
- $\int \tan^3 x \sec x \, dx$
- $\int \csc^5 x \cot^2 x \, dx$



The main idea: Use trig identities to simplify the integrand until we can use u -substitution.

Pythagorean Identity

Problem 1

Compute $\int \sin^3 x \, dx$

Pythagorean Identities:

$$\int \sin^3 x \, dx = \int \sin^2 x \sin x \, dx$$

$$\sin^2 x + \cos^2 x = 1 \quad \star \quad \int (1 - \cos^2 x) \sin x \, dx$$

$$1 + \cot^2 x = \csc^2 x$$

$$\tan^2 x + 1 = \sec^2 x$$

$$= \int \sin x \, dx + \int \cos^2 x \sin x \, dx$$

$\int u^2 du$
(-1)

$$= -\cos x + \frac{1}{3} \cos^3 x + C$$

$u = \cos x$

$$\int (\cos x)^2 \sin x \, dx = \int u^2 \sin x \frac{du}{-\sin x}$$

$$\text{Let } u = \cos x$$

$$du = -\sin x \, dx$$

$$dx = \frac{du}{-\sin x}$$

$$= - \int u^2 \, du$$

$$= -\frac{1}{3} u^3 + C_0$$

$$= -\frac{1}{3} \cos^3 x + C$$

Another Example

Problem 2

Compute $\int \cos^5 x \, dx$

$$\begin{aligned}\int \cos^5 x \cos x \, dx &= \int (\cos^2 x)^2 \cos x \, dx \\&= \int (1 - \sin^2 x)^2 \cos x \, dx = \int (1 - 2\sin^2 x + \sin^4 x) \cos x \, dx \\ \text{Let } u &= \sin x, \, du = \cos x \, dx \\&= \int (1 - 2u^2 + u^4) du = u - \frac{2}{3}u^3 + \frac{1}{5}u^5 + C_0 \\&= \sin x - \frac{2}{3}\sin^3 x + \frac{1}{5}\sin^5 x + C\end{aligned}$$

Power Reduction Formulas

$$(1 - \cos 2x)^2$$

Problem 3

Compute $\int \sin^4 x \, dx$

Power Reduction Formulas

$$\sin^2 x = \frac{1 - \cos 2x}{2}$$

$$\cos^2 x = \frac{1 + \cos 2x}{2}$$

$$\int (\sin^2 x)^2 dx = \int \left(\frac{1 - \cos 2x}{2} \right)^2 dx$$

$$= \frac{1}{4} \int (1 - 2\cos 2x + \cos^2 2x) dx$$

$$= \frac{1}{4} \int dx - \frac{1}{2} \int \cos 2x dx + \frac{1}{4} \int \cos^2 2x dx$$

$$\frac{1}{4} \int \cos^2 2x dx = \frac{1}{4} \int \left(\frac{1 + \cos 4x}{2} \right) dx = \frac{1}{4} \int \left(\frac{1}{2} + \frac{1}{2} \cos 4x \right) dx$$

$$= \int \frac{1}{8} dx + \frac{1}{4} \cdot \frac{1}{8} \int \cos 4x^{(4)} dx$$

$$= \frac{1}{8} x + \frac{1}{32} \sin 4x + C_0$$

$$\int \sin^4 x = \int \frac{1}{4} dx - \frac{1}{2} \int \cos 2x^{(2)} dx + \frac{1}{4} \int \cos^2 2x dx$$

$$= \frac{1}{4} x - \frac{1}{4} \sin 2x + \frac{1}{8} x + \frac{1}{32} \sin 4x + C$$

$$= \frac{3}{8} x - \frac{1}{4} \sin 2x + \frac{1}{32} \sin 4x + C$$

When sine has an odd power

Problem 4

Compute $\int \sin^5 x \cos^2 x \, dx$

$$\begin{aligned}\int \sin^5 x \cos^2 x \, dx &= \int \sin^4 x \cos^2 x \sin x \, dx \\&= \int (\sin^2 x)^2 \cos^2 x \sin x \, dx = \int (1 - \cos^2 x)^2 \cos^2 x \sin x \, dx \\&= \int (1 - 2\cos^2 x + \cos^4 x) \cos^2 x \sin x \, dx \\&= \int (-1)(-1)(\cos^2 x - 2\cos^4 x + \cos^6 x) \sin x \, dx\end{aligned}$$

$u = \cos x$
 $du = -\sin x \, dx$

$$\begin{aligned} &= - \left[\frac{1}{3} \cos^3 x - \frac{2}{5} \cos^5 x + \frac{1}{7} \cos^7 x \right] + C \\ &= -\frac{1}{3} \cos^3 x + \frac{2}{5} \cos^5 x - \frac{1}{7} \cos^7 x + C \end{aligned}$$

When cosine has an odd power

Problem 5

Compute $\int \sin^{-\frac{3}{2}} x \cos^3 x \, dx$

$$\begin{aligned} &= \int \sin^{-\frac{3}{2}} x \cos^2 x \cos x \, dx = \int \sin^{-\frac{3}{2}} x (1 - \sin^2 x) \cos x \, dx \\ &= \int (\sin^{-\frac{3}{2}} x - \sin^{\frac{1}{2}} x) \cos x \, dx \quad \begin{array}{l} u = \sin x \\ du = \cos x \, dx \end{array} \\ &= -2 \sin^{-\frac{1}{2}} x - \frac{2}{3} \sin^{\frac{3}{2}} x + C \end{aligned}$$

When they both have even powers

$$(\sin^2 x)^2$$

Problem 6

Compute $\int \sin^4 x \cos^2 x \, dx$

$$\int \left(\frac{1 - \cos 2x}{2} \right)^2 \left(\frac{1 + \cos 2x}{2} \right) dx$$

$$= \frac{1}{8} \int (1 - \cos 2x - \cos^2 2x + \cos^3 2x) dx$$

$$= \frac{1}{8} \int \left(1 - \cos 2x - \frac{1}{2} - \frac{1}{2} \cos^4 2x + \cos^3 2x \right) dx$$

$$= \frac{1}{8} \int \underbrace{\left(\frac{1}{2} - \cos 2x - \frac{1}{2} \cos^4 2x \right)}_{(I_1)} dx + \frac{1}{8} \underbrace{\int \cos^3 2x dx}_{(I_2)}$$

$$\begin{aligned}
 (I_1): & \int \left(\frac{1}{16} - \frac{1}{8} \cos 2x - \frac{1}{16} \cos 4x \right) dx \\
 &= \int \frac{1}{16} dx - \frac{1}{2} \cdot \frac{1}{8} \int \cos 2x^{(2)} dx - \frac{1}{4} \cdot \frac{1}{16} \int \cos 4x^{(4)} dx \\
 &= \frac{1}{16} x - \frac{1}{16} \sin 2x - \frac{1}{64} \sin 4x + C_1
 \end{aligned}$$

$$\begin{aligned}
 (I_2): & \frac{1}{8} \int \cos^3 2x dx = \frac{1}{8} \int \cos^2 2x \cos 2x dx \\
 &= \frac{1}{8} \int (1 - \sin^2 2x) \cos 2x dx \\
 &= \int \frac{1}{8} dx - \frac{1}{2} \cdot \frac{1}{8} \int \sin^2 2x \cos 2x^{(2)} dx \\
 &= \frac{1}{8} x - \frac{1}{16} \cdot \frac{1}{3} \sin^3 2x + C_2
 \end{aligned}$$

$u = \sin 2x$
 $du = 2 \cos 2x dx$

$$\int \sin^4 x \cos^2 x \, dx = (I_1) + (I_2)$$
$$= \frac{1}{16} x - \frac{1}{64} \sin 4x - \frac{1}{48} \sin^3 2x + C$$

General Strategy

$\int \sin^m x \cos^n x \, dx$	Strategy
m odd and positive, n real	Split off $\sin x$, rewrite the resulting even power of $\sin x$ in terms of $\cos x$, then use $u = \cos x$.
n odd and positive, m real	Split off $\cos x$, rewrite the resulting even power of $\cos x$ in terms of $\sin x$, then use $u = \sin x$.
m and n are both even, non-negative integers	Use half-angle formulas to transform the integrand into a polynomial of $\cos 2x$, then apply the preceding strategies once again to powers of $\cos 2x$ greater than 1.

Reduction Formula (for sine) Derivation

Problem 7

Recursively compute $\int \sin^n x \, dx$

$$\int \sin^8 x \, dx = \int \left(\frac{1 - \cos^2 x}{2} \right)^4$$

Reduction Formulas

Theorem 8

Assume n is a positive integer:

$$1 \quad \int \sin^n x \, dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} \int \sin^{n-2} x \, dx$$

$$2 \quad \int \cos^n x \, dx = \frac{\cos^{n-1} x \sin x}{n} + \frac{n-1}{n} \int \cos^{n-2} x \, dx$$

$$3 \quad \int \tan^n x \, dx = \frac{\tan^{n-1} x}{n-1} - \int \tan^{n-2} x \, dx, \text{ provided } n \neq 1$$

$$4 \quad \int \sec^n x \, dx = \frac{\sec^{n-2} x \tan x}{n-1} + \frac{n-2}{n-1} \int \sec^{n-2} x \, dx, \text{ provided } n \neq 1$$

Powers of Tangent

$$\tan^2 x + 1 = \sec^2 x$$

Problem 9

Compute $\int \tan^3 x \, dx$

$$\begin{aligned}\int \tan^3 x \, dx &= \int \tan^2 x \tan x \, dx \\ &= \int (\sec^2 x - 1) \tan x \, dx \\ &= \int (\tan x \sec^2 x - \tan x) \, dx \\ u &= \tan x \quad \text{--- red arrow pointing to } \tan x \text{ in the next line} \\ du &= \sec^2 x \, dx = \int \tan x (\sec^2 x) \, dx - \int \tan x \, dx \\ &= \frac{1}{2} \tan^2 x - \ln |\sec x| + C\end{aligned}$$

Factoring out $\sec^2 x$

Problem 10

Compute $\int \tan^6 x \sec^4 x \, dx$

$$\begin{aligned}\int \tan^6 x \sec^2 x \sec^2 x \, dx &= \int \tan^6 x (\tan^2 x + 1) \sec^2 x \, dx \\ &= \int (\tan^8 x + \tan^6 x) \sec^2 x \, dx \quad \begin{array}{l} u = \tan x \\ du = \sec^2 x \, dx \end{array} \\ &= \frac{1}{9} \tan^9 x + \frac{1}{7} \tan^7 x + C\end{aligned}$$

Odd power of sec x $\tan x$

Problem 11

Compute $\int \tan^5 x \sec^7 x \, dx$

$$\begin{aligned} &= \int \tan^4 x \sec^6 x (\sec x \tan x) \, dx \quad \text{du} \\ &= \int (\sec^2 x - 1)^2 \sec^6 x [\sec x \tan x] \, dx \\ &\quad \text{Let } u = \sec x ; du = \sec x \tan x \, dx \\ &= \int (u^2 - 1)^2 u^6 \, du = \int (u^4 - 2u^2 + 1) u^6 \, du \end{aligned}$$

$$\begin{aligned} &= \int (u^{10} - 2u^8 + u^6) du \\ &= \frac{1}{11} u^{11} - \frac{2}{9} u^9 + \frac{1}{7} u^7 + C_0 \\ &= \frac{1}{11} \sec^{11} x - \frac{2}{9} \sec^9 x + \frac{1}{7} \sec^7 x + C \end{aligned}$$

General Strategy

$\int \tan^m x \sec^n x \, dx$	Strategy
n even, m real	Split off $\sec^2 x$, rewrite the resulting even power of $\sec x$ in terms of $\tan x$, then use $u = \tan x$.
m odd and positive, n real	Split off $\sec x \tan x$, rewrite the resulting even power of $\tan x$ in terms of $\sec x$, then use $u = \sec x$.
m even and positive, n odd and positive	Rewrite the even power of $\tan x$ in terms of $\sec x$ to produce a polynomial of $\sec x$, then apply Reduction Formula 4 to each term.

Cotangent

Similar strategies can be used when the integrand is powers of cotangent and cosecant.

$$1 + \cot^2 x = \csc^2 x$$

Problem 12

Compute $\int \cot^4 x \, dx$

$$\begin{aligned} \int (\cot^2 x)^2 dx &= \int (\csc^2 x - 1) \cot^2 x \, dx \\ &\stackrel{(-1)}{=} \int \cot^2 x \stackrel{(-1)}{\csc^2 x} - \int \cot^2 x \, dx \\ &= -\frac{1}{3} \cot^3 x - \int (\csc^2 x - 1) dx \quad \begin{array}{l} u = \cot x \\ du = -\csc^2 x \, dx \end{array} \\ &= -\frac{1}{3} \cot^3 x + \int -\csc^2 x \, dx + \int dx \end{aligned}$$

$$= -\frac{1}{3} \cot^3 x + \cot x + x + C$$

Square Roots

$$\int \frac{1}{\sqrt{u}} du = \int u^{-\frac{1}{2}}$$

Problem 13

Compute $\int \sqrt{1 - \sin x} \, dx$

$$\int \frac{\sqrt{1 - \sin x}}{1} \cdot \frac{\sqrt{1 + \sin x}}{\sqrt{1 + \sin x}} dx = \int \frac{\sqrt{1 - \sin^2 x}}{\sqrt{1 + \sin x}} dx$$

$$= \int \frac{\sqrt{\cos^2 x}}{\sqrt{1 + \sin x}} dx = \int \frac{\cos x}{\sqrt{1 + \sin x}} dx$$

$$u = 1 + \sin x \\ du = \cos x \, dx$$

$$= 2\sqrt{1 + \sin x} + C$$

$$\sqrt{1 - \cos x} = \sqrt{2\left(1 - \frac{\cos x}{2}\right)} \\ = \sqrt{2} \cos^2\left(\frac{x}{2}\right)$$

Powers of Secant

$$\int u dv = uv - \int v du \quad \text{"Ultra Violet"}$$

"Voo-doo"

Problem 14

Compute $\int \sec^3 x \, dx$

$$\int \underbrace{\sec x}_u \underbrace{\sec^2 x}_{dv} dx = \sec x \tan x - \int \tan^2 x \sec x \, dx$$

$$u = \sec x \quad v = \tan x$$
$$du = \sec x \tan x \, dx \quad dv = \sec^2 x \, dx$$

$$= \sec x \tan x - \int (\sec^2 x - 1) \sec x \, dx$$

$$\int \sec^3 x = \sec x \tan x - \int \sec^3 x \, dx + \int \sec x \, dx$$

$$+ \int \sec^3 x \, dx$$

$$2 \int \sec^3 x \, dx = \sec x \tan x + \int \sec x \, dx$$

$$2 \int \sec^3 x \, dx = \sec x \tan x + \ln |\sec x + \tan x| + C$$

$$\int \sec^3 x \, dx = \frac{1}{2} [\sec x \tan x + \ln |\sec x + \tan x|] + C$$

Product to Sum Formulas

Theorem 15

- $\sin A \cos B = \frac{1}{2}[\sin(A - B) + \sin(A + B)]$
- $\sin A \sin B = \frac{1}{2}[\cos(A - B) - \cos(A + B)]$
- $\cos A \cos B = \frac{1}{2}[\cos(A - B) + \cos(A + B)]$

$$\begin{aligned}\sin(A+B) &= \sin A \cos B + \sin B \cos A \\ \sin(A-B) &= \sin A \cos B - \sin B \cos A\end{aligned}$$

Using the Product to Sum Formulas

Problem 16

Compute $\int \sin 4x \cos 5x \, dx$

$\underbrace{\sin}_{A} \quad \underbrace{\cos}_{B}$

$$\begin{aligned} &= \int \frac{1}{2} [\sin(-x) + \sin(9x)] \, dx \\ &= (-1) \frac{1}{2} \int \sin(-x) \, dx + \frac{1}{9} \cdot \frac{1}{2} \int \sin(9x) \, dx \\ &= \frac{1}{2} \cos(-x) + \frac{1}{18} [-\cos(9x)] \\ &= \frac{1}{2} \cos x - \frac{1}{18} \cos 9x + C \end{aligned}$$