

Review and derived Artin stacks

Zachary Gardner

`zachary.gardner@bc.edu`

Introduction

Derived arithmetic intersection theory (DAIT) requires a derived input, so let's begin by reviewing derived algebraic geometry (DAG). The key idea of DAG is to build a more “robust” version of algebraic geometry using homotopy theory. Let's review the algebraic inputs first.

Introduction

Derived arithmetic intersection theory (DAIT) requires a derived input, so let's begin by reviewing derived algebraic geometry (DAG). The key idea of DAG is to build a more “robust” version of algebraic geometry using homotopy theory. Let's review the algebraic inputs first.

- ∞ -category dRing of derived (commutative) rings

Introduction

Derived arithmetic intersection theory (DAIT) requires a derived input, so let's begin by reviewing derived algebraic geometry (DAG). The key idea of DAG is to build a more “robust” version of algebraic geometry using homotopy theory. Let's review the algebraic inputs first.

- ∞ -category \mathbf{dRing} of derived (commutative) rings
- $A \in \mathbf{dRing}$ has homotopy groups $\pi_i(A)$ for $i \in \mathbb{Z}$, with $\pi_i(A) = 0$ for $i < 0$

Introduction

Derived arithmetic intersection theory (DAIT) requires a derived input, so let's begin by reviewing derived algebraic geometry (DAG). The key idea of DAG is to build a more “robust” version of algebraic geometry using homotopy theory. Let's review the algebraic inputs first.

- ∞ -category \mathbf{dRing} of derived (commutative) rings
- $A \in \mathbf{dRing}$ has homotopy groups $\pi_i(A)$ for $i \in \mathbb{Z}$, with $\pi_i(A) = 0$ for $i < 0$
- $\pi_0(A)$ is an ordinary commutative ring and each $\pi_i(A)$ is a module over $\pi_0(A)$

Introduction

Derived arithmetic intersection theory (DAIT) requires a derived input, so let's begin by reviewing derived algebraic geometry (DAG). The key idea of DAG is to build a more “robust” version of algebraic geometry using homotopy theory. Let's review the algebraic inputs first.

- ∞ -category \mathbf{dRing} of derived (commutative) rings
- $A \in \mathbf{dRing}$ has homotopy groups $\pi_i(A)$ for $i \in \mathbb{Z}$, with $\pi_i(A) = 0$ for $i < 0$
- $\pi_0(A)$ is an ordinary commutative ring and each $\pi_i(A)$ is a module over $\pi_0(A)$
- A is **discrete** if $\pi_i(A) = 0$ for $i \neq 0$

Introduction

Derived arithmetic intersection theory (DAIT) requires a derived input, so let's begin by reviewing derived algebraic geometry (DAG). The key idea of DAG is to build a more “robust” version of algebraic geometry using homotopy theory. Let's review the algebraic inputs first.

- ∞ -category \mathbf{dRing} of derived (commutative) rings
- $A \in \mathbf{dRing}$ has homotopy groups $\pi_i(A)$ for $i \in \mathbb{Z}$, with $\pi_i(A) = 0$ for $i < 0$
- $\pi_0(A)$ is an ordinary commutative ring and each $\pi_i(A)$ is a module over $\pi_0(A)$
- A is **discrete** if $\pi_i(A) = 0$ for $i \neq 0$
- Full subcategory of discrete derived rings in \mathbf{dRing} is equivalent to ordinary category \mathbf{CRing} of commutative rings

Introduction

Derived arithmetic intersection theory (DAIT) requires a derived input, so let's begin by reviewing derived algebraic geometry (DAG). The key idea of DAG is to build a more “robust” version of algebraic geometry using homotopy theory. Let's review the algebraic inputs first.

- ∞ -category \mathbf{dRing} of derived (commutative) rings
- $A \in \mathbf{dRing}$ has homotopy groups $\pi_i(A)$ for $i \in \mathbb{Z}$, with $\pi_i(A) = 0$ for $i < 0$
- $\pi_0(A)$ is an ordinary commutative ring and each $\pi_i(A)$ is a module over $\pi_0(A)$
- A is **discrete** if $\pi_i(A) = 0$ for $i \neq 0$
- Full subcategory of discrete derived rings in \mathbf{dRing} is equivalent to ordinary category \mathbf{CRing} of commutative rings
- ∞ -over-category \mathbf{dAlg}_A of derived (commutative) A -algebras

Modules

Modules

Fix $A \in \mathbf{dRing}$. Assume first that A is discrete and let \mathbf{Mod}_A denote the ordinary category of (left) A -modules.

Modules

Fix $A \in \mathbf{dRing}$. Assume first that A is discrete and let \mathbf{Mod}_A denote the ordinary category of (left) A -modules.

- Consider the *animation* $\mathbf{Anim}(\mathbf{Mod}_A)$

Modules

Fix $A \in \mathbf{dRing}$. Assume first that A is discrete and let \mathbf{Mod}_A denote the ordinary category of (left) A -modules.

- Consider the *animation* $\mathbf{Anim}(\mathbf{Mod}_A)$
 - ▶ Take product-preserving $(\infty\text{-})$ functors $(F_A)^{\mathrm{op}} \rightarrow \mathbf{Anim}$

Modules

Fix $A \in \mathbf{dRing}$. Assume first that A is discrete and let \mathbf{Mod}_A denote the ordinary category of (left) A -modules.

- Consider the *animation* $\mathbf{Anim}(\mathbf{Mod}_A)$
 - ▶ Take product-preserving $(\infty\text{-})$ functors $(F_A)^{\mathrm{op}} \rightarrow \mathbf{Anim}$
 - ▶ F_A full subcategory of \mathbf{Mod}_A spanned by finitely generated free A -modules

Modules

Fix $A \in \mathbf{dRing}$. Assume first that A is discrete and let \mathbf{Mod}_A denote the ordinary category of (left) A -modules.

- Consider the *animation* $\mathbf{Anim}(\mathbf{Mod}_A)$
 - ▶ Take product-preserving $(\infty\text{-})$ functors $(F_A)^{\mathrm{op}} \rightarrow \mathbf{Anim}$
 - ▶ F_A full subcategory of \mathbf{Mod}_A spanned by finitely generated free A -modules
 - ▶ ∞ -category \mathbf{Anim} of **anima**, generalization of ordinary groupoids

Modules

Fix $A \in \mathbf{dRing}$. Assume first that A is discrete and let \mathbf{Mod}_A denote the ordinary category of (left) A -modules.

- Consider the *animation* $\mathbf{Anim}(\mathbf{Mod}_A)$
 - ▶ Take product-preserving $(\infty\text{-})$ functors $(F_A)^{\mathrm{op}} \rightarrow \mathbf{Anim}$
 - ▶ F_A full subcategory of \mathbf{Mod}_A spanned by finitely generated free A -modules
 - ▶ ∞ -category \mathbf{Anim} of **anima**, generalization of ordinary groupoids
- Can define stable ∞ -category $\mathcal{D}(A)$ using spectral methods or by applying *non-connective animation* procedure to $\mathbf{Anim}(\mathbf{Mod}_A)$

Modules

Fix $A \in \mathbf{dRing}$. Assume first that A is discrete and let \mathbf{Mod}_A denote the ordinary category of (left) A -modules.

- Consider the *animation* $\mathbf{Anim}(\mathbf{Mod}_A)$
 - ▶ Take product-preserving $(\infty\text{-})$ functors $(F_A)^{\mathrm{op}} \rightarrow \mathbf{Anim}$
 - ▶ F_A full subcategory of \mathbf{Mod}_A spanned by finitely generated free A -modules
 - ▶ ∞ -category \mathbf{Anim} of **anima**, generalization of ordinary groupoids
- Can define stable ∞ -category $\mathcal{D}(A)$ using spectral methods or by applying *non-connective animation* procedure to $\mathbf{Anim}(\mathbf{Mod}_A)$
- $M \in \mathcal{D}(A)$ has homotopy groups $\pi_i(M)$ for $i \in \mathbb{Z}$, with $\pi_i(M)$ a (left) $\pi_0(A)$ -module

Modules

Fix $A \in \mathbf{dRing}$. Assume first that A is discrete and let \mathbf{Mod}_A denote the ordinary category of (left) A -modules.

- Consider the *animation* $\mathbf{Anim}(\mathbf{Mod}_A)$
 - ▶ Take product-preserving $(\infty\text{-})$ functors $(F_A)^{\mathrm{op}} \rightarrow \mathbf{Anim}$
 - ▶ F_A full subcategory of \mathbf{Mod}_A spanned by finitely generated free A -modules
 - ▶ ∞ -category \mathbf{Anim} of **anima**, generalization of ordinary groupoids
- Can define stable ∞ -category $\mathcal{D}(A)$ using spectral methods or by applying *non-connective animation* procedure to $\mathbf{Anim}(\mathbf{Mod}_A)$
- $M \in \mathcal{D}(A)$ has homotopy groups $\pi_i(M)$ for $i \in \mathbb{Z}$, with $\pi_i(M)$ a (left) $\pi_0(A)$ -module

Most of this generalizes to any $A \in \mathbf{dRing}$.

Modules

Modules

Fix $M \in \mathcal{D}(A)$ and $n \in \mathbb{Z}$.

Modules

Fix $M \in \mathcal{D}(A)$ and $n \in \mathbb{Z}$.

- M is **n -connective** if $\pi_i(M) = 0$ for every $i < n$

Modules

Fix $M \in \mathcal{D}(A)$ and $n \in \mathbb{Z}$.

- M is **n -connective** if $\pi_i(M) = 0$ for every $i < n$
 - ▶ Yields full subcategory $\mathcal{D}_{\geq n}(A) \subseteq \mathcal{D}(A)$

Modules

Fix $M \in \mathcal{D}(A)$ and $n \in \mathbb{Z}$.

- M is **n -connective** if $\pi_i(M) = 0$ for every $i < n$
 - ▶ Yields full subcategory $\mathcal{D}_{\geq n}(A) \subseteq \mathcal{D}(A)$
- M is **n -coconnective** if $\pi_i(M) = 0$ for every $i > n$

Modules

Fix $M \in \mathcal{D}(A)$ and $n \in \mathbb{Z}$.

- M is **n -connective** if $\pi_i(M) = 0$ for every $i < n$
 - ▶ Yields full subcategory $\mathcal{D}_{\geq n}(A) \subseteq \mathcal{D}(A)$
- M is **n -coconnective** if $\pi_i(M) = 0$ for every $i > n$
 - ▶ Yields full subcategory $\mathcal{D}_{\leq n}(A) \subseteq \mathcal{D}(A)$

Modules

Fix $M \in \mathcal{D}(A)$ and $n \in \mathbb{Z}$.

- M is **n -connective** if $\pi_i(M) = 0$ for every $i < n$
 - ▶ Yields full subcategory $\mathcal{D}_{\geq n}(A) \subseteq \mathcal{D}(A)$
- M is **n -coconnective** if $\pi_i(M) = 0$ for every $i > n$
 - ▶ Yields full subcategory $\mathcal{D}_{\leq n}(A) \subseteq \mathcal{D}(A)$
- M is **discrete** if $\pi_i(M) = 0$ for every $i \neq 0$

Modules

Fix $M \in \mathcal{D}(A)$ and $n \in \mathbb{Z}$.

- M is **n -connective** if $\pi_i(M) = 0$ for every $i < n$
 - ▶ Yields full subcategory $\mathcal{D}_{\geq n}(A) \subseteq \mathcal{D}(A)$
- M is **n -coconnective** if $\pi_i(M) = 0$ for every $i > n$
 - ▶ Yields full subcategory $\mathcal{D}_{\leq n}(A) \subseteq \mathcal{D}(A)$
- M is **discrete** if $\pi_i(M) = 0$ for every $i \neq 0$

This defines a natural t -structure on $\mathcal{D}(A)$ with

$$\mathcal{D}(A)^{\heartsuit} := \mathcal{D}_{\geq 0}(A) \cap \mathcal{D}_{\leq 0}(A) \simeq \text{Mod}_{\pi_0(A)}.$$

Modules

Fix $M \in \mathcal{D}(A)$ and $n \in \mathbb{Z}$.

- M is **n -connective** if $\pi_i(M) = 0$ for every $i < n$
 - ▶ Yields full subcategory $\mathcal{D}_{\geq n}(A) \subseteq \mathcal{D}(A)$
- M is **n -coconnective** if $\pi_i(M) = 0$ for every $i > n$
 - ▶ Yields full subcategory $\mathcal{D}_{\leq n}(A) \subseteq \mathcal{D}(A)$
- M is **discrete** if $\pi_i(M) = 0$ for every $i \neq 0$

This defines a natural t -structure on $\mathcal{D}(A)$ with

$$\mathcal{D}(A)^{\heartsuit} := \mathcal{D}_{\geq 0}(A) \cap \mathcal{D}_{\leq 0}(A) \simeq \mathrm{Mod}_{\pi_0(A)}.$$

For A discrete we have $\mathrm{Anim}(\mathrm{Mod}_A) \simeq \mathcal{D}(A)_{\geq 0}$.

Modules

Modules

- There is a forgetful functor $\mathrm{dAlg}_A \rightarrow \mathcal{D}(A)_{\geq 0}$

Modules

- There is a forgetful functor $\mathrm{dAlg}_A \rightarrow \mathcal{D}(A)_{\geq 0}$
- $\mathcal{D}(A)$ has a symmetric monoidal structure encoded by \otimes_A

Modules

- There is a forgetful functor $\mathrm{dAlg}_A \rightarrow \mathcal{D}(A)_{\geq 0}$
- $\mathcal{D}(A)$ has a symmetric monoidal structure encoded by \otimes_A
- For A discrete, $\mathrm{Ho}(\mathcal{D}(A)) \simeq D(A)$ with \otimes_A restricting to the usual derived tensor product $\otimes_A^{\mathbb{L}}$

Modules

- There is a forgetful functor $\mathrm{dAlg}_A \rightarrow \mathcal{D}(A)_{\geq 0}$
- $\mathcal{D}(A)$ has a symmetric monoidal structure encoded by \otimes_A
- For A discrete, $\mathrm{Ho}(\mathcal{D}(A)) \simeq D(A)$ with \otimes_A restricting to the usual derived tensor product $\otimes_A^{\mathbb{L}}$
- $\phi \in \mathrm{Hom}_{\mathrm{dRing}}(A, B)$ induces adjunction

$$\phi^* : \mathcal{D}(B) \rightleftarrows \mathcal{D}(A) : \phi_*$$

which is *not* (co-)connective in general

Modules

- There is a forgetful functor $\mathrm{dAlg}_A \rightarrow \mathcal{D}(A)_{\geq 0}$
- $\mathcal{D}(A)$ has a symmetric monoidal structure encoded by \otimes_A
- For A discrete, $\mathrm{Ho}(\mathcal{D}(A)) \simeq D(A)$ with \otimes_A restricting to the usual derived tensor product $\otimes_A^{\mathbb{L}}$
- $\phi \in \mathrm{Hom}_{\mathrm{dRing}}(A, B)$ induces adjunction

$$\phi^* : \mathcal{D}(B) \rightleftarrows \mathcal{D}(A) : \phi_*$$

which is *not* (co-)connective in general

- $M \in \mathcal{D}(A)$ is **strong** if the natural morphism

$$\pi_0(M) \otimes_{\pi_0(A)} \pi_i(A) \rightarrow \pi_i(M)$$

of $\pi_0(A)$ -modules is an isomorphism for every $i \in \mathbb{Z}$

Sheaves and Stacks

Sheaves and Stacks

The functor-of-points tells us that we can think of schemes (and, more generally, algebraic spaces) as functors $\mathbf{CRing} \rightarrow \mathbf{Set}$.

Sheaves and Stacks

The functor-of-points tells us that we can think of schemes (and, more generally, algebraic spaces) as functors $\mathbf{CRing} \rightarrow \mathbf{Set}$. Building off of this, ordinary stacks are (pseudo-)functors $\mathbf{CRing} \rightarrow \mathbf{Grpd}$ for \mathbf{Grpd} the ordinary category of groupoids.

Sheaves and Stacks

The functor-of-points tells us that we can think of schemes (and, more generally, algebraic spaces) as functors $\mathbf{CRing} \rightarrow \mathbf{Set}$. Building off of this, ordinary stacks are (pseudo-)functors $\mathbf{CRing} \rightarrow \mathbf{Grpd}$ for \mathbf{Grpd} the ordinary category of groupoids. By extension, derived stacks will be $(\infty\text{-})$ functors $\mathbf{dRing} \rightarrow \mathbf{Anim}$.

Sheaves and Stacks

The functor-of-points tells us that we can think of schemes (and, more generally, algebraic spaces) as functors $\mathbf{CRing} \rightarrow \mathbf{Set}$. Building off of this, ordinary stacks are (pseudo-)functors $\mathbf{CRing} \rightarrow \mathbf{Grpd}$ for \mathbf{Grpd} the ordinary category of groupoids. By extension, derived stacks will be $(\infty\text{-})$ functors $\mathbf{dRing} \rightarrow \mathbf{Anim}$.

We can say more. The ordinary category \mathbf{Sch} of schemes consists of functors $\mathbf{CRing} \rightarrow \mathbf{Set}$ admitting a Zariski open covering by affine schemes and satisfying a descent condition with respect to an appropriate topology (e.g., the fpqc topology).

Sheaves and Stacks

The functor-of-points tells us that we can think of schemes (and, more generally, algebraic spaces) as functors $\mathbf{CRing} \rightarrow \mathbf{Set}$. Building off of this, ordinary stacks are (pseudo-)functors $\mathbf{CRing} \rightarrow \mathbf{Grpd}$ for \mathbf{Grpd} the ordinary category of groupoids. By extension, derived stacks will be $(\infty\text{-})$ functors $\mathbf{dRing} \rightarrow \mathbf{Anim}$.

We can say more. The ordinary category \mathbf{Sch} of schemes consists of functors $\mathbf{CRing} \rightarrow \mathbf{Set}$ admitting a Zariski open covering by affine schemes and satisfying a descent condition with respect to an appropriate topology (e.g., the fpqc topology). We can simply take the category of affine schemes to be $\mathbf{Aff} := \mathbf{CRing}^{\mathrm{op}}$.

Sheaves and Stacks

The functor-of-points tells us that we can think of schemes (and, more generally, algebraic spaces) as functors $\mathbf{CRing} \rightarrow \mathbf{Set}$. Building off of this, ordinary stacks are (pseudo-)functors $\mathbf{CRing} \rightarrow \mathbf{Grpd}$ for \mathbf{Grpd} the ordinary category of groupoids. By extension, derived stacks will be $(\infty\text{-})$ functors $\mathbf{dRing} \rightarrow \mathbf{Anim}$.

We can say more. The ordinary category \mathbf{Sch} of schemes consists of functors $\mathbf{CRing} \rightarrow \mathbf{Set}$ admitting a Zariski open covering by affine schemes and satisfying a descent condition with respect to an appropriate topology (e.g., the fpqc topology). We can simply take the category of affine schemes to be $\mathbf{Aff} := \mathbf{CRing}^{\mathrm{op}}$. This generalizes to the derived setting by taking $\mathbf{dAff} := \mathbf{dRing}^{\mathrm{op}}$. What about the covering and descent conditions?

Sheaves and Stacks

Sheaves and Stacks

We need several ingredients to make sense of sheaves in the derived setting.

Sheaves and Stacks

We need several ingredients to make sense of sheaves in the derived setting.

- **Presheaves:** for ∞ -category \mathcal{C} define $\mathrm{Pre}(\mathcal{C}) := \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathrm{Anim})$

Sheaves and Stacks

We need several ingredients to make sense of sheaves in the derived setting.

- **Presheaves:** for ∞ -category \mathcal{C} define $\mathrm{Pre}(\mathcal{C}) := \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathrm{Anim})$
- **Grothendieck Topology:** yields notion of ∞ -site \mathcal{T}

Sheaves and Stacks

We need several ingredients to make sense of sheaves in the derived setting.

- **Presheaves:** for ∞ -category \mathcal{C} define $\mathrm{Pre}(\mathcal{C}) := \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathrm{Anim})$
- **Grothendieck Topology:** yields notion of ∞ -site \mathcal{T}
- **Sheaf Condition:** for simplicity, work with pretopology $\tilde{\tau}$

Sheaves and Stacks

We need several ingredients to make sense of sheaves in the derived setting.

- **Presheaves:** for ∞ -category \mathcal{C} define $\text{Pre}(\mathcal{C}) := \text{Fun}(\mathcal{C}^{\text{op}}, \text{Anim})$
- **Grothendieck Topology:** yields notion of ∞ -site \mathcal{T}
- **Sheaf Condition:** for simplicity, work with pretopology $\tilde{\tau}$

$\mathcal{F} \in \text{Pre}(\mathcal{T})$ is a **sheaf** if and only if

$$\mathcal{F}(U) \xrightarrow{\sim} \text{hlim} \left(\prod_{\alpha \in \Lambda} \mathcal{F}(U_{\alpha}) \rightrightarrows \prod_{\alpha, \beta \in \Lambda} \mathcal{F}(U_{\alpha} \times_U U_{\beta}) \cdots \right)$$

for every $\mathcal{U} = \{U_{\alpha} \rightarrow U\}_{\alpha \in \Lambda}$ in $\tilde{\tau}$. The simplicial object inside the limit is the **Čech nerve** of \mathcal{U} .

Classical Notions

Classical Notions

We can formulate the topologies we want to work with once we have the right properties of morphisms. For this it is best to start with classical notions, beginning with intuition.

Classical Notions

We can formulate the topologies we want to work with once we have the right properties of morphisms. For this it is best to start with classical notions, beginning with intuition.

- **Flat:** continuously varying fibers

Classical Notions

We can formulate the topologies we want to work with once we have the right properties of morphisms. For this it is best to start with classical notions, beginning with intuition.

- **Flat:** continuously varying fibers
- **Unramified:** like an unramified topological covering, so branches don't meet

Classical Notions

We can formulate the topologies we want to work with once we have the right properties of morphisms. For this it is best to start with classical notions, beginning with intuition.

- **Flat:** continuously varying fibers
- **Unramified:** like an unramified topological covering, so branches don't meet
- **Smooth:** smoothly varying fibers *and* good deformation properties

Classical Notions

We can formulate the topologies we want to work with once we have the right properties of morphisms. For this it is best to start with classical notions, beginning with intuition.

- **Flat:** continuously varying fibers
- **Unramified:** like an unramified topological covering, so branches don't meet
- **Smooth:** smoothly varying fibers *and* good deformation properties
- **Étale:** smooth and unramified, analogous to a local diffeomorphism

Classical Notions

Classical Notions

More rigorously, a map $\phi : A \rightarrow B$ of ordinary rings is

Classical Notions

More rigorously, a map $\phi : A \rightarrow B$ of ordinary rings is

- **flat** if $B \otimes_A (-) : \text{Mod}_A \rightarrow \text{Mod}_B$ is exact;

Classical Notions

More rigorously, a map $\phi : A \rightarrow B$ of ordinary rings is

- **flat** if $B \otimes_A (-) : \text{Mod}_A \rightarrow \text{Mod}_B$ is exact;
- **unramified** if $\Omega_{B/A}^1 = 0$ and ϕ is finitely presented (as an algebra map);

Classical Notions

More rigorously, a map $\phi : A \rightarrow B$ of ordinary rings is

- **flat** if $B \otimes_A (-) : \text{Mod}_A \rightarrow \text{Mod}_B$ is exact;
- **unramified** if $\Omega_{B/A}^1 = 0$ and ϕ is finitely presented (as an algebra map);
- **étale** if $\mathbb{L}_{B/A} \simeq 0$ and ϕ is finitely presented;

Classical Notions

More rigorously, a map $\phi : A \rightarrow B$ of ordinary rings is

- **flat** if $B \otimes_A (-) : \text{Mod}_A \rightarrow \text{Mod}_B$ is exact;
- **unramified** if $\Omega_{B/A}^1 = 0$ and ϕ is finitely presented (as an algebra map);
- **étale** if $\mathbb{L}_{B/A} \simeq 0$ and ϕ is finitely presented;
- **smooth** if $\Omega_{B/A}^1$ is projective, $\mathbb{L}_{B/A}$ is coconnective (discrete?), and ϕ is finitely presented.

Classical Notions

More rigorously, a map $\phi : A \rightarrow B$ of ordinary rings is

- **flat** if $B \otimes_A (-) : \text{Mod}_A \rightarrow \text{Mod}_B$ is exact;
- **unramified** if $\Omega_{B/A}^1 = 0$ and ϕ is finitely presented (as an algebra map);
- **étale** if $\mathbb{L}_{B/A} \simeq 0$ and ϕ is finitely presented;
- **smooth** if $\Omega_{B/A}^1$ is projective, $\mathbb{L}_{B/A}$ is coconnective (discrete?), and ϕ is finitely presented.

Here, $\Omega_{B/A}^1 \in \text{Mod}_B$ is the module of Kähler differentials characterized by

$$\text{Hom}_{\text{Mod}_B}(\Omega_{B/A}^1, M) \cong \text{Der}_A(B, M)$$

natural in $M \in \text{Mod}_B$.

Classical Notions

More rigorously, a map $\phi : A \rightarrow B$ of ordinary rings is

- **flat** if $B \otimes_A (-) : \text{Mod}_A \rightarrow \text{Mod}_B$ is exact;
- **unramified** if $\Omega_{B/A}^1 = 0$ and ϕ is finitely presented (as an algebra map);
- **étale** if $\mathbb{L}_{B/A} \simeq 0$ and ϕ is finitely presented;
- **smooth** if $\Omega_{B/A}^1$ is projective, $\mathbb{L}_{B/A}$ is coconnective (discrete?), and ϕ is finitely presented.

Here, $\Omega_{B/A}^1 \in \text{Mod}_B$ is the module of Kähler differentials characterized by

$$\text{Hom}_{\text{Mod}_B}(\Omega_{B/A}^1, M) \cong \text{Der}_A(B, M)$$

natural in $M \in \text{Mod}_B$. We “left derive” this to get the **cotangent complex** $\mathbb{L}_{B/A} \in \mathcal{D}(B)$, which satisfies a similar universal property.

The Cotangent Complex

The Cotangent Complex

Given $A \in \mathbf{CRing}$ and $M \in \mathbf{Mod}_A$, we have the split square-zero extension $A \oplus M \in \mathbf{CAlg}_A$ with

$$\begin{aligned}\mathrm{Der}_{\mathbb{Z}}(A, M) &\cong \mathrm{Hom}_{\mathbf{CRing}/A}(A, A \oplus M) \\ &= \mathrm{fib}_{\mathrm{id}_A}(\mathrm{Hom}_{\mathbf{CRing}}(A, A \oplus M), \mathrm{Hom}_{\mathbf{CRing}}(A, A))\end{aligned}$$

natural in M (\mathbb{Z} can be omitted). This lets us make sense of derivations without appealing to explicit formulas.

The Cotangent Complex

Given $A \in \mathbf{CRing}$ and $M \in \mathbf{Mod}_A$, we have the split square-zero extension $A \oplus M \in \mathbf{CAlg}_A$ with

$$\begin{aligned}\mathrm{Der}_{\mathbb{Z}}(A, M) &\cong \mathrm{Hom}_{\mathbf{CRing}/A}(A, A \oplus M) \\ &= \mathrm{fib}_{\mathrm{id}_A}(\mathrm{Hom}_{\mathbf{CRing}}(A, A \oplus M), \mathrm{Hom}_{\mathbf{CRing}}(A, A))\end{aligned}$$

natural in M (\mathbb{Z} can be omitted). This lets us make sense of derivations without appealing to explicit formulas.

Given $A \rightarrow B \rightarrow C$ in \mathbf{CRing} , there is an exact sequence

$$C \otimes_B \Omega_{B/A}^1 \rightarrow \Omega_{C/A}^1 \rightarrow \Omega_{C/B}^1 \rightarrow 0$$

in \mathbf{Mod}_C . This lets us make sense of the *relative* $\Omega_{B/A}^1$ in terms of the *absolute* Ω_A^1 and Ω_B^1 .

The Cotangent Complex

The Cotangent Complex

Let $A \in \mathbf{dRing}$, $M \in \mathcal{D}(A)_{\geq 0}$, and $B \in \mathbf{dAlg}_A$.

The Cotangent Complex

Let $A \in \mathbf{dRing}$, $M \in \mathcal{D}(A)_{\geq 0}$, and $B \in \mathbf{dAlg}_A$. We can make sense of $A \oplus M \in \mathbf{dAlg}_A$ (e.g., by using animation) and define $\mathrm{Der}(A, M)$ to be the homotopy fiber of

$$\mathrm{Hom}_{\mathbf{dRing}}(A, A \oplus M) \rightarrow \mathrm{Hom}_{\mathbf{dRing}}(A, A)$$

over id_A .

The Cotangent Complex

Let $A \in \mathbf{dRing}$, $M \in \mathcal{D}(A)_{\geq 0}$, and $B \in \mathbf{dAlg}_A$. We can make sense of $A \oplus M \in \mathbf{dAlg}_A$ (e.g., by using animation) and define $\mathrm{Der}(A, M)$ to be the homotopy fiber of

$$\mathrm{Hom}_{\mathbf{dRing}}(A, A \oplus M) \rightarrow \mathrm{Hom}_{\mathbf{dRing}}(A, A)$$

over id_A . $\mathbb{L}_A \in \mathcal{D}(A)_{\geq 0}$ then corepresents $\mathrm{Der}(A, -) : \mathcal{D}(A)_{\geq 0} \rightarrow \mathbf{Anim}$.

The Cotangent Complex

Let $A \in \mathbf{dRing}$, $M \in \mathcal{D}(A)_{\geq 0}$, and $B \in \mathbf{dAlg}_A$. We can make sense of $A \oplus M \in \mathbf{dAlg}_A$ (e.g., by using animation) and define $\mathrm{Der}(A, M)$ to be the homotopy fiber of

$$\mathrm{Hom}_{\mathbf{dRing}}(A, A \oplus M) \rightarrow \mathrm{Hom}_{\mathbf{dRing}}(A, A)$$

over id_A . $\mathbb{L}_A \in \mathcal{D}(A)_{\geq 0}$ then corepresents $\mathrm{Der}(A, -) : \mathcal{D}(A)_{\geq 0} \rightarrow \mathbf{Anim}$. We can define $\mathbb{L}_{B/A}$ as sitting in a homotopy cofiber sequence

$$B \otimes_A \mathbb{L}_A \rightarrow \mathbb{L}_B \rightarrow \mathbb{L}_{B/A}$$

in $\mathcal{D}(B)$.

The Cotangent Complex

Let $A \in \mathbf{dRing}$, $M \in \mathcal{D}(A)_{\geq 0}$, and $B \in \mathbf{dAlg}_A$. We can make sense of $A \oplus M \in \mathbf{dAlg}_A$ (e.g., by using animation) and define $\mathrm{Der}(A, M)$ to be the homotopy fiber of

$$\mathrm{Hom}_{\mathbf{dRing}}(A, A \oplus M) \rightarrow \mathrm{Hom}_{\mathbf{dRing}}(A, A)$$

over id_A . $\mathbb{L}_A \in \mathcal{D}(A)_{\geq 0}$ then corepresents $\mathrm{Der}(A, -) : \mathcal{D}(A)_{\geq 0} \rightarrow \mathbf{Anim}$. We can define $\mathbb{L}_{B/A}$ as sitting in a homotopy cofiber sequence

$$B \otimes_A \mathbb{L}_A \rightarrow \mathbb{L}_B \rightarrow \mathbb{L}_{B/A}$$

in $\mathcal{D}(B)$. Note that $\pi_0(\mathbb{L}_A) \simeq \Omega^1_{\pi_0(A)}$ and $\mathbb{L}_{B/A} \simeq 0$ if and only if $B \otimes_A \mathbb{L}_A \xrightarrow{\sim} \mathbb{L}_B$.

Derived Notions

Derived Notions

Fixing $A \in \mathbf{dRing}$, we say $P \in \mathcal{D}(A)_{\geq 0}$ is **projective** if $\mathrm{Hom}_{\mathcal{D}(A)_{\geq 0}}(P, -) : \mathcal{D}(A)_{\geq 0} \rightarrow \mathbf{Anim}$ commutes with geometric realization (of simplicial objects). This captures the usual notion when P is discrete.

Derived Notions

Fixing $A \in \mathbf{dRing}$, we say $P \in \mathcal{D}(A)_{\geq 0}$ is **projective** if $\mathrm{Hom}_{\mathcal{D}(A)_{\geq 0}}(P, -) : \mathcal{D}(A)_{\geq 0} \rightarrow \mathbf{Anim}$ commutes with geometric realization (of simplicial objects). This captures the usual notion when P is discrete. Moreover, $\mathbb{L}_{B/A} \in \mathcal{D}(B)_{\geq 0}$ for $B \in \mathbf{dAlg}_A$ is projective if and only if B is **formally infinitesimally smooth** over A . Explicitly, this means that, given any $C \rightarrow \overline{C}$ in \mathbf{dAlg}_A with $\pi_0(C) \rightarrow \pi_0(\overline{C})$ surjective with nilpotent kernel, the induced map

$$\mathrm{Hom}_{\mathbf{dAlg}_A}(B, C) \rightarrow \mathrm{Hom}_{\mathbf{dAlg}_A}(B, \overline{C})$$

is surjective on π_0 .

Derived Notions

Fixing $A \in \mathbf{dRing}$, we say $P \in \mathcal{D}(A)_{\geq 0}$ is **projective** if $\mathrm{Hom}_{\mathcal{D}(A)_{\geq 0}}(P, -) : \mathcal{D}(A)_{\geq 0} \rightarrow \mathbf{Anim}$ commutes with geometric realization (of simplicial objects). This captures the usual notion when P is discrete. Moreover, $\mathbb{L}_{B/A} \in \mathcal{D}(B)_{\geq 0}$ for $B \in \mathbf{dAlg}_A$ is projective if and only if B is **formally infinitesimally smooth** over A . Explicitly, this means that, given any $C \rightarrow \overline{C}$ in \mathbf{dAlg}_A with $\pi_0(C) \rightarrow \pi_0(\overline{C})$ surjective with nilpotent kernel, the induced map

$$\mathrm{Hom}_{\mathbf{dAlg}_A}(B, C) \rightarrow \mathrm{Hom}_{\mathbf{dAlg}_A}(B, \overline{C})$$

is surjective on π_0 . For future reference, we say B is **homotopically finitely presented** if B is compact in \mathbf{dAlg}_A – i.e., $\mathrm{Hom}_{\mathbf{dAlg}_A}(B, -)$ commutes with filtered homotopy colimits.

Derived Notions

Derived Notions

Let's now generalize our earlier classical notions to the derived setting.
What things do we want?

Derived Notions

Let's now generalize our earlier classical notions to the derived setting.
What things do we want?

- Good formal properties – e.g., stability under composition and homotopy pushouts

Derived Notions

Let's now generalize our earlier classical notions to the derived setting.
What things do we want?

- Good formal properties – e.g., stability under composition and homotopy pushouts
- Compatibility with cotangent complexes

Derived Notions

Let's now generalize our earlier classical notions to the derived setting.
What things do we want?

- Good formal properties – e.g., stability under composition and homotopy pushouts
- Compatibility with cotangent complexes
- Computability, using homotopy groups

Derived Notions

Let's now generalize our earlier classical notions to the derived setting. What things do we want?

- Good formal properties – e.g., stability under composition and homotopy pushouts
- Compatibility with cotangent complexes
- Computability, using homotopy groups

A good way to achieve these goals is to ask that $\phi : A \rightarrow B$ in \mathbf{dRing} have property Σ if and only if ϕ is strong and $\pi_0(\phi) : \pi_0(A) \rightarrow \pi_0(B)$ has property Σ (called *strongly* Σ).

Derived Notions

Let's now generalize our earlier classical notions to the derived setting. What things do we want?

- Good formal properties – e.g., stability under composition and homotopy pushouts
- Compatibility with cotangent complexes
- Computability, using homotopy groups

A good way to achieve these goals is to ask that $\phi : A \rightarrow B$ in \mathbf{dRing} have property Σ if and only if ϕ is strong and $\pi_0(\phi) : \pi_0(A) \rightarrow \pi_0(B)$ has property Σ (called *strongly* Σ). With this in mind, we say ϕ is

Derived Notions

Let's now generalize our earlier classical notions to the derived setting. What things do we want?

- Good formal properties – e.g., stability under composition and homotopy pushouts
- Compatibility with cotangent complexes
- Computability, using homotopy groups

A good way to achieve these goals is to ask that $\phi : A \rightarrow B$ in \mathbf{dRing} have property Σ if and only if ϕ is strong and $\pi_0(\phi) : \pi_0(A) \rightarrow \pi_0(B)$ has property Σ (called *strongly* Σ). With this in mind, we say ϕ is

- **flat** if $\phi^* : \mathcal{D}(B)_{\geq 0} \rightarrow \mathcal{D}(A)_{\geq 0}$ commutes with finite homotopy limits;

Derived Notions

Let's now generalize our earlier classical notions to the derived setting. What things do we want?

- Good formal properties – e.g., stability under composition and homotopy pushouts
- Compatibility with cotangent complexes
- Computability, using homotopy groups

A good way to achieve these goals is to ask that $\phi : A \rightarrow B$ in \mathbf{dRing} have property Σ if and only if ϕ is strong and $\pi_0(\phi) : \pi_0(A) \rightarrow \pi_0(B)$ has property Σ (called *strongly* Σ). With this in mind, we say ϕ is

- **flat** if $\phi^* : \mathcal{D}(B)_{\geq 0} \rightarrow \mathcal{D}(A)_{\geq 0}$ commutes with finite homotopy limits;
- **formally étale** if $\mathbb{L}_{A/B} \simeq 0$;

Derived Notions

Let's now generalize our earlier classical notions to the derived setting. What things do we want?

- Good formal properties – e.g., stability under composition and homotopy pushouts
- Compatibility with cotangent complexes
- Computability, using homotopy groups

A good way to achieve these goals is to ask that $\phi : A \rightarrow B$ in \mathbf{dRing} have property Σ if and only if ϕ is strong and $\pi_0(\phi) : \pi_0(A) \rightarrow \pi_0(B)$ has property Σ (called *strongly* Σ). With this in mind, we say ϕ is

- **flat** if $\phi^* : \mathcal{D}(B)_{\geq 0} \rightarrow \mathcal{D}(A)_{\geq 0}$ commutes with finite homotopy limits;
- **formally étale** if $\mathbb{L}_{A/B} \simeq 0$;
- **étale** if it is formally étale and homotopically finitely presented.

Derived Notions

Derived Notions

Proposition

Let $B \in \mathbf{dAlg}_A$ and $M \in \mathcal{D}(A)_{\geq 0}$.

Derived Notions

Proposition

Let $B \in \mathbf{dAlg}_A$ and $M \in \mathcal{D}(A)_{\geq 0}$.

- B is flat if and only if it is strongly flat.

Derived Notions

Proposition

Let $B \in \mathbf{dAlg}_A$ and $M \in \mathcal{D}(A)_{\geq 0}$.

- B is flat if and only if it is strongly flat.
- B is étale if and only if it is strongly étale.

Derived Notions

Proposition

Let $B \in \mathbf{dAlg}_A$ and $M \in \mathcal{D}(A)_{\geq 0}$.

- B is flat if and only if it is strongly flat.
- B is étale if and only if it is strongly étale.
- M is projective if and only if it is strongly projective. Hence, B is formally infinitesimally smooth if and only if it is strongly formally infinitesimally smooth.

Derived Notions

Proposition

Let $B \in \mathbf{dAlg}_A$ and $M \in \mathcal{D}(A)_{\geq 0}$.

- B is flat if and only if it is strongly flat.
- B is étale if and only if it is strongly étale.
- M is projective if and only if it is strongly projective. Hence, B is formally infinitesimally smooth if and only if it is strongly formally infinitesimally smooth.

Smoothness is a more subtle notion. Let us content ourselves for now by saying that B is **smooth** if it is formally infinitesimally smooth, homotopically finitely presented, and if $M \in \mathcal{D}(B)_{\geq 0}$ with $\pi_0(M) = 0$ then $[\mathbb{L}_{B/A}, M] = 0$.

Derived Notions

Proposition

Let $B \in \mathbf{dAlg}_A$ and $M \in \mathcal{D}(A)_{\geq 0}$.

- B is flat if and only if it is strongly flat.
- B is étale if and only if it is strongly étale.
- M is projective if and only if it is strongly projective. Hence, B is formally infinitesimally smooth if and only if it is strongly formally infinitesimally smooth.

Smoothness is a more subtle notion. Let us content ourselves for now by saying that B is **smooth** if it is formally infinitesimally smooth, homotopically finitely presented, and if $M \in \mathcal{D}(B)_{\geq 0}$ with $\pi_0(M) = 0$ then $[\mathbb{L}_{B/A}, M] = 0$. Note that B is smooth if and only if it is strongly smooth. Later we will encounter other notions of smoothness such as *quasi-smoothness*.

Derived Notions

Derived Notions

One last notion to cover. We say $A \rightarrow B$ in \mathbf{dRing} is a **(Zariski) open immersion** if it is flat, homotopically finitely presented, and epic (i.e., $B \otimes_A B \xrightarrow{\sim} B$).

Derived Notions

One last notion to cover. We say $A \rightarrow B$ in \mathbf{dRing} is a **(Zariski) open immersion** if it is flat, homotopically finitely presented, and epic (i.e., $B \otimes_A B \xrightarrow{\sim} B$). We have the implications

$$\text{open immersion} \implies \text{étale} \implies \text{smooth} \implies \text{flat}$$

Derived Notions

One last notion to cover. We say $A \rightarrow B$ in \mathbf{dRing} is a **(Zariski) open immersion** if it is flat, homotopically finitely presented, and epic (i.e., $B \otimes_A B \xrightarrow{\sim} B$). We have the implications

$$\text{open immersion} \implies \text{étale} \implies \text{smooth} \implies \text{flat}$$

We transfer all of our derived notions to affine derived schemes $\mathbf{dAff} := \mathbf{dRing}^{\text{op}}$ in the expected manner.

Derived Notions

One last notion to cover. We say $A \rightarrow B$ in \mathbf{dRing} is a **(Zariski) open immersion** if it is flat, homotopically finitely presented, and epic (i.e., $B \otimes_A B \xrightarrow{\sim} B$). We have the implications

$$\text{open immersion} \implies \text{étale} \implies \text{smooth} \implies \text{flat}$$

We transfer all of our derived notions to affine derived schemes $\mathbf{dAff} := \mathbf{dRing}^{\text{op}}$ in the expected manner. We then define an **étale covering** in $(\mathbf{dAff})_{/\text{Spec } A} \simeq (\mathbf{dRing}_A)^{\text{op}}$ to be a family $\{A \rightarrow A_i\}$ such that

Derived Notions

One last notion to cover. We say $A \rightarrow B$ in \mathbf{dRing} is a **(Zariski) open immersion** if it is flat, homotopically finitely presented, and epic (i.e., $B \otimes_A B \xrightarrow{\sim} B$). We have the implications

$$\text{open immersion} \implies \text{étale} \implies \text{smooth} \implies \text{flat}$$

We transfer all of our derived notions to affine derived schemes $\mathbf{dAff} := \mathbf{dRing}^{\text{op}}$ in the expected manner. We then define an **étale covering** in $(\mathbf{dAff})_{/\text{Spec } A} \simeq (\mathbf{dRing}_A)^{\text{op}}$ to be a family $\{A \rightarrow A_i\}$ such that

- the collection of $A_i \otimes_A (-) : \mathcal{D}(A)_{\geq 0} \rightarrow \mathcal{D}(A_i)_{\geq 0}$ is conservative;

Derived Notions

One last notion to cover. We say $A \rightarrow B$ in \mathbf{dRing} is a **(Zariski) open immersion** if it is flat, homotopically finitely presented, and epic (i.e., $B \otimes_A B \xrightarrow{\sim} B$). We have the implications

$$\text{open immersion} \implies \text{étale} \implies \text{smooth} \implies \text{flat}$$

We transfer all of our derived notions to affine derived schemes $\mathbf{dAff} := \mathbf{dRing}^{\mathrm{op}}$ in the expected manner. We then define an **étale covering** in $(\mathbf{dAff})_{/\mathrm{Spec} A} \simeq (\mathbf{dRing}_A)^{\mathrm{op}}$ to be a family $\{A \rightarrow A_i\}$ such that

- the collection of $A_i \otimes_A (-) : \mathcal{D}(A)_{\geq 0} \rightarrow \mathcal{D}(A_i)_{\geq 0}$ is conservative;
- each $A \rightarrow A_i$ is étale.

Derived Notions

One last notion to cover. We say $A \rightarrow B$ in \mathbf{dRing} is a **(Zariski) open immersion** if it is flat, homotopically finitely presented, and epic (i.e., $B \otimes_A B \xrightarrow{\sim} B$). We have the implications

$$\text{open immersion} \implies \text{étale} \implies \text{smooth} \implies \text{flat}$$

We transfer all of our derived notions to affine derived schemes $\mathbf{dAff} := \mathbf{dRing}^{\text{op}}$ in the expected manner. We then define an **étale covering** in $(\mathbf{dAff})_{/\text{Spec } A} \simeq (\mathbf{dRing}_A)^{\text{op}}$ to be a family $\{A \rightarrow A_i\}$ such that

- the collection of $A_i \otimes_A (-) : \mathcal{D}(A)_{\geq 0} \rightarrow \mathcal{D}(A_i)_{\geq 0}$ is conservative;
- each $A \rightarrow A_i$ is étale.

This gives us the small étale ∞ -site $(\text{Spec } A)_{\text{ét}}$.

Derived Stacks

Derived Stacks

We finally get the ∞ -category $\mathrm{dStk} := \mathrm{Shv}((\mathrm{Spec} \mathbb{Z})_{\mathrm{\acute{e}t}})$ of **derived (étale) stacks**.

Derived Stacks

We finally get the ∞ -category $\mathbf{dStk} := \mathbf{Shv}((\mathrm{Spec} \mathbb{Z})_{\acute{\mathrm{e}}\mathrm{t}})$ of **derived (étale) stacks**. We can think of $X \in \mathbf{dStk}$ as a functor $\mathbf{dRing} \rightarrow \mathbf{Anim}$ such that

Derived Stacks

We finally get the ∞ -category $\mathbf{dStk} := \mathbf{Shv}((\mathrm{Spec} \mathbb{Z})_{\mathrm{\acute{e}t}})$ of **derived (étale) stacks**. We can think of $X \in \mathbf{dStk}$ as a functor $\mathbf{dRing} \rightarrow \mathbf{Anim}$ such that

- $A_1, \dots, A_n \in \mathbf{dRing} \implies X\left(\prod_i A_i\right) \xrightarrow{\sim} \prod_i X(A_i);$

Derived Stacks

We finally get the ∞ -category $\mathrm{dStk} := \mathrm{Shv}((\mathrm{Spec} \mathbb{Z})_{\mathrm{\acute{e}t}})$ of **derived (étale) stacks**. We can think of $X \in \mathrm{dStk}$ as a functor $\mathrm{dRing} \rightarrow \mathrm{Anim}$ such that

- $A_1, \dots, A_n \in \mathrm{dRing} \implies X\left(\prod_i A_i\right) \xrightarrow{\sim} \prod_i X(A_i);$
- $A \rightarrow B$ faithfully flat étale $\implies X(A) \xrightarrow{\sim} \mathrm{hlim}_n X(\check{C}(B/A)^n).$

Here, faithful flatness is defined as you would expect.

Derived Stacks

We finally get the ∞ -category $\mathbf{dStk} := \mathrm{Shv}((\mathrm{Spec} \mathbb{Z})_{\acute{\mathrm{e}}\mathrm{t}})$ of **derived (étale) stacks**. We can think of $X \in \mathbf{dStk}$ as a functor $\mathbf{dRing} \rightarrow \mathbf{Anim}$ such that

- $A_1, \dots, A_n \in \mathbf{dRing} \implies X\left(\prod_i A_i\right) \xrightarrow{\sim} \prod_i X(A_i);$
- $A \rightarrow B$ faithfully flat étale $\implies X(A) \xrightarrow{\sim} \mathrm{hlim}_n X(\check{C}(B/A)^n).$

Here, faithful flatness is defined as you would expect.

Theorem

Affine derived schemes satisfy étale descent – i.e., the Yoneda embedding $\mathbf{dAff} \hookrightarrow \mathrm{Pre}(\mathbf{dAff})$ identifies \mathbf{dAff} with a full subcategory of \mathbf{dStk} .

Derived Schemes

Derived Schemes

Our next task is to identify the ∞ -category dSch of *derived schemes* inside of dStk .

Derived Schemes

Our next task is to identify the ∞ -category dSch of *derived schemes* inside of dStk .

- 1 Generalize notion of open immersion to morphisms of derived stacks

Derived Schemes

Our next task is to identify the ∞ -category dSch of *derived schemes* inside of dStk .

- 1 Generalize notion of open immersion to morphisms of derived stacks
- 2 Generalize notion of Zariski open covering to derived stacks

Derived Schemes

Our next task is to identify the ∞ -category dSch of *derived schemes* inside of dStk .

- 1 Generalize notion of open immersion to morphisms of derived stacks
- 2 Generalize notion of Zariski open covering to derived stacks
- 3 Describe dSch by a Zariski covering condition

Derived Schemes

Our next task is to identify the ∞ -category dSch of *derived schemes* inside of dStk .

- 1 Generalize notion of open immersion to morphisms of derived stacks
- 2 Generalize notion of Zariski open covering to derived stacks
- 3 Describe dSch by a Zariski covering condition

Let $j : U \rightarrow X$ in dStk . If U, X are both affine then we already know what it means for j to be an open immersion.

Derived Schemes

Our next task is to identify the ∞ -category dSch of *derived schemes* inside of dStk .

- 1 Generalize notion of open immersion to morphisms of derived stacks
- 2 Generalize notion of Zariski open covering to derived stacks
- 3 Describe dSch by a Zariski covering condition

Let $j : U \rightarrow X$ in dStk . If U, X are both affine then we already know what it means for j to be an open immersion. Only assuming X is affine, we say j is an **open immersion** if it is monic and there exists a family $\{U_\alpha \rightarrow U\}_{\alpha \in \Lambda}$ in dStk such that

Derived Schemes

Our next task is to identify the ∞ -category dSch of *derived schemes* inside of dStk .

- 1 Generalize notion of open immersion to morphisms of derived stacks
- 2 Generalize notion of Zariski open covering to derived stacks
- 3 Describe dSch by a Zariski covering condition

Let $j : U \rightarrow X$ in dStk . If U, X are both affine then we already know what it means for j to be an open immersion. Only assuming X is affine, we say j is an **open immersion** if it is monic and there exists a family $\{U_\alpha \rightarrow U\}_{\alpha \in \Lambda}$ in dStk such that

- each U_α is affine;

Derived Schemes

Our next task is to identify the ∞ -category dSch of *derived schemes* inside of dStk .

- 1 Generalize notion of open immersion to morphisms of derived stacks
- 2 Generalize notion of Zariski open covering to derived stacks
- 3 Describe dSch by a Zariski covering condition

Let $j : U \rightarrow X$ in dStk . If U, X are both affine then we already know what it means for j to be an open immersion. Only assuming X is affine, we say j is an **open immersion** if it is monic and there exists a family $\{U_\alpha \rightarrow U\}_{\alpha \in \Lambda}$ in dStk such that

- each U_α is affine;
- $U_\alpha \rightarrow U \rightarrow X$ is an open immersion;

Derived Schemes

Our next task is to identify the ∞ -category dSch of *derived schemes* inside of dStk .

- 1 Generalize notion of open immersion to morphisms of derived stacks
- 2 Generalize notion of Zariski open covering to derived stacks
- 3 Describe dSch by a Zariski covering condition

Let $j : U \rightarrow X$ in dStk . If U, X are both affine then we already know what it means for j to be an open immersion. Only assuming X is affine, we say j is an **open immersion** if it is monic and there exists a family $\{U_\alpha \rightarrow U\}_{\alpha \in \Lambda}$ in dStk such that

- each U_α is affine;
- $U_\alpha \rightarrow U \rightarrow X$ is an open immersion;
- $V := \coprod_{\alpha \in \Lambda} U_\alpha \rightarrow U$ satisfies $\mathrm{hcolim}_n \check{C}(V/U)_n \xrightarrow{\sim} U$.

Derived Schemes

Derived Schemes

In the general case, we require that $U \times_X \operatorname{Spec} R \rightarrow \operatorname{Spec} R$ is an open immersion for every $\operatorname{Spec} R \rightarrow X$. One then checks that all of these definitions are compatible.

Derived Schemes

In the general case, we require that $U \times_X \operatorname{Spec} R \rightarrow \operatorname{Spec} R$ is an open immersion for every $\operatorname{Spec} R \rightarrow X$. One then checks that all of these definitions are compatible.

- Define **Zariski covering** of $X \in \mathbf{dStk}$ to be a family $\{j_\alpha : U_\alpha \rightarrow X\}$ with each j_α an open immersion and $\coprod_\alpha U_\alpha \rightarrow X$ an effective epimorphism

Derived Schemes

In the general case, we require that $U \times_X \operatorname{Spec} R \rightarrow \operatorname{Spec} R$ is an open immersion for every $\operatorname{Spec} R \rightarrow X$. One then checks that all of these definitions are compatible.

- Define **Zariski covering** of $X \in \mathbf{dStk}$ to be a family $\{j_\alpha : U_\alpha \rightarrow X\}$ with each j_α an open immersion and $\coprod_\alpha U_\alpha \rightarrow X$ an effective epimorphism
- Covering is **affine** if each U_α is affine

Derived Schemes

In the general case, we require that $U \times_X \operatorname{Spec} R \rightarrow \operatorname{Spec} R$ is an open immersion for every $\operatorname{Spec} R \rightarrow X$. One then checks that all of these definitions are compatible.

- Define **Zariski covering** of $X \in \mathbf{dStk}$ to be a family $\{j_\alpha : U_\alpha \rightarrow X\}$ with each j_α an open immersion and $\coprod_\alpha U_\alpha \rightarrow X$ an effective epimorphism
- Covering is **affine** if each U_α is affine
- ∞ -category \mathbf{dSch} of **derived schemes** is full subcategory of \mathbf{dStk} of derived stacks admitting affine Zariski coverings

Derived Schemes

In the general case, we require that $U \times_X \operatorname{Spec} R \rightarrow \operatorname{Spec} R$ is an open immersion for every $\operatorname{Spec} R \rightarrow X$. One then checks that all of these definitions are compatible.

- Define **Zariski covering** of $X \in \mathbf{dStk}$ to be a family $\{j_\alpha : U_\alpha \rightarrow X\}$ with each j_α an open immersion and $\coprod_\alpha U_\alpha \rightarrow X$ an effective epimorphism
- Covering is **affine** if each U_α is affine
- ∞ -category \mathbf{dSch} of **derived schemes** is full subcategory of \mathbf{dStk} of derived stacks admitting affine Zariski coverings

We immediately get that \mathbf{dAff} is a full subcategory of \mathbf{dSch} .

Derived Schemes

In the general case, we require that $U \times_X \operatorname{Spec} R \rightarrow \operatorname{Spec} R$ is an open immersion for every $\operatorname{Spec} R \rightarrow X$. One then checks that all of these definitions are compatible.

- Define **Zariski covering** of $X \in \mathbf{dStk}$ to be a family $\{j_\alpha : U_\alpha \rightarrow X\}$ with each j_α an open immersion and $\coprod_\alpha U_\alpha \rightarrow X$ an effective epimorphism
- Covering is **affine** if each U_α is affine
- ∞ -category \mathbf{dSch} of **derived schemes** is full subcategory of \mathbf{dStk} of derived stacks admitting affine Zariski coverings

We immediately get that \mathbf{dAff} is a full subcategory of \mathbf{dSch} . Any derived stack X has an underlying **classical stack** X_{cl} , basically characterized by $(\operatorname{Spec} A)_{\text{cl}} \simeq \operatorname{Spec} \pi_0(A)$.

Extending Notions

Extending Notions

Remark

For clarity, we will use normal font (X , Y , etc.) to refer to derived schemes and calligraphic font (\mathcal{X} , \mathcal{Y} , etc.) to refer to general derived stacks.

Extending Notions

Remark

For clarity, we will use normal font (X, Y , etc.) to refer to derived schemes and calligraphic font (\mathcal{X}, \mathcal{Y} , etc.) to refer to general derived stacks.

We say $X \rightarrow Y$ in \mathbf{dSch} is **smooth (resp., flat, étale)** if there exist affine Zariski coverings $\{\mathrm{Spec} B_i \rightarrow X\}$ and $\{\mathrm{Spec} A_{j_i} \rightarrow Y\}$ and commutative squares

$$\begin{array}{ccc} \mathrm{Spec} B_i & \longrightarrow & X \\ \downarrow & & \downarrow \\ \mathrm{Spec} A_{j_i} & \longrightarrow & Y \end{array}$$

such that each $\mathrm{Spec} B_i \rightarrow \mathrm{Spec} A_{j_i}$ is smooth (resp., flat, étale).

Types of Stacks

Types of Stacks

Among all classical stacks, there are certain classes with nice properties. We give here some intuition before jumping into derived variants.

Types of Stacks

Among all classical stacks, there are certain classes with nice properties. We give here some intuition before jumping into derived variants.

- **Algebraic Spaces:** Essentially what we get if we take quotients of classical schemes by étale equivalence relations (good since the étale topology is good)

Types of Stacks

Among all classical stacks, there are certain classes with nice properties. We give here some intuition before jumping into derived variants.

- **Algebraic Spaces:** Essentially what we get if we take quotients of classical schemes by étale equivalence relations (good since the étale topology is good)
- **Artin Stacks:** Good setting for many geometric moduli problems (e.g., moduli of elliptic curves) because they encode many “stacky” groupoid quotients

Types of Stacks

Among all classical stacks, there are certain classes with nice properties. We give here some intuition before jumping into derived variants.

- **Algebraic Spaces:** Essentially what we get if we take quotients of classical schemes by étale equivalence relations (good since the étale topology is good)
- **Artin Stacks:** Good setting for many geometric moduli problems (e.g., moduli of elliptic curves) because they encode many “stacky” groupoid quotients
- **Deligne-Mumford (DM) Stacks:** Good for working with stack quotients of schemes whose automorphism groups are finite groups (analogous to orbifolds)

Derived Artin Stacks

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

- **0-Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y \in \mathbf{dSch}$ for every $Y \in \mathbf{dSch}/_{\mathcal{Y}}$;

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

- **0-Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y \in \mathbf{dSch}$ for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$;
- **0-smooth** if it is 0-Artin and $\mathcal{X} \times_{\mathcal{Y}} Y \rightarrow Y$ is smooth for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

- **0-Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y \in \mathbf{dSch}$ for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$;
- **0-smooth** if it is 0-Artin and $\mathcal{X} \times_{\mathcal{Y}} Y \rightarrow Y$ is smooth for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.

Moreover, \mathcal{X} is **0-Artin** if it is a derived scheme.

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

- **0-Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y \in \mathbf{dSch}$ for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$;
- **0-smooth** if it is 0-Artin and $\mathcal{X} \times_{\mathcal{Y}} Y \rightarrow Y$ is smooth for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.

Moreover, \mathcal{X} is **0-Artin** if it is a derived scheme. Fix now $n > 0$ and assume we have defined the relevant notions up to $n - 1$.

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

- **0-Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y \in \mathbf{dSch}$ for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$;
- **0-smooth** if it is 0-Artin and $\mathcal{X} \times_{\mathcal{Y}} Y \rightarrow Y$ is smooth for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.

Moreover, \mathcal{X} is **0-Artin** if it is a derived scheme. Fix now $n > 0$ and assume we have defined the relevant notions up to $n - 1$.

- \mathcal{X} is **n -Artin** if there exists $X \in \mathbf{dSch}_{/\mathcal{X}}$ such that $X \rightarrow \mathcal{X}$ is $(n - 1)$ -smooth and epic.

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

- **0-Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y \in \mathbf{dSch}$ for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$;
- **0-smooth** if it is 0-Artin and $\mathcal{X} \times_{\mathcal{Y}} Y \rightarrow Y$ is smooth for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.

Moreover, \mathcal{X} is **0-Artin** if it is a derived scheme. Fix now $n > 0$ and assume we have defined the relevant notions up to $n - 1$.

- \mathcal{X} is **n -Artin** if there exists $X \in \mathbf{dSch}_{/\mathcal{X}}$ such that $X \rightarrow \mathcal{X}$ is $(n - 1)$ -smooth and epic.
- $\mathcal{X} \rightarrow \mathcal{Y}$ is **n -Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y$ is n -Artin for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

- **0-Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y \in \mathbf{dSch}$ for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$;
- **0-smooth** if it is 0-Artin and $\mathcal{X} \times_{\mathcal{Y}} Y \rightarrow Y$ is smooth for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.

Moreover, \mathcal{X} is **0-Artin** if it is a derived scheme. Fix now $n > 0$ and assume we have defined the relevant notions up to $n - 1$.

- \mathcal{X} is **n -Artin** if there exists $X \in \mathbf{dSch}_{/\mathcal{X}}$ such that $X \rightarrow \mathcal{X}$ is $(n - 1)$ -smooth and epic.
- $\mathcal{X} \rightarrow \mathcal{Y}$ is **n -Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y$ is n -Artin for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.
- $\mathcal{X} \rightarrow \mathcal{Y}$ is **n -smooth** if it is n -Artin and for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$ there exists $X \in \mathbf{dSch}_{/\mathcal{X} \times_{\mathcal{Y}} Y}$ such that

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

- **0-Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y \in \mathbf{dSch}$ for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$;
- **0-smooth** if it is 0-Artin and $\mathcal{X} \times_{\mathcal{Y}} Y \rightarrow Y$ is smooth for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.

Moreover, \mathcal{X} is **0-Artin** if it is a derived scheme. Fix now $n > 0$ and assume we have defined the relevant notions up to $n - 1$.

- \mathcal{X} is **n -Artin** if there exists $X \in \mathbf{dSch}_{/\mathcal{X}}$ such that $X \rightarrow \mathcal{X}$ is $(n - 1)$ -smooth and epic.
- $\mathcal{X} \rightarrow \mathcal{Y}$ is **n -Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y$ is n -Artin for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$.
- $\mathcal{X} \rightarrow \mathcal{Y}$ is **n -smooth** if it is n -Artin and for every $Y \in \mathbf{dSch}_{/\mathcal{Y}}$ there exists $X \in \mathbf{dSch}_{/\mathcal{X} \times_{\mathcal{Y}} Y}$ such that
 - ▶ $X \rightarrow \mathcal{X} \times_{\mathcal{Y}} Y$ is $(n - 1)$ -smooth and epic;

Derived Artin Stacks

We will proceed by induction. We say $\mathcal{X} \rightarrow \mathcal{Y}$ in \mathbf{dStk} is

- **0-Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y \in \mathbf{dSch}$ for every $Y \in \mathbf{dSch}/_{\mathcal{Y}}$;
- **0-smooth** if it is 0-Artin and $\mathcal{X} \times_{\mathcal{Y}} Y \rightarrow Y$ is smooth for every $Y \in \mathbf{dSch}/_{\mathcal{Y}}$.

Moreover, \mathcal{X} is **0-Artin** if it is a derived scheme. Fix now $n > 0$ and assume we have defined the relevant notions up to $n - 1$.

- \mathcal{X} is **n -Artin** if there exists $X \in \mathbf{dSch}/_{\mathcal{X}}$ such that $X \rightarrow \mathcal{X}$ is $(n - 1)$ -smooth and epic.
- $\mathcal{X} \rightarrow \mathcal{Y}$ is **n -Artin** if $\mathcal{X} \times_{\mathcal{Y}} Y$ is n -Artin for every $Y \in \mathbf{dSch}/_{\mathcal{Y}}$.
- $\mathcal{X} \rightarrow \mathcal{Y}$ is **n -smooth** if it is n -Artin and for every $Y \in \mathbf{dSch}/_{\mathcal{Y}}$ there exists $X \in \mathbf{dSch}/_{\mathcal{X} \times_{\mathcal{Y}} Y}$ such that
 - ▶ $X \rightarrow \mathcal{X} \times_{\mathcal{Y}} Y$ is $(n - 1)$ -smooth and epic;
 - ▶ $X \rightarrow \mathcal{X} \times_{\mathcal{Y}} Y \rightarrow Y$ is smooth.

Derived Artin Stacks

Derived Artin Stacks

Fix $\mathcal{X} \in \mathbf{dStk}$.

Derived Artin Stacks

Fix $\mathcal{X} \in \mathbf{dStk}$.

- \mathcal{X} is a **derived Artin stack** if it is n -Artin for some $n \geq 0$.

Derived Artin Stacks

Fix $\mathcal{X} \in \mathbf{dStk}$.

- \mathcal{X} is a **derived Artin stack** if it is n -Artin for some $n \geq 0$.
- \mathcal{X} is a **derived Deligne-Mumford (DM) stack** if it is a derived Artin stack and there exists an étale surjection $X \rightarrow \mathcal{X}$ with $X \in \mathbf{dSch}$.

Derived Artin Stacks

Fix $\mathcal{X} \in \mathbf{dStk}$.

- \mathcal{X} is a **derived Artin stack** if it is n -Artin for some $n \geq 0$.
- \mathcal{X} is a **derived Deligne-Mumford (DM) stack** if it is a derived Artin stack and there exists an étale surjection $X \rightarrow \mathcal{X}$ with $X \in \mathbf{dSch}$.

We take *derived algebraic stack* to be synonymous with derived Artin stack.

Derived Artin Stacks

Fix $\mathcal{X} \in \mathbf{dStk}$.

- \mathcal{X} is a **derived Artin stack** if it is n -Artin for some $n \geq 0$.
- \mathcal{X} is a **derived Deligne-Mumford (DM) stack** if it is a derived Artin stack and there exists an étale surjection $X \rightarrow \mathcal{X}$ with $X \in \mathbf{dSch}$.

We take *derived algebraic stack* to be synonymous with derived Artin stack. Note that one can give equivalent descriptions of derived Artin and DM stacks in terms of representability of the diagonal and derived quotient stacks.