

Virtual intersection theories

Zachary Gardner

`zachary.gardner@bc.edu`

Our goal is to describe a general context for intersection theories and virtual fundamental classes, mostly following the axiomatic framework of Kiem and Park. We start with a mostly technical consideration. Let St_k denote the (2-)category of qs LFT algebraic stacks over k , for k a field of characteristic zero (which need not be algebraically closed). We fix an *admissible* full (2-)subcategory $V \subseteq \mathrm{St}_k$, which is by definition required to satisfy the following properties.

Our goal is to describe a general context for intersection theories and virtual fundamental classes, mostly following the axiomatic framework of Kiem and Park. We start with a mostly technical consideration. Let St_k denote the (2-)category of qs LFT algebraic stacks over k , for k a field of characteristic zero (which need not be algebraically closed). We fix an *admissible* full (2-)subcategory $V \subseteq \mathrm{St}_k$, which is by definition required to satisfy the following properties.

- $\emptyset, \mathrm{Spec} k \in V$.

Our goal is to describe a general context for intersection theories and virtual fundamental classes, mostly following the axiomatic framework of Kiem and Park. We start with a mostly technical consideration. Let St_k denote the (2-)category of qs LFT algebraic stacks over k , for k a field of characteristic zero (which need not be algebraically closed). We fix an *admissible* full (2-)subcategory $V \subseteq \mathrm{St}_k$, which is by definition required to satisfy the following properties.

- $\emptyset, \mathrm{Spec} k \in V$.
- V is closed under finite coproducts.

Our goal is to describe a general context for intersection theories and virtual fundamental classes, mostly following the axiomatic framework of Kiem and Park. We start with a mostly technical consideration. Let St_k denote the (2-)category of qs LFT algebraic stacks over k , for k a field of characteristic zero (which need not be algebraically closed). We fix an *admissible* full (2-)subcategory $V \subseteq \mathrm{St}_k$, which is by definition required to satisfy the following properties.

- $\emptyset, \mathrm{Spec} k \in V$.
- V is closed under finite coproducts.
- V is closed under fiber products.

Our goal is to describe a general context for intersection theories and virtual fundamental classes, mostly following the axiomatic framework of Kiem and Park. We start with a mostly technical consideration. Let St_k denote the (2-)category of qs LFT algebraic stacks over k , for k a field of characteristic zero (which need not be algebraically closed). We fix an *admissible* full (2-)subcategory $V \subseteq \mathrm{St}_k$, which is by definition required to satisfy the following properties.

- $\emptyset, \mathrm{Spec} k \in V$.
- V is closed under finite coproducts.
- V is closed under fiber products.
- If $X \rightarrow Y$ is a qs morphism in St_k with $Y \in V$ then $X \in V$.

Our goal is to describe a general context for intersection theories and virtual fundamental classes, mostly following the axiomatic framework of Kiem and Park. We start with a mostly technical consideration. Let St_k denote the (2-)category of qs LFT algebraic stacks over k , for k a field of characteristic zero (which need not be algebraically closed). We fix an *admissible* full (2-)subcategory $V \subseteq \mathrm{St}_k$, which is by definition required to satisfy the following properties.

- $\emptyset, \mathrm{Spec} k \in V$.
- V is closed under finite coproducts.
- V is closed under fiber products.
- If $X \rightarrow Y$ is a qs morphism in St_k with $Y \in V$ then $X \in V$.

Good examples of admissible categories are St_k as well as the categories Sch_k of LFT k -schemes and QSch_k its full subcategory of qs schemes (viewing both as 2-categories in the trivial way).

The following constitute the data of an *intersection theory* for V .

The following constitute the data of an *intersection theory* for V .

- For $X \in V$, we have a \mathbb{Z} -graded abelian group $H_*(X)$.

The following constitute the data of an *intersection theory* for V .

- For $X \in V$, we have a \mathbb{Z} -graded abelian group $H_*(X)$.
- *Projective pushforward*: For $f : X \rightarrow Y$ projective, we have a graded map

$$f_* : H_*(X) \rightarrow H_*(Y).$$

The following constitute the data of an *intersection theory* for V .

- For $X \in V$, we have a \mathbb{Z} -graded abelian group $H_*(X)$.
- *Projective pushforward*: For $f : X \rightarrow Y$ projective, we have a graded map

$$f_* : H_*(X) \rightarrow H_*(Y).$$

- *Smooth pullback*: For $f : X \rightarrow Y$ smooth of constant relative dimension e , we have a graded map

$$f^* : H_*(Y) \rightarrow H_{*+e}(X).$$

The following constitute the data of an *intersection theory* for V .


- For $X \in V$, we have a \mathbb{Z} -graded abelian group $H_*(X)$.
- *Projective pushforward*: For $f : X \rightarrow Y$ projective, we have a graded map

$$f_* : H_*(X) \rightarrow H_*(Y).$$

- *Smooth pullback*: For $f : X \rightarrow Y$ smooth of constant relative dimension e , we have a graded map

$$f^* : H_*(Y) \rightarrow H_{*+e}(X).$$

Note that the context of more narrow theories we also have access to more general *proper* pushforwards and *flat* pullbacks.

¹Here, $X \times Y$ is the product formed in V and so is naturally fibered over $\text{Spec } k$. 

- *Refined Gysin pullback*: For a Cartesian square

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ g' \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Y \end{array}$$

with f a regular immersion of constant codimension c , we have a graded map

$$f^! : H_*(Y') \rightarrow H_{*-c}(X').$$

¹Here, $X \times Y$ is the product formed in V and so is naturally fibered over $\text{Spec } k$.

- *Refined Gysin pullback*: For a Cartesian square

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ g' \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Y \end{array}$$

with f a regular immersion of constant codimension c , we have a graded map

$$f^! : H_*(Y') \rightarrow H_{*-c}(X').$$

- *Exterior product*: For $X, Y \in V$, we have a bilinear graded map

$$\times : H_*(X) \otimes_{\mathbb{Z}} H_*(Y) \rightarrow H_*(X \times Y)$$

which is commutative and associative with a distinguished element $1 \in H_0(\operatorname{Spec} k)$ as unit.¹

¹Here, $X \times Y$ is the product formed in V and so is naturally fibered over $\operatorname{Spec} k$.

Some of the conditions we place on these data are obvious.

Some of the conditions we place on these data are obvious.

- $H_*(\emptyset) = 0$.

Some of the conditions we place on these data are obvious.

- $H_*(\emptyset) = 0$.
- Projective pushforward and smooth pullback are functorial (the former covariant and the latter contravariant).

Some of the conditions we place on these data are obvious.

- $H_*(\emptyset) = 0$.
- Projective pushforward and smooth pullback are functorial (the former covariant and the latter contravariant).
- Refined Gysin pullback is (contravariantly) functorial.

Some of the conditions we place on these data are obvious.

- $H_*(\emptyset) = 0$.
- Projective pushforward and smooth pullback are functorial (the former covariant and the latter contravariant).
- Refined Gysin pullback is (contravariantly) functorial.
- Exterior products are compatible with projective pushforward, smooth pullback, and refined Gysin pullback.

Some of the conditions we place on these data are obvious.

- $H_*(\emptyset) = 0$.
- Projective pushforward and smooth pullback are functorial (the former covariant and the latter contravariant).
- Refined Gysin pullback is (contravariantly) functorial.
- Exterior products are compatible with projective pushforward, smooth pullback, and refined Gysin pullback.
- Given $X, Y \in V$, there are obvious maps $i : X \rightarrow X \sqcup Y$ and $j : Y \rightarrow X \sqcup Y$ and the induced map

$$(i_*, j_*) : H_*(X) \oplus H_*(Y) \rightarrow H_*(X \sqcup Y)$$

is an isomorphism.

Some of the conditions we place on these data are obvious.

- $H_*(\emptyset) = 0$.
- Projective pushforward and smooth pullback are functorial (the former covariant and the latter contravariant).
- Refined Gysin pullback is (contravariantly) functorial.
- Exterior products are compatible with projective pushforward, smooth pullback, and refined Gysin pullback.
- Given $X, Y \in V$, there are obvious maps $i : X \rightarrow X \sqcup Y$ and $j : Y \rightarrow X \sqcup Y$ and the induced map

$$(i_*, j_*) : H_*(X) \oplus H_*(Y) \rightarrow H_*(X \sqcup Y)$$

is an isomorphism.

The other conditions are more mysterious or at least require more geometric context. But before that, some examples!

G-Theory Operations

G-Theory Operations

Given $X \in V$, endow $G_0(X)$ with a grading by considering

$$G_0(X) \otimes_{\mathbb{Z}} \mathbb{Z}[\beta^{\pm 1}] \cong \bigoplus_{n \in \mathbb{Z}} G_0(X) \cdot \beta^n$$

with $G_0(X)$ in degree 0 and β in degree 1.

G-Theory Operations

Given $X \in V$, endow $G_0(X)$ with a grading by considering

$$G_0(X) \otimes_{\mathbb{Z}} \mathbb{Z}[\beta^{\pm 1}] \cong \bigoplus_{n \in \mathbb{Z}} G_0(X) \cdot \beta^n$$

with $G_0(X)$ in degree 0 and β in degree 1.

- *Projective pushforward*: For $f : X \rightarrow Y$ projective, we have f_* with formula

$$f_* : [E] \cdot \beta^n \mapsto [Rf_* E] \cdot \beta^n.$$

G-Theory Operations

Given $X \in V$, endow $G_0(X)$ with a grading by considering

$$G_0(X) \otimes_{\mathbb{Z}} \mathbb{Z}[\beta^{\pm 1}] \cong \bigoplus_{n \in \mathbb{Z}} G_0(X) \cdot \beta^n$$

with $G_0(X)$ in degree 0 and β in degree 1.

- *Projective pushforward*: For $f : X \rightarrow Y$ projective, we have f_* with formula

$$f_* : [E] \cdot \beta^n \mapsto [Rf_* E] \cdot \beta^n.$$

- *Smooth pullback*: For $f : X \rightarrow Y$ smooth of constant relative dimension e , we have f^* with formula

$$f^* : [E] \cdot \beta^n \mapsto [f^* E] \cdot \beta^{n+e}.$$

G-Theory Operations

G-Theory Operations

- *Refined Gysin pullback*: This is a bit subtle but the trick is to work with derived tensor products for suitable Cartesian squares. The only thing to do is some kind of push-pull construction.

G-Theory Operations

- *Refined Gysin pullback*: This is a bit subtle but the trick is to work with derived tensor products for suitable Cartesian squares. The only thing to do is some kind of push-pull construction.
- *Exterior product*: For $X, Y \in V$, we have \times with formula

$$\times : ([E] \cdot \beta^n) \otimes ([E'] \cdot \beta^m) \mapsto [E \boxtimes E'] \cdot \beta^{n+m}.$$

This is commutative and associative with a distinguished element $1 \in G_0(\mathrm{Spec} k) \cong \mathbb{Z}$ as unit.


G-Theory Operations

- *Refined Gysin pullback*: This is a bit subtle but the trick is to work with derived tensor products for suitable Cartesian squares. The only thing to do is some kind of push-pull construction.
- *Exterior product*: For $X, Y \in V$, we have \times with formula

$$\times : ([E] \cdot \beta^n) \otimes ([E'] \cdot \beta^m) \mapsto [E \boxtimes E'] \cdot \beta^{n+m}.$$

This is commutative and associative with a distinguished element $1 \in G_0(\operatorname{Spec} k) \cong \mathbb{Z}$ as unit. Note that, by definition, $E \boxtimes E'$ is given by the tensor product $\operatorname{pr}_X^{-1} E \otimes_{\mathcal{O}_{X \times Y}} \operatorname{pr}_Y^{-1} E'$ for $\operatorname{pr}_X : X \times Y \rightarrow X$ and $\operatorname{pr}_Y : X \times Y \rightarrow Y$ the projection maps.

Algebraic Cycles

² “Nice” means separated, Noetherian, finite dimensional, and excellent. 

Algebraic Cycles

Our second and arguably most important example comes from Chow theory. For simplicity, let Sch denote the category of “nice” schemes.² We will try to do things carefully to clarify some things and make our lives easier in the future. In particular, we will be careful about matters of (co-)dimension as well as rational equivalence.

² “Nice” means separated, Noetherian, finite dimensional, and excellent. 

Algebraic Cycles

Our second and arguably most important example comes from Chow theory. For simplicity, let Sch denote the category of “nice” schemes.² We will try to do things carefully to clarify some things and make our lives easier in the future. In particular, we will be careful about matters of (co-)dimension as well as rational equivalence.

- For *any* (general) scheme X , we let $Z(X)$ denote the free abelian group on the set of closed integral subschemes of X , whose elements are called (*algebraic*) *cycles* on X .

² “Nice” means separated, Noetherian, finite dimensional, and excellent. 

Algebraic Cycles

Our second and arguably most important example comes from Chow theory. For simplicity, let Sch denote the category of “nice” schemes.² We will try to do things carefully to clarify some things and make our lives easier in the future. In particular, we will be careful about matters of (co-)dimension as well as rational equivalence.

- For *any* (general) scheme X , we let $Z(X)$ denote the free abelian group on the set of closed integral subschemes of X , whose elements are called (*algebraic*) *cycles* on X .
- The closed integral subschemes of X correspond bijectively with the (topological, field-valued) points of X . This is given by looking at generic points ζ and we have

$$Z(X) \cong \bigoplus_{\zeta \in X} \mathbb{Z}.$$

² “Nice” means separated, Noetherian, finite dimensional, and excellent. 

(Co-)dimension

(Co-)dimension

As promised, let's think about (co-)dimension.

(Co-)dimension

As promised, let's think about (co-)dimension.

- If X is Noetherian we can grade $Z(X)$ by codimension with pieces $Z^p(X)$, with elements of $Z^1(X)$ called *Weil divisors*.

(Co-)dimension

As promised, let's think about (co-)dimension.

- If X is Noetherian we can grade $Z(X)$ by codimension with pieces $Z^p(X)$, with elements of $Z^1(X)$ called *Weil divisors*.
- If all the closed integral subschemes of X also have finite dimension then we can grade $Z(X)$ by dimension with pieces $Z_p(X)$.

(Co-)dimension

As promised, let's think about (co-)dimension.

- If X is Noetherian we can grade $Z(X)$ by codimension with pieces $Z^p(X)$, with elements of $Z^1(X)$ called *Weil divisors*.
- If all the closed integral subschemes of X also have finite dimension then we can grade $Z(X)$ by dimension with pieces $Z_p(X)$.
- If X is Noetherian, catenary, and equidimensional of finite dimension d then we can switch between the two gradings via $Z_p(X) = Z^{d-p}(X)$.

(Co-)dimension

As promised, let's think about (co-)dimension.

- If X is Noetherian we can grade $Z(X)$ by codimension with pieces $Z^p(X)$, with elements of $Z^1(X)$ called *Weil divisors*.
- If all the closed integral subschemes of X also have finite dimension then we can grade $Z(X)$ by dimension with pieces $Z_p(X)$.
- If X is Noetherian, catenary, and equidimensional of finite dimension d then we can switch between the two gradings via $Z_p(X) = Z^{d-p}(X)$.

We define the sets $X^{(p)}$ and $X_{(p)}$ so that $Z^p(X) \cong \bigoplus_{x \in X^{(p)}} \mathbb{Z}$ and $Z_p(X) \cong \bigoplus_{x \in X_{(p)}} \mathbb{Z}$ – these are useful for bookkeeping purposes.

(Co-)dimension

As promised, let's think about (co-)dimension.

- If X is Noetherian we can grade $Z(X)$ by codimension with pieces $Z^p(X)$, with elements of $Z^1(X)$ called *Weil divisors*.
- If all the closed integral subschemes of X also have finite dimension then we can grade $Z(X)$ by dimension with pieces $Z_p(X)$.
- If X is Noetherian, catenary, and equidimensional of finite dimension d then we can switch between the two gradings via $Z_p(X) = Z^{d-p}(X)$.

We define the sets $X^{(p)}$ and $X_{(p)}$ so that $Z^p(X) \cong \bigoplus_{x \in X^{(p)}} \mathbb{Z}$ and $Z_p(X) \cong \bigoplus_{x \in X_{(p)}} \mathbb{Z}$ – these are useful for bookkeeping purposes.

Geometrically, we can glue to get the flasque cycle sheaves \mathcal{Z}_X^p .

Rational Equivalence

Rational Equivalence

For general X , we have the sheaf $\mathcal{O}_X^{\text{reg}}$ of regular elements (which can be defined at the level of stalks). To this we associate $\mathcal{K}_X := \mathcal{O}_X[(\mathcal{O}_X^{\text{reg}})^{-1}]$ and $\mathcal{D}\text{iv}_X := \mathcal{K}_X^\times / \mathcal{O}_X^\times$. These give rise to the additive group of *Cartier divisors* $\text{Div}(X) := H^0(X, \mathcal{D}\text{iv}_X)$.

Rational Equivalence

For general X , we have the sheaf $\mathcal{O}_X^{\text{reg}}$ of regular elements (which can be defined at the level of stalks). To this we associate $\mathcal{K}_X := \mathcal{O}_X[(\mathcal{O}_X^{\text{reg}})^{-1}]$ and $\mathcal{D}\text{iv}_X := \mathcal{K}_X^\times / \mathcal{O}_X^\times$. These give rise to the additive group of *Cartier divisors* $\text{Div}(X) := H^0(X, \mathcal{D}\text{iv}_X)$.

- There is a unique Zariski sheaf morphism $\text{div} : \mathcal{D}\text{iv}_X \rightarrow \mathcal{Z}_X^1$ satisfying $\text{div}(f) = [\mathcal{O}_U/(f)]$ for $U \subseteq X$ open and $f \in \mathcal{O}_X^{\text{reg}}(U)$.

Rational Equivalence

For general X , we have the sheaf $\mathcal{O}_X^{\text{reg}}$ of regular elements (which can be defined at the level of stalks). To this we associate $\mathcal{K}_X := \mathcal{O}_X[(\mathcal{O}_X^{\text{reg}})^{-1}]$ and $\mathcal{D}\text{iv}_X := \mathcal{K}_X^\times / \mathcal{O}_X^\times$. These give rise to the additive group of *Cartier divisors* $\text{Div}(X) := H^0(X, \mathcal{D}\text{iv}_X)$.

- There is a unique Zariski sheaf morphism $\text{div} : \mathcal{D}\text{iv}_X \rightarrow \mathcal{Z}_X^1$ satisfying $\text{div}(f) = [\mathcal{O}_U/(f)]$ for $U \subseteq X$ open and $f \in \mathcal{O}_X^{\text{reg}}(U)$.
- If X is regular (or even locally factorial) then div is an isomorphism.

Rational Equivalence

For general X , we have the sheaf $\mathcal{O}_X^{\text{reg}}$ of regular elements (which can be defined at the level of stalks). To this we associate $\mathcal{K}_X := \mathcal{O}_X[(\mathcal{O}_X^{\text{reg}})^{-1}]$ and $\mathcal{D}\text{iv}_X := \mathcal{K}_X^\times / \mathcal{O}_X^\times$. These give rise to the additive group of *Cartier divisors* $\text{Div}(X) := H^0(X, \mathcal{D}\text{iv}_X)$.

- There is a unique Zariski sheaf morphism $\text{div} : \mathcal{D}\text{iv}_X \rightarrow \mathcal{Z}_X^1$ satisfying $\text{div}(f) = [\mathcal{O}_U/(f)]$ for $U \subseteq X$ open and $f \in \mathcal{O}_X^{\text{reg}}(U)$.
- If X is regular (or even locally factorial) then div is an isomorphism.
- If X is integral then \mathcal{K}_X is the constant sheaf associated to $K(X)$ and so we have $\text{div} : K(X)^\times \rightarrow Z^1(X)$.

Rational Equivalence

For general X , we have the sheaf $\mathcal{O}_X^{\text{reg}}$ of regular elements (which can be defined at the level of stalks). To this we associate $\mathcal{K}_X := \mathcal{O}_X[(\mathcal{O}_X^{\text{reg}})^{-1}]$ and $\mathcal{D}\text{iv}_X := \mathcal{K}_X^\times / \mathcal{O}_X^\times$. These give rise to the additive group of *Cartier divisors* $\text{Div}(X) := H^0(X, \mathcal{D}\text{iv}_X)$.

- There is a unique Zariski sheaf morphism $\text{div} : \mathcal{D}\text{iv}_X \rightarrow \mathcal{Z}_X^1$ satisfying $\text{div}(f) = [\mathcal{O}_U/(f)]$ for $U \subseteq X$ open and $f \in \mathcal{O}_X^{\text{reg}}(U)$.
- If X is regular (or even locally factorial) then div is an isomorphism.
- If X is integral then \mathcal{K}_X is the constant sheaf associated to $K(X)$ and so we have $\text{div} : K(X)^\times \rightarrow Z^1(X)$.

Define $R(X) := \bigoplus_{\zeta \in X} k(\zeta)^\times$, which can be graded by (co-)dimension as appropriate.

Rational Equivalence

For general X , we have the sheaf $\mathcal{O}_X^{\text{reg}}$ of regular elements (which can be defined at the level of stalks). To this we associate $\mathcal{K}_X := \mathcal{O}_X[(\mathcal{O}_X^{\text{reg}})^{-1}]$ and $\text{Div}_X := \mathcal{K}_X^\times / \mathcal{O}_X^\times$. These give rise to the additive group of *Cartier divisors* $\text{Div}(X) := H^0(X, \text{Div}_X)$.

- There is a unique Zariski sheaf morphism $\text{div} : \text{Div}_X \rightarrow \mathcal{Z}_X^1$ satisfying $\text{div}(f) = [\mathcal{O}_U/(f)]$ for $U \subseteq X$ open and $f \in \mathcal{O}_X^{\text{reg}}(U)$.
- If X is regular (or even locally factorial) then div is an isomorphism.
- If X is integral then \mathcal{K}_X is the constant sheaf associated to $K(X)$ and so we have $\text{div} : K(X)^\times \rightarrow \mathcal{Z}^1(X)$.

Define $R(X) := \bigoplus_{\zeta \in X} k(\zeta)^\times$, which can be graded by (co-)dimension as appropriate. The amalgamation of $\text{div} : K(Z)^\times \rightarrow \mathcal{Z}^1(Z)$ for $Z \subseteq X$ closed integral induces $\text{div} : R(X) \rightarrow \mathcal{Z}(X)$ and we set $\text{CH}(X) := \text{coker}(\text{div} : R(X) \rightarrow \mathcal{Z}(X))$. Being *rationaly equivalent to zero* is captured by the image of div .

Rational Equivalence

Rational Equivalence

Suppose now that $X \in \text{Sch}$.

Rational Equivalence

Suppose now that $X \in \text{Sch}$.

- The map div is of pure degree -1 for the dimension grading and we set $\text{CH}_p(X) := \text{coker}(\text{div} : R_{p+1}(X) \rightarrow Z_p(X))$.

Rational Equivalence

Suppose now that $X \in \text{Sch}$.

- The map div is of pure degree -1 for the dimension grading and we set $\text{CH}_p(X) := \text{coker}(\text{div} : R_{p+1}(X) \rightarrow Z_p(X))$.
- The map div is not generally of pure degree $+1$ for the codimension grading but we can get around this by defining $\text{CH}^p(X)$ to be the cokernel of

$$\text{div} : \bigoplus_{x \in X^{(p-1)}} k(x)^\times \rightarrow \bigoplus_{x \in X^{(p)}} \mathbb{Z}, \quad \sum_x \{f_x\} \mapsto \sum_x \text{div}(\{f_x\}).$$

Rational Equivalence

Suppose now that $X \in \text{Sch}$.

- The map div is of pure degree -1 for the dimension grading and we set $\text{CH}_p(X) := \text{coker}(\text{div} : R_{p+1}(X) \rightarrow Z_p(X))$.
- The map div is not generally of pure degree $+1$ for the codimension grading but we can get around this by defining $\text{CH}^p(X)$ to be the cokernel of

$$\text{div} : \bigoplus_{x \in X^{(p-1)}} k(x)^\times \rightarrow \bigoplus_{x \in X^{(p)}} \mathbb{Z}, \quad \sum_x \{f_x\} \mapsto \sum_x \text{div}(\{f_x\}).$$

- If X is equidimensional of dimension d then these notions are related via $\text{CH}_p(X) \cong \text{CH}^{d-p}(X)$.

Rational Equivalence

Suppose now that $X \in \text{Sch}$.

- The map div is of pure degree -1 for the dimension grading and we set $\text{CH}_p(X) := \text{coker}(\text{div} : R_{p+1}(X) \rightarrow Z_p(X))$.
- The map div is not generally of pure degree $+1$ for the codimension grading but we can get around this by defining $\text{CH}^p(X)$ to be the cokernel of

$$\text{div} : \bigoplus_{x \in X^{(p-1)}} k(x)^\times \rightarrow \bigoplus_{x \in X^{(p)}} \mathbb{Z}, \quad \sum_x \{f_x\} \mapsto \sum_x \text{div}(\{f_x\}).$$

- If X is equidimensional of dimension d then these notions are related via $\text{CH}_p(X) \cong \text{CH}^{d-p}(X)$.

Note that there is another equivalent approach to rational equivalence that works for varieties over a field. Heuristically, two cycles α, β are rationally equivalent if there is a family of cycles $\{\zeta_t\}_{t \in \mathbb{P}^1}$ with $\zeta_0 = \alpha$ and $\zeta_\infty = \beta$. This can be made more precise using the language of flat families.

Chow Operations

Chow Operations

Let $f : X \rightarrow Y$ be a map of sufficiently nice schemes.

Chow Operations

Let $f : X \rightarrow Y$ be a map of sufficiently nice schemes.

- If f is flat then we have the pullback

$$f^* : Z^p(Y) \rightarrow Z^p(X), \quad [Z] \mapsto [\mathcal{O}_X \otimes_{\mathcal{O}_Y} \mathcal{O}_Z],$$

which must be interpreted in terms of support and length

Chow Operations

Let $f : X \rightarrow Y$ be a map of sufficiently nice schemes.

- If f is flat then we have the pullback

$$f^* : Z^p(Y) \rightarrow Z^p(X), \quad [Z] \mapsto [\mathcal{O}_X \otimes_{\mathcal{O}_Y} \mathcal{O}_Z],$$

which must be interpreted in terms of support and length

- If f is proper then we have the pushforward $f_* : Z_p(X) \rightarrow Z_p(Y)$ given explicitly by

$$[Z] \mapsto \begin{cases} [K(Z) : K(f(Z))][f(Z)], & \dim f(Z) = \dim Z, \\ 0, & \dim f(Z) < \dim Z. \end{cases}$$

Chow Operations

Let $f : X \rightarrow Y$ be a map of sufficiently nice schemes.

- If f is flat then we have the pullback

$$f^* : Z^p(Y) \rightarrow Z^p(X), \quad [Z] \mapsto [\mathcal{O}_X \otimes_{\mathcal{O}_Y} \mathcal{O}_Z],$$

which must be interpreted in terms of support and length

- If f is proper then we have the pushforward $f_* : Z_p(X) \rightarrow Z_p(Y)$ given explicitly by

$$[Z] \mapsto \begin{cases} [K(Z) : K(f(Z))][f(Z)], & \dim f(Z) = \dim Z, \\ 0, & \dim f(Z) < \dim Z. \end{cases}$$

This can be written succinctly as $[Z] \mapsto \deg(f|_Z)[f(Z)]$.

Chow Operations

Let $f : X \rightarrow Y$ be a map of sufficiently nice schemes.

- If f is flat then we have the pullback

$$f^* : Z^p(Y) \rightarrow Z^p(X), \quad [Z] \mapsto [\mathcal{O}_X \otimes_{\mathcal{O}_Y} \mathcal{O}_Z],$$

which must be interpreted in terms of support and length

- If f is proper then we have the pushforward $f_* : Z_p(X) \rightarrow Z_p(Y)$ given explicitly by

$$[Z] \mapsto \begin{cases} [K(Z) : K(f(Z))][f(Z)], & \dim f(Z) = \dim Z, \\ 0, & \dim f(Z) < \dim Z. \end{cases}$$

This can be written succinctly as $[Z] \mapsto \deg(f|_Z)[f(Z)]$.

Both of these constructions are invariant under rational equivalence and so descend to maps of Chow groups.

Chow Operations

³Defining the exterior product over a more general base should be possible using some kind of fibral procedure.

Chow Operations

We ultimately want to grade by dimension and this affects pullbacks. If $f : X \rightarrow Y$ is smooth of constant relative dimension e then we have

$$f^* : \mathrm{CH}_*(Y) \rightarrow \mathrm{CH}_{*+e}(X), \quad [\eta] \mapsto [f^{-1}(\eta)]$$

and this construction is functorial.

³Defining the exterior product over a more general base should be possible using some kind of fibral procedure.

Chow Operations

We ultimately want to grade by dimension and this affects pullbacks. If $f : X \rightarrow Y$ is smooth of constant relative dimension e then we have

$$f^* : \mathrm{CH}_*(Y) \rightarrow \mathrm{CH}_{*+e}(X), \quad [\eta] \mapsto [f^{-1}(\eta)]$$

and this construction is functorial.

Working over k , the *exterior product*

$$\times : \mathrm{CH}_*(X) \otimes_{\mathbb{Z}} \mathrm{CH}_*(Y) \rightarrow \mathrm{CH}_*(X \times Y)$$

is given by the recipe $[Z_1] \otimes [Z_2] \mapsto [Z_1 \times Z_2]$ if k is algebraically closed and for general k in terms of the irreducible components of $Z_1 \times Z_2$.³

³Defining the exterior product over a more general base should be possible using some kind of fibral procedure.

Chow Operations

We ultimately want to grade by dimension and this affects pullbacks. If $f : X \rightarrow Y$ is smooth of constant relative dimension e then we have

$$f^* : \mathrm{CH}_*(Y) \rightarrow \mathrm{CH}_{*+e}(X), \quad [\eta] \mapsto [f^{-1}(\eta)]$$

and this construction is functorial.

Working over k , the *exterior product*

$$\times : \mathrm{CH}_*(X) \otimes_{\mathbb{Z}} \mathrm{CH}_*(Y) \rightarrow \mathrm{CH}_*(X \times Y)$$

is given by the recipe $[Z_1] \otimes [Z_2] \mapsto [Z_1 \times Z_2]$ if k is algebraically closed and for general k in terms of the irreducible components of $Z_1 \times Z_2$.³

The only thing left to describe for Chow theory is the refined Gysin pullback. This is a bit subtle and so we will come back to it later.

³Defining the exterior product over a more general base should be possible using some kind of fibral procedure.

Chow Operations

Chow Operations

The following is an important piece of inspiration.

Theorem

Chow Operations

The following is an important piece of inspiration.

Theorem

Let Sm_k denote the category of smooth k -varieties.

Chow Operations

The following is an important piece of inspiration.

Theorem

Let Sm_k denote the category of smooth k -varieties. There exists a unique contravariant graded ring structure \cdot on CH^ with the following properties.*

Chow Operations

The following is an important piece of inspiration.

Theorem

Let Sm_k denote the category of smooth k -varieties. There exists a unique contravariant graded ring structure \cdot on CH^ with the following properties.*

- 1 *Compatibility with flat pullbacks, in the sense that flat morphisms induce ring homomorphisms.*

Chow Operations

The following is an important piece of inspiration.

Theorem

Let Sm_k denote the category of smooth k -varieties. There exists a unique contravariant graded ring structure \cdot on CH^ with the following properties.*

- ① Compatibility with flat pullbacks, in the sense that flat morphisms induce ring homomorphisms.*
- ② Compatibility with the Cartier divisor intersection product $\mathrm{CH}^1(X) \times \mathrm{CH}^p(X) \rightarrow \mathrm{CH}^{p+1}(X)$ for $X \in \mathrm{Sm}_k$.*

Chow Operations

The following is an important piece of inspiration.

Theorem

Let Sm_k denote the category of smooth k -varieties. There exists a unique contravariant graded ring structure \cdot on CH^ with the following properties.*

- ① *Compatibility with flat pullbacks, in the sense that flat morphisms induce ring homomorphisms.*
- ② *Compatibility with the Cartier divisor intersection product $\mathrm{CH}^1(X) \times \mathrm{CH}^p(X) \rightarrow \mathrm{CH}^{p+1}(X)$ for $X \in \mathrm{Sm}_k$.*
- ③ *Given $X \in \mathrm{Sm}_k$ and $V, W \subseteq X$ closed integral, $[V \times_k X] \cdot [X \times_k W] = [V \times_k W]$ working with cycles on $X \times_k X$.*

Chow Operations

The following is an important piece of inspiration.

Theorem

Let Sm_k denote the category of smooth k -varieties. There exists a unique contravariant graded ring structure \cdot on CH^ with the following properties.*

- ① *Compatibility with flat pullbacks, in the sense that flat morphisms induce ring homomorphisms.*
- ② *Compatibility with the Cartier divisor intersection product $\mathrm{CH}^1(X) \times \mathrm{CH}^p(X) \rightarrow \mathrm{CH}^{p+1}(X)$ for $X \in \mathrm{Sm}_k$.*
- ③ *Given $X \in \mathrm{Sm}_k$ and $V, W \subseteq X$ closed integral, $[V \times_k X] \cdot [X \times_k W] = [V \times_k W]$ working with cycles on $X \times_k X$.*
- ④ *Projection formula: Let $f : X \rightarrow Y$ be proper with $X, Y \in \mathrm{Sm}_k$, $\alpha \in \mathrm{CH}^*(X)$, and $\beta \in \mathrm{CH}^*(Y)$. Then, $f_*(\alpha \cdot f^*(\beta)) = f_*(\alpha) \cdot \beta$.*

Chow Operations

The following is an important piece of inspiration.

Theorem

Let Sm_k denote the category of smooth k -varieties. There exists a unique contravariant graded ring structure \cdot on CH^* with the following properties.

- ① *Compatibility with flat pullbacks, in the sense that flat morphisms induce ring homomorphisms.*
- ② *Compatibility with the Cartier divisor intersection product $\mathrm{CH}^1(X) \times \mathrm{CH}^p(X) \rightarrow \mathrm{CH}^{p+1}(X)$ for $X \in \mathrm{Sm}_k$.*
- ③ *Given $X \in \mathrm{Sm}_k$ and $V, W \subseteq X$ closed integral, $[V \times_k X] \cdot [X \times_k W] = [V \times_k W]$ working with cycles on $X \times_k X$.*
- ④ *Projection formula: Let $f : X \rightarrow Y$ be proper with $X, Y \in \mathrm{Sm}_k$, $\alpha \in \mathrm{CH}^*(X)$, and $\beta \in \mathrm{CH}^*(Y)$. Then, $f_*(\alpha \cdot f^*(\beta)) = f_*(\alpha) \cdot \beta$.*
- ⑤ *Homotopy invariance: Let $p : V \rightarrow X$ be a smooth vector bundle. Then, $p^* : \mathrm{CH}^*(X) \xrightarrow{\sim} \mathrm{CH}^*(V)$.*

Returning to our intersection theory H_* for V , what are the remaining properties we want to impose?

Returning to our intersection theory H_* for V , what are the remaining properties we want to impose? First, we have the base change condition which says that if

$$\begin{array}{ccc} X & \xrightarrow{g} & Y \\ q \downarrow & & \downarrow p \\ Z & \xrightarrow{f} & W \end{array}$$

is a Cartesian square with f projective and p smooth then $p^* \circ f_* = g_* \circ q^*$.

Consider the following behemoth commutative diagram

$$\begin{array}{ccccc}
 X'' & \xrightarrow{f''} & Y'' & \longrightarrow & S \\
 g'' \downarrow & & \downarrow g' & & \downarrow g \\
 X' & \xrightarrow{f'} & Y' & \longrightarrow & T \\
 h \downarrow & & \downarrow & & \\
 X & \xrightarrow{f} & Y & &
 \end{array}$$

with all squares Cartesian.

Consider the following behemoth commutative diagram

$$\begin{array}{ccccc}
 X'' & \xrightarrow{f''} & Y'' & \longrightarrow & S \\
 g'' \downarrow & & \downarrow g' & & \downarrow g \\
 X' & \xrightarrow{f'} & Y' & \longrightarrow & T \\
 h \downarrow & & \downarrow & & \\
 X & \xrightarrow{f} & Y & &
 \end{array}$$

with all squares Cartesian. Assume that f is a regular immersion (of constant codimension c), so that we have $f^! : H_*(Y') \rightarrow H_{*-c}(X')$.

Consider the following behemoth commutative diagram

$$\begin{array}{ccccc}
 X'' & \xrightarrow{f''} & Y'' & \longrightarrow & S \\
 g'' \downarrow & & \downarrow g' & & \downarrow g \\
 X' & \xrightarrow{f'} & Y' & \longrightarrow & T \\
 h \downarrow & & \downarrow & & \\
 X & \xrightarrow{f} & Y & &
 \end{array}$$

with all squares Cartesian. Assume that f is a regular immersion (of constant codimension c), so that we have $f^! : H_*(Y') \rightarrow H_{*-c}(X')$. We place several compatibilities on $f^!$.

Consider the following behemoth commutative diagram

$$\begin{array}{ccccc}
 X'' & \xrightarrow{f''} & Y'' & \longrightarrow & S \\
 g'' \downarrow & & \downarrow g' & & \downarrow g \\
 X' & \xrightarrow{f'} & Y' & \longrightarrow & T \\
 h \downarrow & & \downarrow & & \\
 X & \xrightarrow{f} & Y & &
 \end{array}$$

with all squares Cartesian. Assume that f is a regular immersion (of constant codimension c), so that we have $f^! : H_*(Y') \rightarrow H_{*-c}(X')$. We place several compatibilities on $f^!$.

- If g' is projective (so that g'' is automatically projective) then we demand $f^! \circ g_*^! = g_*'' \circ f^!$.

Consider the following behemoth commutative diagram

$$\begin{array}{ccccc}
 X'' & \xrightarrow{f''} & Y'' & \longrightarrow & S \\
 g'' \downarrow & & \downarrow g' & & \downarrow g \\
 X' & \xrightarrow{f'} & Y' & \longrightarrow & T \\
 h \downarrow & & \downarrow & & \\
 X & \xrightarrow{f} & Y & &
 \end{array}$$

with all squares Cartesian. Assume that f is a regular immersion (of constant codimension c), so that we have $f^! : H_*(Y') \rightarrow H_{*-c}(X')$. We place several compatibilities on $f^!$.

- If g' is projective (so that g'' is automatically projective) then we demand $f^! \circ g_*^! = g''_* \circ f^!$.
- If g' is smooth (so that g'' is automatically smooth) then we demand $f^! \circ g'^* = g''^* \circ f^!$.

Consider the following behemoth commutative diagram

$$\begin{array}{ccccc}
 X'' & \xrightarrow{f''} & Y'' & \longrightarrow & S \\
 g'' \downarrow & & \downarrow g' & & \downarrow g \\
 X' & \xrightarrow{f'} & Y' & \longrightarrow & T \\
 h \downarrow & & \downarrow & & \\
 X & \xrightarrow{f} & Y & &
 \end{array}$$

with all squares Cartesian. Assume that f is a regular immersion (of constant codimension c), so that we have $f^! : H_*(Y') \rightarrow H_{*-c}(X')$. We place several compatibilities on $f^!$.

- If g' is projective (so that g'' is automatically projective) then we demand $f^! \circ g_*^! = g''_* \circ f^!$.
- If g' is smooth (so that g'' is automatically smooth) then we demand $f^! \circ g'^* = g''^* \circ f^!$.
- If g is a regular immersion then we demand $f^! \circ g^! = g^! \circ f^!$.

Suppose that f' is also a regular immersion, and let $E := h^*N_{X/Y}/N_{X'/Y'}$ be the *excess normal bundle*.

Suppose that f' is also a regular immersion, and let $E := h^* N_{X/Y} / N_{X'/Y'}$ be the *excess normal bundle*. We demand $f^! = \eta \circ f'^!$ for $\eta := 0_E^! \circ 0_{E*}$ with $0_E : X'' \rightarrow g''^* E$ the zero section. This is often called the *excess intersection formula*.

Suppose that f' is also a regular immersion, and let $E := h^* N_{X/Y} / N_{X'/Y'}$ be the *excess normal bundle*. We demand $f^! = \eta \circ f'^!$ for $\eta := 0_E^! \circ 0_{E*}$ with $0_E : X'' \rightarrow g''^* E$ the zero section. This is often called the *excess intersection formula*.

Remark

I believe this construction is closely linked to Grothendieck-Riemann-Roch but have not looked into the details. More on this later...

Suppose that f' is also a regular immersion, and let $E := h^* N_{X/Y} / N_{X'/Y'}$ be the *excess normal bundle*. We demand $f^! = \eta \circ f'^!$ for $\eta := 0_E^! \circ 0_{E*}$ with $0_E : X'' \rightarrow g''^* E$ the zero section. This is often called the *excess intersection formula*.

Remark

I believe this construction is closely linked to Grothendieck-Riemann-Roch but have not looked into the details. More on this later...

The latter construction of η is linked to other constructions. For L a line bundle on X , the zero section $0 : X \rightarrow L$ defines the *first Chern class* homomorphism

$$c_1(L) := 0^! \circ 0_* : H_*(X) \rightarrow H_{*-1}(X),$$

which you should check is well defined in the sense that 0 is a projective regular immersion.

Given $i : D = s^{-1}(0) \hookrightarrow X$ an effective Cartier divisor (for s some nonzero section of a line bundle on X), we have the *divisor intersection product*

$$D \cdot := i^! : H_*(X) \rightarrow H_{*-1}(D).$$

Given $i : D = s^{-1}(0) \hookrightarrow X$ an effective Cartier divisor (for s some nonzero section of a line bundle on X), we have the *divisor intersection product*

$$D \cdot := i^! : H_*(X) \rightarrow H_{*-1}(D).$$

Recall that this construction played a role in the earlier Chow ring structure theorem.

Given $i : D = s^{-1}(0) \hookrightarrow X$ an effective Cartier divisor (for s some nonzero section of a line bundle on X), we have the *divisor intersection product*

$$D \cdot := i^! : H_*(X) \rightarrow H_{*-1}(D).$$

Recall that this construction played a role in the earlier Chow ring structure theorem. Intersections with Cartier divisors provide a rich supply of geometric information, and historically were studied first in the quest to build an algebraic intersection theory.

Given $i : D = s^{-1}(0) \hookrightarrow X$ an effective Cartier divisor (for s some nonzero section of a line bundle on X), we have the *divisor intersection product*

$$D \cdot := i^! : H_*(X) \rightarrow H_{*-1}(D).$$

Recall that this construction played a role in the earlier Chow ring structure theorem. Intersections with Cartier divisors provide a rich supply of geometric information, and historically were studied first in the quest to build an algebraic intersection theory.

Remark

I really should provide example computations for this and subsequent constructions, but alas!

Given $i : D = s^{-1}(0) \hookrightarrow X$ an effective Cartier divisor (for s some nonzero section of a line bundle on X), we have the *divisor intersection product*

$$D \cdot := i^! : H_*(X) \rightarrow H_{*-1}(D).$$

Recall that this construction played a role in the earlier Chow ring structure theorem. Intersections with Cartier divisors provide a rich supply of geometric information, and historically were studied first in the quest to build an algebraic intersection theory.

Remark

I really should provide example computations for this and subsequent constructions, but alas!

Let's round out our discussion of properties of H_* .

- *Excision sequence*: Given a closed immersion $i : Z \hookrightarrow X$ with complement $j : U \hookrightarrow X$, we demand that the sequence

$$H_*(Z) \xrightarrow{i_*} H_*(X) \xrightarrow{j^*} H_*(U) \longrightarrow 0$$

is exact.

- *Excision sequence*: Given a closed immersion $i : Z \hookrightarrow X$ with complement $j : U \hookrightarrow X$, we demand that the sequence

$$H_*(Z) \xrightarrow{i_*} H_*(X) \xrightarrow{j^*} H_*(U) \longrightarrow 0$$

is exact.

- *Extended homotopy*: Given $E \in \text{Vect}_r(X)$ and $p : V \rightarrow X$ an E -torsor, we demand $p^* : H_*(X) \xrightarrow{\sim} H_{*+r}(V)$.

- *Excision sequence*: Given a closed immersion $i : Z \hookrightarrow X$ with complement $j : U \hookrightarrow X$, we demand that the sequence

$$H_*(Z) \xrightarrow{i_*} H_*(X) \xrightarrow{j^*} H_*(U) \longrightarrow 0$$

is exact.

- *Extended homotopy*: Given $E \in \text{Vect}_r(X)$ and $p : V \rightarrow X$ an E -torsor, we demand $p^* : H_*(X) \xrightarrow{\sim} H_{*+r}(V)$.
- *Projective bundle formula*: Let $E \in \text{Vect}_r(X)$ with associated projective bundle $p : \mathbb{P}(E) \rightarrow X$.

- *Excision sequence*: Given a closed immersion $i : Z \hookrightarrow X$ with complement $j : U \hookrightarrow X$, we demand that the sequence

$$H_*(Z) \xrightarrow{i_*} H_*(X) \xrightarrow{j^*} H_*(U) \longrightarrow 0$$

is exact.


- *Extended homotopy*: Given $E \in \text{Vect}_r(X)$ and $p : V \rightarrow X$ an E -torsor, we demand $p^* : H_*(X) \xrightarrow{\sim} H_{*+r}(V)$.
- *Projective bundle formula*: Let $E \in \text{Vect}_r(X)$ with associated projective bundle $p : \mathbb{P}(E) \rightarrow X$. We demand that the induced map

$$\bigoplus_{i=0}^{r-1} H_{*-r+1+i}(X) \rightarrow H_*(\mathbb{P}(E)), \quad (\xi_i) \mapsto \sum_i c_1(\mathcal{O}_{\mathbb{P}(E)}(1))^i \cdot (p^* \xi_i)$$

is an isomorphism.

⁴This should remind you of the theory of quasi-smooth morphisms.


- Let $f : X \rightarrow Y$ be an lci morphism of constant relative dimension d and choose a factorization $f = h \circ g$ with $g : X \rightarrow Z$ a regular closed immersion and $h : Z \rightarrow Y$ smooth.

⁴This should remind you of the theory of quasi-smooth morphisms. 

- Let $f : X \rightarrow Y$ be an lci morphism of constant relative dimension d and choose a factorization $f = h \circ g$ with $g : X \rightarrow Z$ a regular closed immersion and $h : Z \rightarrow Y$ smooth. We demand that the *lci pullback*

$$f^* := g^! \circ h^* : H_*(Y) \rightarrow H_{*+d}(X)$$

is independent of the choice of factorization.⁴


⁴This should remind you of the theory of quasi-smooth morphisms. 

- Let $f : X \rightarrow Y$ be an lci morphism of constant relative dimension d and choose a factorization $f = h \circ g$ with $g : X \rightarrow Z$ a regular closed immersion and $h : Z \rightarrow Y$ smooth. We demand that the *lci pullback*

$$f^* := g^! \circ h^* : H_*(Y) \rightarrow H_{*+d}(X)$$

is independent of the choice of factorization.⁴

- *Detection by smooth schemes*: Fixing $X \in \text{QSch}_k$, consider the pushforwards f_* associated to projective $f : Y \rightarrow X$ with Y smooth and quasi-projective over k .


⁴This should remind you of the theory of quasi-smooth morphisms. 

- Let $f : X \rightarrow Y$ be an lci morphism of constant relative dimension d and choose a factorization $f = h \circ g$ with $g : X \rightarrow Z$ a regular closed immersion and $h : Z \rightarrow Y$ smooth. We demand that the *lci pullback*

$$f^* := g^! \circ h^* : H_*(Y) \rightarrow H_{*+d}(X)$$

is independent of the choice of factorization.⁴

- Detection by smooth schemes:* Fixing $X \in \text{QSch}_k$, consider the pushforwards f_* associated to projective $f : Y \rightarrow X$ with Y smooth and quasi-projective over k . Then, there is a natural map $\varinjlim_{Y \rightarrow X} H_*(Y) \rightarrow H_*(X)$ for $Y \rightarrow X$ as above, with transition maps arising from projective morphisms $Y \rightarrow Y'$ over X .

⁴This should remind you of the theory of quasi-smooth morphisms. 

- Let $f : X \rightarrow Y$ be an lci morphism of constant relative dimension d and choose a factorization $f = h \circ g$ with $g : X \rightarrow Z$ a regular closed immersion and $h : Z \rightarrow Y$ smooth. We demand that the *lci pullback*

$$f^* := g^! \circ h^* : H_*(Y) \rightarrow H_{*+d}(X)$$

is independent of the choice of factorization.⁴

- Detection by smooth schemes:* Fixing $X \in \mathbf{QSch}_k$, consider the pushforwards f_* associated to projective $f : Y \rightarrow X$ with Y smooth and quasi-projective over k . Then, there is a natural map $\varinjlim_{Y \rightarrow X} H_*(Y) \rightarrow H_*(X)$ for $Y \rightarrow X$ as above, with transition maps arising from projective morphisms $Y \rightarrow Y'$ over X . We demand that this natural map out of the colimit is an isomorphism.

⁴This should remind you of the theory of quasi-smooth morphisms. 