
📝 Unit 7 Practice 2 — Data Analysis

Complete independently · NC.8.SP.1–4 · Show all work

Name _____ **Date** _____ **Period** _____

Remember: Describe a scatter plot: Up or down? Tight or spread out? Any outliers?


Question 1 Score 3



Looking at the **non-outlier** points, what correlation is shown?

- ○ Negative correlation
- ○ Positive correlation
- ○ No correlation
- ○ Perfect correlation

Question 2 Score 3



In a scatterplot for "temperature vs. hot coffee sales," which type of correlation would you expect?

- ○ Strong positive
- ○ Strong negative
- ○ No correlation
- ○ Perfect positive


Question 3 Score 4



This scatterplot shows athletes' skill level as weeks without practice increase. Describe the full correlation.

- ○ Weak positive correlation — practice doesn't matter
- ○ Strong negative — the longer they skip practice, the lower their skill
- ○ No correlation — the points are scattered
- ○ Moderate positive — skill improves with rest

Question 4 Score 4



Student scores a 95 on a test after studying 8 hours. The trend line predicts 87 points at 8 hours. This student's score above the line makes it:

- ○ A point above the trend line (the line under-predicts this student)
- ○ An outlier below the trend line
- ○ The y-intercept
- ○ The maximum score


Question 5 Score 4



A trend line is described as "strong and positive." What does each word mean?

- ○ Strong = steep slope · Positive = y is positive
- ○ Strong = points cluster tightly near the line · Positive = trend goes upward
- ○ Strong = many data points · Positive = all y-values are positive
- ○ Strong = the line is long · Positive = x values increase

Question 6 Score 5




Both go up in summer. Is one causing the other, or is something else behind both?

Ice cream sales and pool visits both increase in summer. A student says "ice cream causes people to visit pools." What concept explains the real relationship?

- ○ Causation — ice cream makes people want to swim
- ○ Correlation does not imply causation — both are caused by hot weather
- ○ The data must be wrong
- ○ There is no correlation

Interpret in context: slope = "per unit increase in x" · y-intercept = "when x = 0"


Question 1 Score 3



The model is y = 8x + 30 (x = training weeks, y = max push-ups in 1 min). Find the green dot: how many push-ups after **5 weeks**?

- ○ 38 push-ups
- ○ 55 push-ups
- ○ 70 push-ups
- ○ 80 push-ups


Question 2 Score 3



In y = 8x + 30 (training weeks vs. push-ups), interpret the slope of 8.

- ○ The athlete starts at 8 push-ups
- ○ Training takes 8 weeks total
- ○ For each additional week of training, push-up count increases by about 8
- ○ The slope means 8 athletes were tested


Question 3 Score 4



Interpret the y-intercept of 30 in the same model (y = 8x + 30).

- ○ 30 is the maximum push-up count
- ○ It takes 30 weeks to improve
- ○ At 0 weeks of training, a person can do about 30 push-ups
- ○ 30 people were surveyed

Question 4 Score 4



Temperature vs. electric heating bills: y = −2x + 120 (x = °F above 50, y = bill in $). What does slope −2 mean?

- ○ The bill increases $2 per degree above 50°F
- ○ The bill decreases by $2
- ○ For each degree above 50°F, the bill
- ○ The starting bill is −$2
- ○ The bill is always negative


Question 5 Score 4



Model: y = 4x + 8 (x = hours tutoring, y = points gained). The data was collected for x = 1 to x = 6. Is x = 10 inside or outside the data? Is that prediction safe?

- ○ Inside the data — safe
- ○ Inside the data — less safe
- ○ Outside the data — less safe
- ○ Outside the data — safe

Question 6 Score 5



Using y = 4x + 8 from Q5, which is the most reliable prediction?

- ○ x = 0 hours (no data there)
- ○ x = 20 hours (beyond data)
- ○ x = 4 hours (within the data range)
- ○ x = −2 hours (impossible)

📋 Practice 2 Two-Way Table

120 8th grade students surveyed on sport preference split by **lunch period**:

	Basketball 🏀	Soccer ⚽	Other 🏃	Total
1st Lunch	24	**18**	18	**60**
2nd Lunch	30	12	18	**60**
Total	**54**	**30**	**36**	**120**

Question 1 Score 3

	Basket-ball	Soccer	Other	Total
1st Lunch	24	18	18	60
2nd Lunch	30	12	18	60
Total	54	30	36	120

Find where the 1st Lunch row and the Soccer column cross.

What is the joint frequency of 1st lunch students who prefer Soccer?

- ○ 30
- ○ 12
- ○ 18
- ○ 60

Question 2 Score 3

	Basket-ball	Soccer	Other	Total
1st Lunch	24	18	18	60
2nd Lunch	30	12	18	60
Total	54	30	36	120

Marginal = the total at the edge of the Basketball column.

What is the marginal frequency for Basketball?

- ○ 24
- ○ 54
- ○ 30
- ○ 120

Question 3 Score 4

	Basket-ball	Soccer	Other	Total
1st Lunch	24	18	18	60
2nd Lunch	30	12	18	60
Total	54	30	36	120

1st Lunch: $\frac{24}{60}$ = ?

What percent of 1st lunch students prefer Basketball?

- ○ 40%
- ○ 20%
- ○ 45%
- ○ 50%

Question 4 Score 4

	Basket-ball	Soccer	Other	Total
1st Lunch	24	18	18	60
2nd Lunch	30	12	18	60
Total	54	30	36	120

2nd Lunch: $\frac{30}{60}$ = ?

What percent of 2nd lunch students prefer Basketball?

- ○ 40%
- ○ 45%
- ○ 50%
- ○ 25%

Question 5 Score 4


	Basket-ball	Soccer	Other	Total
1st Lunch	24	18	18	60
2nd Lunch	30	12	18	60
Total	54	30	36	120

Soccer: $\frac{30}{120} = ?$

What is the relative frequency of students who prefer Soccer (out of 120)?

- ○ 25%
- ○ 15%
- ○ 30%
- ○ 18%

Question 6 Score 5



Based on the basketball percentages (40% vs 50%), is there an association?

- ○ No — the totals are both 60
- ○ No — 40 and 50 are close enough
- ○ Yes — 2nd lunch students prefer basketball more (50% vs 40%)
- ○ Yes — but only if we add more data

Question 7 Score 5

	Basket-ball	Soccer	Other	Total
1st Lunch	24	18	18	60
2nd Lunch	30	12	18	60
Total	54	30	36	120


Everyone who picked Other, out of all 120 students.

If a student is chosen randomly from the 120 surveyed, what is the probability they prefer "Other" sports?

- ○ $^{36}/_{90} = 40\%$
- ○ $^{36}/_{120} = 30\%$
- ○ $^{36}/_{60} = 60\%$
- ○ $^{18}/_{120} = 15\%$

Extended thinking: These questions require connecting multiple concepts. Take your time and show your reasoning.


Question 1 Score 4



The line y = −3.5x + 75 models temperature (°F) at altitude x (in 1,000s of feet). At 8,000 ft (x = 8), predict the temperature.

- ○ 47°F
- ○ 28°F
- ○ 75°F
- ○ 47°C


Question 2 Score 4



Line of best fit: y = 12x + 5. If x increases by 3 units, by how much does y increase?

- ○ 12
- ○ 36
- ○ 15
- ○ 5

Question 3 Score 5



Using the same model (y = −3.5x + 75): at what altitude does the model predict 0°F? Round to the nearest 1,000 ft.

- ○ x ≈ 21,000 ft
- ○ x ≈ 15,000 ft
- ○ x ≈ 75,000 ft
- ○ x ≈ 10,000 ft

Question 4 Score 5



Two researchers study "firefighters at scene vs. fire damage." They find a strong positive correlation. Researcher A says: "More firefighters cause MORE damage." Researcher B says: "More firefighters are sent to BIGGER fires, which also cause more damage." Who is correct?

- ○ Researcher A — the data shows clear causation
- ○ Researcher B — larger fires need more firefighters AND cause more damage (lurking variable)
- ○ Both are wrong — there is no correlation
- ○ Both are correct — it depends on the day

📝 Worked Examples

Study these step-by-step solutions before attempting the problems above.

Two-way table: 45 boys like pizza out of 120 total

Step 1: Joint relative freq

```
45 ÷ 120 = 0.375 = 37.5%
```

	Like pizza	Do not like	Total
Boys	45	15	60
Girls			60
Total			120

Joint: $\frac{45}{120} = 37.5\%$

Step 2: Conditional: of 60 boys

```
45 ÷ 60 = 0.75 = 75% of boys like pizza
```


	Like pizza	Do not like	Total
Boys	45	15	60
Girls			60
Total			120

Of boys: $\frac{45}{60} = 75\%$

🔍 Common Mistakes

Can you spot what went wrong?

Correlation: points go up-left to down-right



✗ **Student:** Student says: positive correlation

✓ **Fix:** WRONG! Downhill = NEGATIVE correlation. Uphill = positive.

💡 *Think of it like a hill: going up = positive, going down = negative.*

⚡ Key Tips for This Unit

✅ Positive: both increase. Negative: one up, other down. None: random.

✅ Line of best fit: approx equal points above and below.

✅ Outlier: far from the pattern. Can distort the trend line.

✅ Joint RF = cell÷grand total. Conditional RF = cell÷row or column total.