

Assignment 2

1.1 Are either $\lceil \lg n \rceil!$ or $\lceil \lg \lg n \rceil!$ polynomially bounded?

Polynomially bounded means $f_n = O(n^k)$ for some constant k (e.g., whether $f_n \leq c \cdot n^k$ for constants c and k as n approaches ∞). For the first function $\lceil \lg n \rceil!$, without loss of generality, assume $n = 2^a$ (where $a \in \mathbb{N}$).

$$\begin{aligned}\lceil \lg n \rceil! &\leq c \cdot n^k \\ \lg(2^a)! &\leq c \cdot (2^a)^k \\ a! &\leq c \cdot 2^{ak}\end{aligned}$$

The statement $a! \leq c \cdot 2^{ak}$ is a contradiction, as the factorial function $a!$ is not exponentially bounded. Therefore, $\lceil \lg n \rceil!$ is not polynomially bounded (via proof by contradiction). For the second function $\lceil \lg \lg n \rceil!$, without loss of generality, assume $n = 2^{2^a}$ (where $a \in \mathbb{N}$).

$$\begin{aligned}\lceil \lg \lg n \rceil! &\leq c \cdot n^k \\ \lg \lg(2^{2^a})! &\leq c \cdot (2^{2^a})^k \\ a! &\leq c \cdot 2^{k \cdot 2^a} \\ 1 \cdot 2 \cdot 3 \cdots a &\leq c \cdot (2^{2k} \cdot 2^{4k} \cdot 2^{8k} \cdots 2^{2^a \cdot k})\end{aligned}$$

The statement $1 \cdot 2 \cdot 3 \cdots a \leq c \cdot (2^{2k} \cdot 2^{4k} \cdot 2^{8k} \cdots 2^{2^a \cdot k})$ is obviously true. Therefore $\lceil \lg \lg n \rceil!$ is polynomially bounded (via direct proof).

- 1.2 Use induction to prove $F_i = \frac{\phi^i - \hat{\phi}^i}{\sqrt{5}}$; where $F_i = F_{i-2} + F_{i-1}$, and ϕ is the golden ratio $\frac{1+\sqrt{5}}{2}$.
- 1.3 Show that $k \lg k = \Theta(n)$ implies $k = \Theta\left(\frac{n}{n \ln n}\right)$.
- 1.4 Are either 2^{n+1} or 2^{2n} big- O of 2^n ?
- 1.5 For each pair of functions (A, B) , indicate whether A is O, o, Ω, ω , or Θ of B . Assume $k \geq 1$, $\epsilon > 0$, $c > 1$ are constants.

	A	B	O	o	Ω	ω	Θ
a.	$\lg^k n$	n^ϵ	yes	yes			
b.	n^k	c^n	yes	yes			
c.	\sqrt{n}	$n^{\sin n}$					
d.	2^n	$2^{n/2}$			yes	yes	
e.	$n^{\lg c}$	$c^{\lg n}$	yes		yes		yes
f.	$\lg(n!)$	$\lg(n^n)$	yes		yes		yes

1.6 Order the following functions such that $f_1 = \Omega(f_2)$, $f_2 = \Omega(f_3)$, ..., $f_{29} = \Omega(f_{30})$, and partition them into equivalence classes such that each function is big- Θ of each other.

$$\begin{aligned}
f_1 &= 2^{2^{n+1}} = \Omega(2^{2^n}), & n \lg n &= \Omega(2^{\lg n}), \\
f_2 &= 2^{2^n} = \Omega((n+1)!), & 2^{\lg n} &= \Omega(n), \\
f_3 &= (n+1)! = \Omega(n!), & n &= \Omega\left((\sqrt{2})^{\lg n}\right), \\
f_4 &= n! = \Omega(e^n), & (\sqrt{2})^{\lg n} &= \Omega(\sqrt{n}), \\
f_5 &= e^n = \Omega(n \cdot 2^n), & \sqrt{n} &= \Omega(2^{\sqrt{2} \lg n}), \\
f_6 &= n \cdot 2^n = \Omega(2^n), & 2^{\sqrt{2} \lg n} &= \Omega(\lg^2 n), \\
f_7 &= 2^n = \Omega\left(\left(\frac{3}{2}\right)^n\right), & \lg^2 n &= \Omega(\ln n), \\
f_8 &= \left(\frac{3}{2}\right)^n = \Omega(n^{\lg \lg n}), & \ln n &= \Omega(\sqrt{\lg n}), \\
f_9 &= n^{\lg \lg n} = \Omega((\lg n)^{\lg n}), & \sqrt{\lg n} &= \Omega(\ln \ln n), \\
f_{10} &= (\lg n)^{\lg n} = \Omega((\lg n)!), & \ln \ln n &= \Omega(2^{\lg^* n}), \\
f_{11} &= (\lg n)! = \Omega(N^3), & 2^{\lg^* n} &= \Omega(\lg^* n), \\
f_{12} &= N^3 = \Omega(n^2), & \lg^* n &= \Omega(\lg^*(\lg n)), \\
f_{13} &= n^2 = \Omega(4^{\lg n}), & \lg^*(\lg n) &= \Omega(\lg(\lg^* n)), \\
f_{14} &= 4^{\lg n} = \Omega(\lg(n!)), & \lg(\lg^* n) &= \Omega\left(n^{\frac{1}{\lg n}}\right), \\
f_{15} &= \lg(n!) = \Omega(n \lg n), & n^{\frac{1}{\lg n}} &
\end{aligned}$$