

# Assignment 2

## 1.1 Are either $\lceil \lg n \rceil!$ or $\lceil \lg \lg n \rceil!$ polynomially bounded?

Polynomially bounded means  $f_n = O(n^k)$  for some constant  $k$  (e.g., whether  $f_n \leq c \cdot n^k$  for constants  $c$  and  $k$  as  $n$  approaches  $\infty$ ).

For the first function  $\lceil \lg n \rceil!$ , without loss of generality, assume  $n = 2^a$  (where  $a \in \mathbb{N}$ ). The statement  $a! \leq c \cdot 2^{ak}$  (see below) is a contradiction, as the factorial function  $a!$  is not exponentially bounded. Therefore,  $\lceil \lg n \rceil!$  is not polynomially bounded (via proof by contradiction).

$$\lceil \lg n \rceil! \leq c \cdot n^k$$

$$\lg(2^a)! \leq c \cdot (2^a)^k$$

$$a! \leq c \cdot 2^{ak}$$

For the second function  $\lceil \lg \lg n \rceil!$ , without loss of generality, assume  $n = 2^{2^a}$  (where  $a \in \mathbb{N}$ ). The statement  $1 \cdot 2 \cdot 3 \cdots a \leq c \cdot (2^{2k} \cdot 2^{4k} \cdot 2^{8k} \cdots 2^{2^a k})$  (see below) is obviously true. Therefore  $\lceil \lg \lg n \rceil!$  is polynomially bounded (via direct proof).

$$\lceil \lg \lg n \rceil! \leq c \cdot n^k$$

$$\lg \lg(2^{2^a})! \leq c \cdot (2^{2^a})^k$$

$$a! \leq c \cdot 2^{k \cdot 2^a}$$

$$1 \cdot 2 \cdot 3 \cdots a \leq c \cdot (2^{2k} \cdot 2^{4k} \cdot 2^{8k} \cdots 2^{2^a k})$$

## 1.2 Use induction to prove $F_i = \frac{\phi^i - \hat{\phi}^i}{\sqrt{5}}$ , where $F_i = F_{i-2} + F_{i-1}$ , and $\phi$ is the golden ratio $\frac{1+\sqrt{5}}{2}$ .

To prove by induction, write out the expressions  $f_n$  and  $f_{n+1}$  (note:  $f_{n+1}$  is the same as  $f_n$ , but with  $(n+1)$  substituted everywhere in place of  $n$ ). Next, if applicable, re-write the expression  $f_{n+1}$  in terms of  $f_n$  then perform algebraic manipulations on the expression until you reach some variation of  $f_{n+1} = f_{n+1}$ . Lastly, show that the expression  $f_c$  also holds for some

constant  $c$ . The algebra is called "the inductive step", and the calculation for on the constant is called "the base case".

In this problem, the expression to prove is  $F_i = \frac{\phi^i - \hat{\phi}^i}{\sqrt{5}}$ , where  $\phi = \frac{1+\sqrt{5}}{2}$ . Start by demonstrating the expression holds for constants  $c = 0$  and  $c = 1$ .

$$F_0 = \frac{\phi^0 - \hat{\phi}^0}{\sqrt{5}} = \frac{1 - 1}{\sqrt{5}} = 0$$

$$F_1 = \frac{\phi^1 - \hat{\phi}^1}{\sqrt{5}} = \frac{\frac{1+\sqrt{5}}{2} - \frac{1-\sqrt{5}}{2}}{\sqrt{5}} = \frac{(1+\sqrt{5}) - (1-\sqrt{5})}{2\sqrt{5}} = 1$$

After showing the expression holds for some base cases  $F_0$  and  $F_1$ , the next step is algebra. Setup the expression  $F_n$  in terms of  $F_{n-1}$ , then solve (see below).

$$F_i = \frac{\phi^i - \hat{\phi}^i}{\sqrt{5}} = F_{i-1} + F_{i-2} \quad F_{i-1} = \frac{\phi^{i-1} - \hat{\phi}^{i-1}}{\sqrt{5}}$$

$$F_i = F_{i-1} + F_{i-2}$$

$$\begin{aligned} \frac{\phi^i - \hat{\phi}^i}{\sqrt{5}} &= \frac{\phi^{i-1} - \hat{\phi}^{i-1}}{\sqrt{5}} + \frac{\phi^{i-2} - \hat{\phi}^{i-2}}{\sqrt{5}} \\ &= \frac{\phi^{i-1} + \hat{\phi}^{i-1} - \phi^{i-2} - \hat{\phi}^{i-2}}{\sqrt{5}} \\ &= \frac{\phi^{i-1} - \phi^{i-2} + \hat{\phi}^{i-1} - \hat{\phi}^{i-2}}{\sqrt{5}} \\ &= \frac{[(\phi \cdot \phi^{i-2}) + \phi^{i-2}] - [(\hat{\phi} \cdot \hat{\phi}^{i-2}) + \hat{\phi}^{i-2}]}{\sqrt{5}} \\ &= \frac{\phi^{i-2}(\phi + 1) - \hat{\phi}^{i-2}(\hat{\phi} + 1)}{\sqrt{5}} \\ &= \frac{\phi^{i-2}(\phi^2) - \hat{\phi}^{i-2}(\hat{\phi}^2)}{\sqrt{5}} \\ &= \frac{\phi^i - \hat{\phi}^i}{\sqrt{5}} \end{aligned}$$

Since we have shown  $F_i$  is obtainable via  $F_{i-1}$ , we have completed the inductive step. Since both the inductive step and base cases have been shown, the proof by induction is complete.