Unit 1: Introduction to data

Data Collection +
 Observational studies & experiments

Sta 104 - Summer 2018, Term 1

Duke University, Department of Statistical Science

Prof. White

Slides posted at https://www2.stat.duke.edu/courses/Summer18/sta104.001-1/

Sampling is natural



- ▶ When you taste a spoonful of soup and decide the spoonful you tasted isn't salty enough, that's *exploratory analysis*
- ▶ If you generalize and conclude that your entire soup needs salt, that's an *inference*
- ► For your inference to be valid, the spoonful you tasted (the sample) needs to be *representative* of the entire pot (the population)

▶ Ultimate goal: make inferences about populations

- ▶ Caveat: populations are difficult or impossible to access
- ➤ Solution: use a sample from that population, and use *statistics* from that sample to make inferences about the unknown population *parameters*
- ► The better (more *representative*) sample we have, the more reliable our estimates and more accurate our inferences will be

Suppose we want to know how many offspring female lemurs have, on average. It's not feasible to obtain offspring data from on all female lemurs, so we use data from the Duke Lemur Center. We use the sample mean from these data as an estimate for the unknown population mean. Can you see any limitations to using data from the Duke Lemur Center to make inferences about all lemurs?


2. Ideally use a simple random sample, stratify to control for a variable, and cluster to make sampling easier

Simple random:


Drawing names from a hat



Stratified: homogenous strata Stratify to control for SES




Cluster: heterogenous clusters Sample all chosen clusters



Multistage:

Random sample in chosen clusters



2

Clicker question

A city council has requested a household survey be conducted in a suburban area of their city. The area is broken into many distinct and unique neighborhoods, some including large homes, some with only apartments, and others a diverse mixture of housing structures. Which approach would likely be the least effective?

- Simple random sampling
- (b) Stratified sampling, where each stratum is a neighborhood
- © Cluster sampling, where each cluster is a neighborhood

- ► Non-response: If only a small fraction of the randomly sampled people choose to respond to a survey, the sample may no longer be representative of the population
- Voluntary response: Occurs when the sample consists of people who volunteer to respond because they have strong opinions on the issue since such a sample will also not be representative of the population
- ► Convenience sample: Individuals who are easily accessible are more likely to be included in the sample

4

5

Clicker question

A school district is considering whether it will no longer allow high school students to park at school after two recent accidents where students were severely injured. As a first step, they survey parents by mail, asking them whether or not the parents would object to this policy change. Of 6,000 surveys that go out, 1,200 are returned. Of these 1,200 surveys that were completed, 960 agreed with the policy change and 240 disagreed. Which of the following statements are true?

- Some of the mailings may have never reached the parents.
- Overall, the school district has strong support from parents to move forward with the policy approval.
- It is possible that majority of the parents of high school students disagree with the policy change.
- The survey results are unlikely to be biased because all parents were mailed a survey.
- a Only I
- (b) I and II
- I and III
- d III and
- Only IV

nd (e) Only

What type of study is this? What is the scope of inference (causality / generalizability)?

Facebook Tinkers With Users' Emotions in News Feed Experiment, Stirring Outcry

By VINDU GOEL JUNE 29, 2014

The New Hork Times

In an academic paper published in conjunction with two university researchers, the company reported that, for one week in January 2012, it had altered the number of positive and negative posts in the news feeds of 689,003 randomly selected users to see what effect the changes had on the tone of the posts the recipients then wrote.

The researchers found that moods were contagious. The people who saw more positive posts responded by writing more positive posts. Similarly, seeing more negative content prompted the viewers to be more negative in their own posts.

A study that surveyed a random sample of otherwise healthy adults found that people are more likely to get muscle cramps when they're stressed. The study also noted that people drink more coffee and sleep less when they're stressed. What type of study is this?

What is the conclusion of the study?

Can this study be used to conclude a causal relationship between increased stress and muscle cramps?

- ▶ We would like to design an experiment to investigate if increased stress causes muscle cramps:
 - Treatment: increased stress
 - Control: no or baseline stress
- ▶ It is suspected that the effect of stress might be different on younger and older people: *block* for age.

Why is this important? Can you think of other variables to block for?

8

9

11

6. Random sampling helps generalizability, random assignment helps causality

most ideal No random Random observational experiment assignment assignment studies No causal conclusion, Causal conclusion. Random correlation statement Generalizability generalized to the whole sampling generalized to the whole population. population. No random No causal conclusion, No Causal conclusion, correlation statement only only for the sample. generalizability sampling for the sample. bad most Causation Correlation observational experiments studies

Summary of main ideas

- 1. ??
- 2. ??
- 3. ??
- 4. ??
- 5. ??
- 6. ??

10