


ENOSF1 ENOSF1 chr18:670318-674406 · LIHC, LUAD and KIRC · OS

ENOSF1 chr18:670318-674406: exon chr18:670,318–674,406 (4,089 nt); HIT index from −1 (used as a first exon) to 1 (used as a last exon), about 0 for an internal exon · minus strand




LIHC







Cox model · 342 patients, 115 events · HIT index q 0.014 *

HIT index + host expression + age + sex + stage (stage Stage III, Stage IIIB, Stage IIIC, Stage IV, Stage IVA, Stage IVB merged into Stage I (24 patients, 13 events); stage baseline I absent; Stage I+Stage III+Stage IIIB+Stage IIIC+Stage …




LUAD







Cox model · 483 patients, 174 events · HIT index q 0.038 *

HIT index + host expression + age + sex + stage (stage Stage I, Stage II merged into Stage IB (6 patients, 2 events); stage baseline I absent; Stage IB+Stage I+Stage II used; non-proportional hazards: stage Stage IIA (p 0.012) …




KIRC







Cox model · 517 patients, 171 events · HIT index q 0.063

HIT index + host expression + age + sex + stage (stage baseline I absent; Stage I used; non-proportional hazards: stage Stage III (p 0.003), stage Stage IV (p 3.1e-4))



Cox models: Numeric covariates per SD of the fit cohort; categories against the level named.





Exon · Region measured by HIT index · Higher in tumour · Lower in tumour · KM split · Paired Δ · Unpaired Δ · Cox p < 0.05 · Cox p ≥ 0.05 · Cohort shown at left · Model term p < 0.05 · p ≥ 0.05 · * q < 0.05 (Benjamini–Hochberg) · † non-proportional hazards (p < 0.05) · ‡ fewer than 20 events: low power

q: Benjamini–Hochberg within ENOSF1 for OS, one family per kind of test over all its HIT-index events × cohorts (within patients 13, all samples 15, KM 14, Cox, forest 14, model rows 14 tests); none for families under 10.