

ENOSF1 expression · LIHC, LUAD and KIRC · OS


The forest shows ENOSF1 expression in every cohort with a test; the rows below show the cohorts drawn on its splicing pages



Host-gene expression · ENOSF1 itself: tumour vs normal, a KM split at its median, and Cox on expression + age + sex + stage

LIHC · ENOSF1 expression







Cox model · 343 patients, 116 events


expression + age + sex + stage (stage Stage III, Stage IIIB, Stage IIIC, Stage IV, Stage IVA, Stage IVB merged into Stage I (25 patients, 14 events); stage baseline I absent; Stage I+Stage III+Stage IIIB+Stage IIIC+Stage IV+Stage …

	HR (95% CI)	p
Expr. (per SD)	0.97 (0.81–1.17)	0.77
age (per SD)	1.10 (0.91–1.33)	0.33
sex (vs female)		
male	0.77 (0.53–1.13)	0.18
stage (vs Stage I+Stage III+Stage IIIB+Stage IIIC+Stage IV+Stage IVA+Stage IVB)		
Stage II	1.14 (0.71–1.82)	0.59
Stage IIIA	2.29 (1.49–3.51)	1.5e-4



LUAD · ENOSF1 expression






Cox model · 483 patients, 174 events


expression + age + sex + stage (stage Stage I, Stage II merged into Stage IB (6 patients, 2 events); stage baseline I absent; Stage IB+Stage I+Stage II used; non-proportional hazards: sex male (p 0.038), stage Stage IIA (p 0.018), stage …

	HR (95% CI)	p
Expr. (per SD)	0.93 (0.81–1.08)	0.37
age (per SD)	1.14 (0.97–1.33)	0.11
sex (vs female)		
male	0.94 (0.69–1.27)	0.68 †
stage (vs Stage IB+Stage I+Stage II)		
Stage IA	0.91 (0.56–1.50)	0.72
Stage IIA	2.91 (1.67–5.05)	1.5e-4 †
Stage IIB	2.16 (1.34–3.49)	0.002
Stage IIIA	3.49 (2.19–5.59)	1.8e-7 †
Stage IIIB	2.37 (0.99–5.67)	0.053 †
Stage IV	4.07 (2.23–7.44)	4.9e-6



KIRC · ENOSF1 expression






Cox model · 519 patients, 171 events

expression + age + sex + stage (stage baseline I absent; Stage I used; non-proportional hazards: stage Stage III (p 0.003), stage Stage IV (p 2.6e-4))

	HR (95% CI)	p
Expr. (per SD)	1.03 (0.88–1.20)	0.74
age (per SD)	1.48 (1.24–1.77)	1.5e-5
sex (vs female)		
male	0.94 (0.68–1.30)	0.71
stage (vs Stage I)		
Stage II	1.33 (0.71–2.47)	0.37
Stage III	2.58 (1.71–3.90)	6.2e-6 †
Stage IV	6.92 (4.69–10.21)	1.9e-22 †



Higher in tumour · Lower in tumour · KM split · Paired Δ · Unpaired Δ · Cox p < 0.05 · Cox p ≥ 0.05
Cohort drawn below · Model term p < 0.05 · p ≥ 0.05 · † non-proportional hazards (p < 0.05) · ‡ fewer than 20 events: low power

Host-gene expression of ENOSF1 on the expression table's scale; not adjusted for multiple testing. A tumour–normal hit needs p < 0.05 and |Δ median| > 1. The HR is per SD of expression in each cohort, with age, sex, stage.