


GLIPR1 GLIPR1 chr12:75498694-75503863 · BRCA, STAD and COAD · OS

GLIPR1 chr12:75498694-75503863: exon chr12:75,498,694–75,503,863 (5,170 nt); HIT index from −1 (used as a first exon) to 1 (used as a last exon), about 0 for an internal exon · plus strand




BRCA







Cox model · 1062 patients, 146 events · HIT index q 0.11

HIT index + host expression + age + sex + stage (stage Stage IB, Stage II, Stage III merged into Stage IIA (14 patients, 2 events); stage baseline I absent; Stage IIA+Stage IB+Stage II+Stage III used; non-proportional hazards: stage …




STAD







Cox model · 335 patients, 134 events · HIT index q 0.11

HIT index + host expression + age + sex + stage (stage Stage I, Stage III merged into Stage IIIA (4 patients, 3 events); stage baseline I absent; Stage IIIA+Stage I+Stage III used; non-proportional hazards: stage Stage IA (p 0.021))




COAD






Cox model · 253 patients, 58 events · HIT index q 0.11

HIT index + host expression + age + sex + stage (stage Stage IA, Stage IIB, Stage IIC, Stage III, Stage IIIA, Stage IVB merged into Stage IIA (16 patients, 1 events); stage baseline I absent; Stage IIA+Stage IA+Stage IIB+Stage …



Cox models: Numeric covariates per SD of the fit cohort; categories against the level named.



Exon · Region measured by HIT index · Higher in tumour · Lower in tumour · KM split · Paired Δ · Unpaired Δ · Cox p < 0.05 · Cox p ≥ 0.05 · Cohort shown at left · Model term p < 0.05 · p ≥ 0.05 · * q < 0.05 (Benjamini–Hochberg) · † non-proportional hazards (p < 0.05) · ‡ fewer than 20 events: low power

q: Benjamini–Hochberg within GLIPR1 for OS, one family per kind of test over all its HIT-index events × cohorts (within patients 13, all samples 15, KM 14, Cox, forest 14, model rows 14 tests); none for families under 10.