

GLIPR1 expression · BRCA, STAD and COAD · OS

The forest shows GLIPR1 expression in every cohort with a test; the rows below show the cohorts drawn on its splicing pages




Host-gene expression · GLIPR1 itself: tumour vs normal, a KM split at its median, and Cox on expression + age + sex + stage

BRCA · GLIPR1 expression




Cox model · 1062 patients, 146 events

expression + age + sex + stage (stage Stage IB, Stage II, Stage III merged into Stage IIA (14 patients, 2 events); stage baseline I absent; Stage IIA+Stage IB+Stage II+Stage III used; non-proportional hazards: stage Stage IIIB (p …




STAD · GLIPR1 expression




Cox model · 335 patients, 134 events

expression + age + sex + stage (stage Stage I, Stage III merged into Stage IIIA (4 patients, 3 events); stage baseline I absent; Stage IIIA+Stage I+Stage III used; non-proportional hazards: stage Stage IA (p 0.019))




COAD · GLIPR1 expression



Cox model · 253 patients, 58 events

expression + age + sex + stage (stage Stage IA, Stage IIB, Stage IIC, Stage III, Stage IIIA, Stage IVB merged into Stage IIA (16 patients, 1 events); stage baseline I absent; Stage IIA+Stage IA+Stage IIB+Stage IIC+Stage III+Stage …



Higher in tumour · Lower in tumour · KM split · Paired Δ · Unpaired Δ · Cox p < 0.05 · Cox p ≥ 0.05
Cohort drawn below · Model term p < 0.05 · p ≥ 0.05 · † non-proportional hazards (p < 0.05) · ‡ fewer than 20 events: low power

Host-gene expression of GLIPR1 on the expression table's scale; not adjusted for multiple testing. A tumour–normal hit needs p < 0.05 and |Δ median| > 1. The HR is per SD of expression in each cohort, with age, sex, stage.