

HSPA12A HSPA12A chr10:116834935-116835022 · KIRC, BRCA and LUSC · OS

HSPA12A chr10:116834935-116835022: mutually exclusive exons chr10:116,834,935–116,835,022 (88 nt) and chr10:116,707,200–116,707,285 (86 nt) between exons chr10:116,705,151–116,705,278 and chr10:116,849,566–116,849,710; PSI = use of the first · minus strand




KIRC




Cox model · 517 patients, 171 events · PSI q 0.10

PSI + host expression + age + sex + stage (stage baseline I absent; Stage I used; non-proportional hazards: stage Stage III (p 0.001), stage Stage IV (p 2.0e-4))

	HR (95% CI)	p
PSI (per SD)	1.25 (1.06–1.46)	0.007
Host expr. (per SD)	1.23 (1.02–1.48)	0.027
age (per SD)	1.50 (1.26–1.79)	5.3e-6
sex (vs female)		
male	0.97 (0.70–1.35)	0.85
stage (vs Stage I)		
Stage II	1.31 (0.70–2.44)	0.40
Stage III	2.55 (1.69–3.85)	8.4e-6 †
Stage IV	7.22 (4.84–10.76)	3.4e-22 †




BRCA




Cox model · 1060 patients, 146 events · PSI q 0.13

PSI + host expression + age + sex + stage (stage Stage IB, Stage II, Stage III merged into Stage IIA (14 patients, 2 events); stage baseline I absent; Stage IIA+Stage IB+Stage II+Stage III used; non-proportional hazards: host …

	HR (95% CI)	p
PSI (per SD)	0.82 (0.69–0.97)	0.019
Host expr. (per SD)	0.80 (0.68–0.95)	0.009 †
age (per SD)	1.60 (1.35–1.91)	1.5e-7
sex (vs female)		
male	0.41 (0.06–3.04)	0.38
stage (vs Stage IIA+Stage IB+Stage II+Stage III)		
Stage I	0.80 (0.42–1.53)	0.50
Stage IA	0.40 (0.12–1.29)	0.12
Stage IIB	1.38 (0.85–2.24)	0.19
Stage IIIA	1.98 (1.18–3.31)	0.010
Stage IIIB	3.09 (1.39–6.85)	0.006 †
Stage IIIC	2.73 (1.30–5.74)	0.008
Stage IV	8.33 (4.49–15.47)	1.8e-11
Stage X	2.56 (1.05–6.27)	0.039




LUSC




Cox model · 482 patients, 205 events · PSI q 0.13

PSI + host expression + age + sex + stage (stage Stage I, Stage II, Stage III, Stage IV merged into Stage IB (15 patients, 12 events); stage baseline I absent; Stage IB+Stage I+Stage II+Stage III+Stage IV used; non-proportional hazards: …

	HR (95% CI)	p
PSI (per SD)	0.85 (0.73–0.99)	0.035 †
Host expr. (per SD)	1.02 (0.87–1.19)	0.79
age (per SD)	1.21 (1.05–1.40)	0.010
sex (vs female)		
male	1.17 (0.85–1.62)	0.34
stage (vs Stage IB+Stage I+Stage II+Stage III+Stage IV)		
Stage IA	0.75 (0.49–1.14)	0.18 †
Stage IIA	0.94 (0.57–1.53)	0.79
Stage IIB	0.96 (0.65–1.41)	0.82
Stage IIIA	1.12 (0.73–1.73)	0.59
Stage IIIB	1.75 (0.87–3.52)	0.12



Cox models: Numeric covariates per SD of the fit cohort; categories against the level named.



Exon · Region measured by PSI · Other exon · Junction of the PSI form · of the other form
Higher in tumour · Lower in tumour · KM split · Paired Δ · Unpaired Δ · Cox p < 0.05 · Cox p ≥ 0.05
Cohort shown at left · Model term p < 0.05 · p ≥ 0.05 · * q < 0.05 (Benjamini–Hochberg) · † non-proportional hazards (p < 0.05)
‡ fewer than 20 events: low power

q: Benjamini–Hochberg within HSPA12A for OS, one family per kind of test over all its events × cohorts (within patients 13, all samples 15, KM 14, Cox, forest 14, model rows 14 tests); none for families under 10.