

HSPA12A expression · KIRC, BRCA and LUSC · OS

The forest shows HSPA12A expression in every cohort with a test; the rows below show the cohorts drawn on its splicing pages



Host-gene expression · HSPA12A itself: tumour vs normal, a KM split at its median, and Cox on expression + age + sex + stage


KIRC · HSPA12A expression




Cox model · 519 patients, 171 events

expression + age + sex + stage (stage baseline I absent; Stage I used; non-proportional hazards: stage Stage III (p 0.001), stage Stage IV (p 1.7e-4))

	HR (95% CI)	p
Expr. (per SD)	1.12 (0.94–1.33)	0.20
age (per SD)	1.50 (1.26–1.79)	7.5e-6
sex (vs female)		
male	0.97 (0.70–1.35)	0.87
stage (vs Stage I)		
Stage II	1.36 (0.73–2.54)	0.33
Stage III	2.62 (1.74–3.95)	4.4e-6 †
Stage IV	7.32 (4.90–10.91)	1.8e-22 †




BRCA · HSPA12A expression




Cox model · 1062 patients, 146 events

expression + age + sex + stage (stage Stage IB, Stage II, Stage III merged into Stage IIA (14 patients, 2 events); stage baseline I absent; Stage IIA+Stage IB+Stage II+Stage III used; non-proportional hazards: expression (p 0.004) …

	HR (95% CI)	p
Expr. (per SD)	0.84 (0.72–0.99)	0.036 †
age (per SD)	1.58 (1.32–1.88)	4.3e-7
sex (vs female)		
male	0.41 (0.06–3.05)	0.38
stage (vs Stage IIA+Stage IB+Stage II+Stage III)		
Stage I	0.84 (0.44–1.59)	0.58
Stage IA	0.38 (0.12–1.23)	0.11
Stage IIB	1.34 (0.83–2.18)	0.23
Stage IIIA	1.93 (1.15–3.23)	0.012
Stage IIIB	2.91 (1.31–6.45)	0.009 †
Stage IIIC	2.68 (1.27–5.63)	0.009
Stage IV	8.04 (4.34–14.89)	3.4e-11
Stage X	2.42 (0.99–5.92)	0.052



LUSC · HSPA12A expression



Cox model · 489 patients, 209 events

expression + age + sex + stage (stage Stage I, Stage II, Stage III, Stage IV merged into Stage IB (16 patients, 13 events); stage baseline I absent; Stage IB+Stage I+Stage II+Stage III+Stage IV used; non-proportional hazards: stage …

	HR (95% CI)	p
Expr. (per SD)	1.09 (0.94–1.26)	0.24
age (per SD)	1.20 (1.04–1.39)	0.011
sex (vs female)		
male	1.18 (0.85–1.63)	0.31
stage (vs Stage IB+Stage I+Stage II+Stage III+Stage IV)		
Stage IA	0.72 (0.48–1.09)	0.12 †
Stage IIA	0.91 (0.56–1.49)	0.72
Stage IIB	0.97 (0.66–1.43)	0.87
Stage IIIA	1.20 (0.79–1.82)	0.39
Stage IIIB	1.67 (0.83–3.35)	0.15

Higher in tumour · Lower in tumour · KM split · Paired Δ · Unpaired Δ · Cox p < 0.05 · Cox p ≥ 0.05

Cohort drawn below · Model term p < 0.05 · p ≥ 0.05 · † non-proportional hazards (p < 0.05) · ‡ fewer than 20 events: low power

Host-gene expression of HSPA12A on the expression table's scale; not adjusted for multiple testing. A tumour–normal hit needs p < 0.05 and |Δ median| > 1. The HR is per SD of expression in each cohort, with age, sex, stage.