

MACROH2A1 MACROH2A1 chr5:135352946-135353045 · KIRC, STAD and LUSC · OS

MACROH2A1 chr5:135352946-135353045: mutually exclusive exons chr5:135,352,946–135,353,045 (100 nt) and chr5:135,350,823–135,350,913 (91 nt) between exons chr5:135,345,968–135,346,057 and chr5:135,360,497–135,360,607; PSI = use of the first · minus strand




KIRC




Cox model · 519 patients, 171 events · PSI q 0.058

PSI + host expression + age + sex + stage (stage baseline I absent; Stage I used; non-proportional hazards: stage Stage III (p 0.003), stage Stage IV (p 8.8e-4))




STAD




Cox model · 335 patients, 134 events · PSI q 0.058

PSI + host expression + age + sex + stage (stage Stage I, Stage III merged into Stage IIIA (4 patients, 3 events); stage baseline I absent; Stage IIIA+Stage I+Stage III used; non-proportional hazards: stage Stage IA (p 0.009))




LUSC




Cox model · 489 patients, 209 events · PSI q 0.070

PSI + host expression + age + sex + stage (stage Stage I, Stage II, Stage III, Stage IV merged into Stage IB (16 patients, 13 events); stage baseline I absent; Stage IB+Stage I+Stage II+Stage III+Stage IV used; non-proportional hazards: …



Cox models: Numeric covariates per SD of the fit cohort; categories against the level named.



Exon · Region measured by PSI · Other exon · Junction of the PSI form · of the other form
Higher in tumour · Lower in tumour · KM split · Paired Δ · Unpaired Δ · Cox p < 0.05 · Cox p ≥ 0.05
Cohort shown at left · Model term p < 0.05 · p ≥ 0.05 · * q < 0.05 (Benjamini–Hochberg) · † non-proportional hazards (p < 0.05)
‡ fewer than 20 events: low power

q: Benjamini–Hochberg within MACROH2A1 for OS, one family per kind of test over all its events × cohorts (within patients 13, all samples 15, KM 14, Cox, forest 14, model rows 14 tests); none for families under 10.