

MACROH2A1 expression · KIRC, STAD and LUSC · OS


The forest shows MACROH2A1 expression in every cohort with a test; the rows below show the cohorts drawn on its splicing pages



Host-gene expression · MACROH2A1 itself: tumour vs normal, a KM split at its median, and Cox on expression + age + sex + stage

KIRC · MACROH2A1 expression







Cox model · 519 patients, 171 events


expression + age + sex + stage (stage baseline I absent; Stage I used; non-proportional hazards: stage Stage III (p 0.003), stage Stage IV (p 4.7e-4))

	HR (95% CI)	p
Expr. (per SD)	1.06 (0.90–1.24)	0.50
age (per SD)	1.47 (1.24–1.76)	1.6e-5
sex (vs female)		
male	0.93 (0.67–1.27)	0.64
stage (vs Stage I)		
Stage II	1.32 (0.71–2.47)	0.38
Stage III	2.52 (1.67–3.81)	1.2e-5 †
Stage IV	6.74 (4.55–10.00)	2.0e-21 †



STAD · MACROH2A1 expression







Cox model · 335 patients, 134 events


expression + age + sex + stage (stage Stage I, Stage III merged into Stage IIIA (4 patients, 3 events); stage baseline I absent; Stage IIIA+Stage I+Stage III used; non-proportional hazards: stage Stage IA (p 0.012))

	HR (95% CI)	p
Expr. (per SD)	0.95 (0.79–1.15)	0.63
age (per SD)	1.39 (1.15–1.69)	6.8e-4
sex (vs female)		
male	1.29 (0.89–1.87)	0.18
stage (vs Stage IIIA+Stage I+Stage III)		
Stage IA	0.24 (0.06–0.99)	0.049 †
Stage IB	0.54 (0.25–1.14)	0.11
Stage II	0.73 (0.35–1.50)	0.39
Stage IIA	0.62 (0.29–1.32)	0.22
Stage IIB	0.77 (0.41–1.45)	0.42
Stage IIIB	0.98 (0.56–1.71)	0.94
Stage IIIC	1.34 (0.72–2.50)	0.35
Stage IV	2.06 (1.16–3.67)	0.014



LUSC · MACROH2A1 expression






Cox model · 489 patients, 209 events

expression + age + sex + stage (stage Stage I, Stage II, Stage III, Stage IV merged into Stage IB (16 patients, 13 events); stage baseline I absent; Stage IB+Stage I+Stage II+Stage III+Stage IV used; non-proportional hazards: stage …

	HR (95% CI)	p
Expr. (per SD)	1.02 (0.89–1.18)	0.75
age (per SD)	1.21 (1.04–1.39)	0.011
sex (vs female)		
male	1.20 (0.86–1.66)	0.28
stage (vs Stage IB+Stage I+Stage II+Stage III+Stage IV)		
Stage IA	0.72 (0.47–1.09)	0.12 †
Stage IIA	0.91 (0.56–1.49)	0.71
Stage IIB	0.99 (0.67–1.46)	0.95
Stage IIIA	1.21 (0.79–1.83)	0.38
Stage IIIB	1.69 (0.84–3.39)	0.14



Higher in tumour · Lower in tumour · KM split · Paired Δ · Unpaired Δ · Cox p < 0.05 · Cox p ≥ 0.05
Cohort drawn below · Model term p < 0.05 · p ≥ 0.05 · † non-proportional hazards (p < 0.05) · ‡ fewer than 20 events: low power

Host-gene expression of MACROH2A1 on the expression table's scale; not adjusted for multiple testing. A tumour–normal hit needs p < 0.05 and |Δ median| > 1. The HR is per SD of expression in each cohort, with age, sex, stage.