

RGS1 RGS1 chr1:192575930-192576284 · HNSC, LIHC and LUAD · OS

RGS1 chr1:192575930-192576284: retained intron chr1:192,575,930–192,576,284 (355 nt) between exons chr1:192,575,773–192,575,929 and chr1:192,576,285–192,576,365; PSI = its retention · plus strand




HNSC




Cox model · 497 patients, 215 events · PSI q 0.038 *

PSI + host expression + age + sex + stage (stage Stage IVC merged into Stage IVA (2 patients, 2 events); stage baseline I absent; Stage IVA+Stage IVC used; narrow PSI range (SD 0.021 < 0.05))




LIHC




Cox model · 325 patients, 108 events · PSI q 0.038 *

PSI + host expression + age + sex + stage (stage Stage III, Stage IIIB, Stage IIIC, Stage IV, Stage IVA, Stage IVB merged into Stage I (22 patients, 12 events); stage baseline I absent; Stage I+Stage III+Stage IIIB+Stage IIIC+Stage …




LUAD




Cox model · 483 patients, 174 events · PSI q 0.038 *

PSI + host expression + age + sex + stage (stage Stage I, Stage II merged into Stage IB (6 patients, 2 events); stage baseline I absent; Stage IB+Stage I+Stage II used; narrow PSI range (SD 0.036 < 0.05); non-proportional hazards: …



Cox models: Numeric covariates per SD of the fit cohort; categories against the level named.



Exon · Region measured by PSI · Junction of the PSI form · of the other form
Higher in tumour · Lower in tumour · KM split · Paired Δ · Unpaired Δ · Cox p < 0.05 · Cox p ≥ 0.05
Cohort shown at left · Model term p < 0.05 · p ≥ 0.05 · * q < 0.05 (Benjamini–Hochberg) · † non-proportional hazards (p < 0.05)
‡ fewer than 20 events: low power

q: Benjamini–Hochberg within RGS1 for OS, one family per kind of test over all its events × cohorts (within patients 13, all samples 15, KM 14, Cox, forest 14, model rows 14 tests); none for families under 10.