
BRCA · ZNF276 HIT
Lobular
antisense FANCA ρ −0.66; host ρ +0.16 → FANCA readout
THCA · YY1 HIT
Follicular
antisense chr14:100.28 Mb, not measured; host ρ −0.53
UCEC · ERCC2 HIT
Serous
antisense KLC3 ρ −0.88; host ρ +0.21 → KLC3 readout
SARC · SLMAP HIT
LMS
no overlap; host ρ −0.76
TGCT · PRODH ALE
Non-seminoma
no overlap; host ρ −0.38
MESO · NDE1 MXE
Biphasic
no overlap; host ρ +0.34
STAD · GCSAM A3SS
Diffuse
no overlap; host ρ +0.01
LGG · IVD ALE
Astrocytoma
no overlap; host ρ −0.20
THYM · SAA4 ALE
A/AB
in readthrough SAA2-SAA4; host ρ +0.40
ESCA · NHERF4 RI
Adenocarcinoma
no overlap; host ρ −0.72
CESC · MNX1 A5SS
Adenocarcinoma
no overlap; host ρ −0.60
PCPG · RINL HIT
Paraganglioma
antisense chr19:38.87 Mb, not measured; host ρ −0.22
LGG_IDH · KHNYN HIT
IDHwt
antisense SDR39U1 ρ −0.28; host ρ +0.80
UCEC_TCGA · ERCC2 HIT
CN-high
antisense KLC3 ρ −0.89; host ρ +0.21 → KLC3 readout
STAD_TCGA · CCNDBP1 A3SS
MSI
no overlap; host ρ +0.15
COAD_TCGA · CCNDBP1 A3SS
MSI
no overlap; host ρ +0.06
PRAD_ETS · HLA-DMB SE
ERG
no overlap; host ρ −0.81
BRCA_ER · SEC16A MXE
ER-negative
no overlap; host ρ +0.52
HNSC_HPV · VRK2 HIT
HPV-positive
antisense FANCL ρ −0.60; host ρ +0.02 → FANCL readout
−1.00
−0.75
−0.50
−0.25
0.00
0.25
0.50
0.75
1.00
median over a class's patients: PSI or usage (0 to 1); HIT index (−1 to 1)