


Question 5

Part A - Visualization

By inspection of the following plot, the data is linearly separable...




Part B

Using CVX to solve the primal problem for SVM, I found:

$$w = [-0.5, 1.5]$$

$$b = -2.0$$

The following shows a plot of the hyperplane defined by w and b overlaid with the data...



Part C

Suppose there are N data points that have feature vectors of dimension M .

As evidenced by the attached scratchwork that investigates the case of $N = 3$, the latter part of $W(a)$ in the dual problem can be decomposed as $(\frac{1}{2})a^T P a$ where for the matrix P we have

$$P(i,j) = Y_i Y_j \langle X_i, X_j \rangle$$

Making this substitution into MATLAB then applying CVX to solve the dual problem of SVM, I found the following non-zero a 's and their corresponding support vectors:

$$a_{28} = 0.853301$$

$$\text{Support vector: } [7.000000, 3.000000]$$

$$a_{29} = 0.396669$$

$$\text{Support vector: } [10.000000, 4.000000]$$

$$a_{30} = 0.201101$$

$$\text{Support vector: } [0.000000, 2.000000]$$

$$a_{31} = 1.048869$$


$$\text{Support vector: } [9.000000, 5.000000]$$

I used these values to calculate w and b , which are reported as follows:

$$w = [-0.5, 1.5]$$

$$b = -2.0$$

There are 4 support vectors, circled on the following graph...



Code Attachment

```
% Import data
data = readtable('/Users/zackberger/Desktop/ML/HW/HW_4/Q5/Data.csv');
data = data{:,~3};

% Separate data into column vectors
X = data(:,[1,2]);
Y = data(:,3);

x1 = data(:,1);
x2 = data(:,2);

[num_data, data_dim] = size(X);

% Optimize primal SVM problem with CVX
cvx_begin

    variable w_primal(2);
    variable b_primal;

    minimize( norm(w_primal) );

    subject to
        Y.*(X*w_primal + b_primal) >= 1;

cvx_end

w_primal;
b_primal;

% Calculate the matrix P, where  $P(i,j) = Y_i Y_j \langle X_i, X_j \rangle$ 
P = zeros(num_data,num_data);
for i = 1:num_data
    for j = 1:num_data
        P(i,j) = X(i,:)*X(j,:)'*Y(i)*Y(j);
    end
end

one_vector = ones(num_data, 1);

% Optimize dual SVM problem with CVX
cvx_begin

    variable alpha_dual(num_data);

    minimize( (1/2)*alpha_dual'*P*alpha_dual - (one_vector'*alpha_dual) );
```

```

subject to
    alpha_dual >= 0;
    Y.*alpha_dual == 0;
cvx_end

% Loop through alpha_dual and for non-zero alpha, print corresponding support vector
num_support_vectors = 0;
for i = 1:num_data
    if alpha_dual(i) > .0001
        fprintf('alpha %i = %f\n', i, alpha_dual(i));
        fprintf('Support vector: [%f, %f]\n\n', X(i,1), X(i,2));
        num_support_vectors = num_support_vectors + 1;
    end
end

% Use alpha_dual to compute w and b
w_dual = [0, 0];
for i = 1:num_data
    if alpha_dual(i) > .0001
        w_dual = w_dual + alpha_dual(i)*X(i,:)*Y(i);
    end
end

b_dual = 0
for i = 1:num_data
    if alpha_dual(i) > .0001
        b_dual = b_dual + (1/num_support_vectors) * (Y(i) - w_dual*X(i,:).');
    end
end

w_dual
b_dual

% Graph results!
hold on

% Scatter plot feature vectors
for i = 1: numel(Y)
    if Y(i) == 1
        scatter(x1(i), x2(i), 'filled', 'b')
    else
        scatter(x1(i), x2(i), 'filled', 'r')
    end
end

% Plot primal decision boundary (w1*x1 + w2*x2 + b = 0)
% x1_axis = 0:1/100:10;
% x2_axis = (-b_primal - w_primal(1)*x1_axis) / w_primal(2);

```

```

% plot(x1_axis, x2_axis, 'g');

% Plot dual decision boundary ( $w_1x_1 + w_2x_2 + b = 0$ )
x1_axis = 0:1/100:10;
x2_axis = (-b_dual - w_dual(1)*x1_axis) / w_dual(2);
plot(x1_axis, x2_axis, 'g');

% Create graph details
title("Data");
xlabel("x1 feature");
ylabel("x2 feature");

L(1) = plot(nan, nan, 'b:');
L(2) = plot(nan, nan, 'r:');
L(3) = plot(nan, nan, 'g:');
legend(L, {'class 1', 'class -1', 'decision boundary'})

hold off

```