

COMP/EECE 7/8740 Neural Networks


Topics:

- Computational Graph
- How to compute the gradients during back-prop.?
- Notation + example

Md Zahangir Alom
Department of Computer Science
University of Memphis, TN

How do we compute gradients?

- Analytic or “Manual” Differentiation
- Symbolic Differentiation
- Numerical Differentiation
- Automatic Differentiation (AD)
 - Forward mode AD
 - Reverse mode AD
 - aka “backpropagation”



current W:

[0.34,
-1.11,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25347

gradient dW:

[?,
?,
?,
?,
?,
?,
?,
?,
?,...]

current W:

[0.34,
-1.11,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25347

W + h (first dim):

[0.34 + **0.0001**,
-1.11,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25322

gradient dW:

[?,
?,
?,
?,
?,
?,
?,
?,
?,...]

current W:

[0.34,
-1.11,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25347

W + h (first dim):

[0.34 + **0.0001**,
-1.11,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25322

gradient dW:

[-2.5,
?,
?,

$$(1.25322 - 1.25347)/0.0001 = -2.5$$

$$\frac{df(x)}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

?,
?,...]

current W:

[0.34,
-1.11,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25347

W + h (second dim):

[0.34,
-1.11 + **0.0001**,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25353

gradient dW:

[-2.5,
?,
?,
?,
?,
?,
?,
?,
?,...]

current W:

[0.34,
-1.11,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25347

W + h (second dim):

[0.34,
-1.11 + **0.0001**,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25353

gradient dW:

[-2.5,
0.6,
?,
?,

$$(1.25353 - 1.25347)/0.0001 = 0.6$$

$$\frac{df(x)}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

?,...]

current W:

[0.34,
-1.11,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25347

W + h (third dim):

[0.34,
-1.11,
0.78 + **0.0001**,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25347

gradient dW:

[-2.5,
0.6,
?,
?,
?,
?,
?,
?,
?,...]

current W:

[0.34,
-1.11,
0.78,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25347

W + h (third dim):

[0.34,
-1.11,
0.78 + **0.0001**,
0.12,
0.55,
2.81,
-3.1,
-1.5,
0.33,...]

loss 1.25347

gradient dW:

[-2.5,
0.6,
0,
?,
?,
?,...]


$$(1.25347 - 1.25347)/0.0001 = 0$$

$$\frac{df(x)}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

?,...]

Numerical vs Analytic Gradients

$$\frac{df(x)}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

- **Numerical gradient:** slow :(), approximate :(), easy to write :)
- **Analytic gradient:** fast :), exact :), error-prone :(
- **In practice:** Derive analytic gradient, check your implementation with numerical gradient.

This is called a **gradient check**.

Neural Turing Machine

inputs

loss







Figure reproduced with permission from a [Twitter post](#) by Andrej Karpathy.

Computational Graphs

- **A computational graph** is a way to represent a math function in the language of graph theory. Recall the premise of **graph** theory:
 - Nodes are connected by edges, and
 - Everything in the **graph** is either a node or an edge.
- In computational graph
 - Nodes are either input values or functions for combining values.
 - Edges receive their weights as the data flows through the graph.
 - Outbound edges from an input node are weighted with that input value
 - outbound nodes from a function node are weighted by combining the weights of the inbound edges using the specified function.


Directed Acyclic Graphs (DAGs)

- Exactly what the name suggests
 - Directed edges
 - No (directed) cycles
 - Underlying undirected cycles okay




Example

$$f(x_1, x_2) = x_1 x_2 + \sin(x_1)$$




Logistic Regression as a Cascade

Given a library of simple functions




Compose into a
→
complicate function

$$-\log \left(\frac{1}{1 + e^{-\mathbf{w}^\top \mathbf{x}}} \right)$$




Computational Graph and NNs




Neural Network Training

- Step 1: Compute Loss on mini-batch [F-Pass]




Neural Network Training

- Step 1: Compute Loss on mini-batch [F-Pass]




Neural Network Training

- Step 1: Compute Loss on mini-batch [F-Pass]




Neural Network Training

- Step 1: Compute Loss on mini-batch [F-Pass]
- Step 2: Compute gradients wrt parameters [B-Pass]




Neural Network Training

- Step 1: Compute Loss on mini-batch [F-Pass]
- Step 2: Compute gradients wrt parameters [B-Pass]




Neural Network Training

- Step 1: Compute Loss on mini-batch [F-Pass]
- Step 2: Compute gradients wrt parameters [B-Pass]



Neural Network Training

- Step 1: Compute Loss on mini-batch [F-Pass]
- Step 2: Compute gradients wrt parameters [B-Pass]
- Step 3: Use gradient to update parameters




$$\theta \leftarrow \theta - \eta \frac{dL}{d\theta}$$

Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$



Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$


e.g. $x = -2$, $y = 5$, $z = -4$



Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$




Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$

Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$



Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial f}$$

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial f}$$

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial z}$$

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial z}$$

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial q}$$

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial q}$$

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial y}$$

Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

Upstream
gradient

Local
gradient

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial y}$$

Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

Upstream
gradient

Local
gradient

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial x}$$

Chain rule:

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$

Upstream
gradient

Local
gradient

Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial x}$$


Chain rule:

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$

Upstream
gradient


Local
gradient

Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$




Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$




Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$f(x) = e^x$	\rightarrow	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	\rightarrow	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	\rightarrow	$\frac{df}{dx} = a$		$f_c(x) = c + x$	\rightarrow	$\frac{df}{dx} = 1$

Example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$




$$\begin{aligned} f(x) &= e^x & \rightarrow & \quad \frac{df}{dx} = e^x \\ f_a(x) &= ax & \rightarrow & \quad \frac{df}{dx} = a \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{1}{x} & \rightarrow & \quad \frac{df}{dx} = -1/x^2 \\ f_c(x) &= c + x & \rightarrow & \quad \frac{df}{dx} = 1 \end{aligned}$$

Example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$




$$\begin{aligned} f(x) &= e^x & \rightarrow & \frac{df}{dx} = e^x \\ f_a(x) &= ax & \rightarrow & \frac{df}{dx} = a \end{aligned}$$

$$\begin{aligned} f(x) &= \frac{1}{x} & \rightarrow & \frac{df}{dx} = -1/x^2 \\ f_c(x) &= c + x & \rightarrow & \frac{df}{dx} = 1 \end{aligned}$$

Example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	\rightarrow	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	\rightarrow	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	\rightarrow	$\frac{df}{dx} = a$		$f_c(x) = c + x$	\rightarrow	$\frac{df}{dx} = 1$

Example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	\rightarrow	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	\rightarrow	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	\rightarrow	$\frac{df}{dx} = a$		$f_c(x) = c + x$	\rightarrow	$\frac{df}{dx} = 1$

Example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$

Example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



[upstream gradient] x [local gradient]

$$[0.2] \times [1] = 0.2$$

$$[0.2] \times [1] = 0.2 \text{ (both inputs!)}$$

$$f(x) = e^x$$

\rightarrow

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

\rightarrow

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

\rightarrow

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

\rightarrow

$$\frac{df}{dx} = 1$$

Example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	\rightarrow	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	\rightarrow	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	\rightarrow	$\frac{df}{dx} = a$		$f_c(x) = c + x$	\rightarrow	$\frac{df}{dx} = 1$

Example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



[upstream gradient] x [local gradient]

$$x_0: [0.2] \times [2] = 0.4$$

$$w_0: [0.2] \times [-1] = -0.2$$

$f(x) = e^x$	\rightarrow	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	\rightarrow	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	\rightarrow	$\frac{df}{dx} = a$		$f_c(x) = c + x$	\rightarrow	$\frac{df}{dx} = 1$

Example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2)}}$$

Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!

$$\sigma(x) = \frac{1}{1 + e^{-x}}$$

sigmoid function

$$\frac{d\sigma(x)}{dx} = \frac{e^{-x}}{(1 + e^{-x})^2} = \left(\frac{1 + e^{-x} - 1}{1 + e^{-x}} \right) \left(\frac{1}{1 + e^{-x}} \right) = (1 - \sigma(x)) \sigma(x)$$



Example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2)}}$$

Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!

$$\sigma(x) = \frac{1}{1 + e^{-x}}$$

sigmoid function

$$\frac{d\sigma(x)}{dx} = \frac{e^{-x}}{(1 + e^{-x})^2} = \left(\frac{1 + e^{-x} - 1}{1 + e^{-x}} \right) \left(\frac{1}{1 + e^{-x}} \right) = (1 - \sigma(x)) \sigma(x)$$




[upstream gradient] x [local gradient]
 $[1.00] \times [(1 - 0.73) (0.73)] = 0.2$


Patterns in backward flow

$f(\cdot, \cdot)$

$$= 2(xy + \max\{z, w\})$$




Patterns in backward flow



Patterns in backward flow

add gate gradient distributor



$$w_3 = w_1 + w_2$$

$$\frac{\partial w_3}{\partial w_1} = 1$$

$$\boxed{\frac{\partial f}{\partial w_1}} = \boxed{\frac{\partial f}{\partial w_3}} \cdot \boxed{\frac{\partial w_3}{\partial w_1}}$$

Patterns in backward flow


$$\underline{w}_2 = \max\{\underline{z}, \underline{w}\}$$

$$\max(0, x) \text{ derivative } \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x \leq 0 \end{cases}$$

add gate: gradient distributor

Q: What is a max gate?


$\frac{\partial f}{\partial z}$
 $\frac{\partial f}{\partial w}$



Patterns in backward flow

add gate: gradient distributor

max gate: gradient router




Patterns in backward flow

add gate: gradient distributor

max gate: gradient router

Q: What is a mul gate?




$$\frac{\partial f}{\partial x} = \left[\frac{\partial f}{\partial w_1} \right] \left[\frac{\partial w_1}{\partial x} \right]$$

Patterns in backward flow

add gate: gradient distributor

max gate: gradient router


mul gate: gradient switcher



A vectorized example: $f(x, W) = \|W \cdot x\|^2 = \sum_{i=1}^n (W \cdot x)_i^2$

$$\begin{bmatrix} 0.1 & 0.5 \\ -0.3 & 0.8 \end{bmatrix} W$$


$$\begin{bmatrix} 0.2 \\ 0.4 \end{bmatrix} x$$



$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = \|q\|^2 = q_1^2 + \cdots + q_n^2$$


A vectorized example: $f(x, W) = \|W \cdot x\|^2 = \sum_{i=1}^n (W \cdot x)_i^2$



$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \dots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \dots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = \|q\|^2 = q_1^2 + \dots + q_n^2$$

A vectorized example: $f(x, W) = \|W \cdot x\|^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = \|q\|^2 = q_1^2 + \cdots + q_n^2$$

$$\frac{\partial f}{\partial q_i} = 2q_i$$

$$\nabla_q f = 2q$$

A vectorized example: $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$\nabla_W f = 2q \cdot x^T$$

$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\begin{aligned} \frac{\partial q_k}{\partial W_{i,j}} &= \mathbf{1}_{k=i} x_j \\ \frac{\partial f}{\partial W_{i,j}} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial W_{i,j}} \\ &= \sum_k (2q_k) (\mathbf{1}_{k=i} x_j) \\ &= 2q_i x_j \end{aligned}$$

A vectorized example: $f(x, W) = \|W \cdot x\|^2 = \sum_{i=1}^n (W \cdot x)_i^2$



Summary

- We have learned computational graphs and gradient computation methods
- How to apply computational graphs to calculate the gradient during back-propagation
- Notations & examples
- What is next ?
 - Linear classifiers
 - Loss functions : Hinge loss
 - Softmax Classifier (Multinomial Logistic Regression)