Inheritance

Mark scheme and worked solutions for 60 questions: 30 multiple choice, 30 structured.

1 ? Part 1

Core · Chromosomes, genes and proteins · [1]

Answer: B

Explanation

Chromosomes are made of DNA. Starch and cellulose are carbohydrates, and fat is a lipid; none is the chromosome's genetic material.

2 ? Part 1

Core · Chromosomes, genes and proteins · [1]

Answer: D

Explanation

A gene is a length of DNA coding for a protein. It is not a cell, membrane or the protein it encodes.

3 ? Part 1

Core · Chromosomes, genes and proteins · [1]

Answer: A

Explanation

Alleles are alternative forms of one gene. Ribosomes make proteins, gametes are reproductive cells, and enzymes are proteins with catalytic roles.

4 ? Part 1

Core · Chromosomes, genes and proteins · [1]

Answer: C

Explanation

Human males normally have XY and females XX in body cells. A single X is a gamete complement, not the normal male body-cell pair.

Topical Hub
topical-hub.pages.dev

5 ? Part 1

Core | Chromosomes, genes and proteins | [1]

Answer: D

Explanation

One X from each gamete gives XX. The sperm in this example contributes no Y and each normal gamete contributes one sex chromosome.

6 ? Part 1

Supplement | Chromosomes, genes and proteins | [1]

Answer: B

Explanation

The order of DNA bases determines the order of amino acids. Cell position and starch do not specify that sequence.

7 ? Part 1

Supplement | Chromosomes, genes and proteins | [1]

Answer: C

Explanation

Different amino-acid sequences can produce different folding and shapes. Identical function or active sites is not guaranteed.

8 ? Part 1

Supplement | Chromosomes, genes and proteins | [1]

Answer: A

Explanation

DNA controls production of proteins including carriers. A changed carrier may alter transport without removing DNA or changing every protein.

9 ? Part 1

Supplement | Chromosomes, genes and proteins | [1]

Answer: B

Explanation

mRNA copies genetic information and moves to the cytoplasm. Chromosomes normally remain in the nucleus during this process.

Topical Hub
topical-hub.pages.dev

10 ? Part 1 Supplement Chromosomes, genes and proteins [1]

Answer: C

Explanation

Ribosomes assemble amino acids in the order specified by mRNA. The other locations are not the site of protein assembly.

11 ? Part 1 Supplement Chromosomes, genes and proteins [1]

Answer: D

Explanation

Most body cells have the same genes but express different subsets. Specialisation does not normally require loss of all DNA or entirely different genomes.

12 ? Part 1 Supplement Chromosomes, genes and proteins [1]

Answer: A

Explanation

Haploid means one set of chromosomes. Diploid means two sets; neither term describes a protein or a pair of cells.

13 ? Part 1 Supplement Chromosomes, genes and proteins [1]

Answer: C

Explanation

Diploid nuclei contain two chromosome sets. This refers to chromosome sets, not a fixed total of one chromosome or amino acids.

14 ? Part 1 Supplement Chromosomes, genes and proteins [1]

Answer: A

Explanation

There are 23 pairs, giving 46 chromosomes. Twenty-three is the normal haploid number; 44 counts only the autosomes.

Topical Hub
topical-hub.pages.dev

15 ? Part 1 Supplement Chromosomes, genes and proteins [1]

Answer: B

Explanation

The mRNA base sequence specifies the amino-acid sequence. Oxygen availability can affect metabolism but does not encode the sequence.

16 ? Part 1 Core Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- Chromosomes are made of DNA. (1)
- A gene is a length of DNA that codes for a protein. (1)

(b) **[2]**

- An allele is an alternative form of a gene. (1)
- Different alleles have differences in their DNA sequence. (1)

Worked solution

(a) Chromosomes are made of DNA. A gene is a length of DNA that codes for a protein.

(b) An allele is an alternative form of a gene. Different alleles have differences in their DNA sequence.

17 ? Part 1 Core Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- The mother has XX. (1)
- The father has XY. (1)

(b) **[2]**

- Eggs each contain an X chromosome. (1)
- Sperm contain either X or Y, giving XX or XY after fertilisation. (1)

Worked solution

(a) The mother has XX. The father has XY.

(b) Eggs each contain an X chromosome. Sperm contain either X or Y, giving XX or XY after fertilisation.

Topical Hub
topical-hub.pages.dev

18 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- The DNA base sequence differs between the alleles. (1)
- The base sequence determines the order of amino acids in the protein. (1)

(b) **[2]**

- A changed amino-acid sequence can alter the protein's shape. (1)
- A changed active-site shape may reduce binding of the usual substrate. (1)

Worked solution

(a) The DNA base sequence differs between the alleles. The base sequence determines the order of amino acids in the protein.

(b) A changed amino-acid sequence can alter the protein's shape. A changed active-site shape may reduce binding of the usual substrate.

19 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- The gene controls production of a protein, the enzyme. (1)
- The enzyme catalyses a reaction, so its production affects that reaction. (1)

(b) **[2]**

- Membrane carrier proteins. (1)
- Receptors for neurotransmitters. (1)

Worked solution

(a) The gene controls production of a protein, the enzyme. The enzyme catalyses a reaction, so its production affects that reaction.

(b) Membrane carrier proteins. Receptors for neurotransmitters.

Topical Hub
topical-hub.pages.dev

20 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- An mRNA copy of the gene is made in the nucleus. (1)
- The mRNA moves from the nucleus to the cytoplasm. (1)

(b) **[2]**

- The mRNA passes through a ribosome. (1)
- The ribosome assembles amino acids in the sequence specified by the mRNA. (1)

Worked solution

(a) An mRNA copy of the gene is made in the nucleus. The mRNA moves from the nucleus to the cytoplasm.

(b) The mRNA passes through a ribosome. The ribosome assembles amino acids in the sequence specified by the mRNA.

21 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- Most body cells in one individual contain the same genes. (1)
- A gene can be present without being expressed. (1)

(b) **[2]**

- Different cells express different genes. (1)
- They produce the particular proteins needed for their specific functions. (1)

Worked solution

(a) Most body cells in one individual contain the same genes. A gene can be present without being expressed.

(b) Different cells express different genes. They produce the particular proteins needed for their specific functions.

Topical Hub
topical-hub.pages.dev

22 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- Diploid means containing two sets of chromosomes. (1)
- The haploid number is 8. (1)

(b) **[2]**

- A haploid nucleus contains one set of chromosomes. (1)
- One set is half the diploid number, so 16 divided by 2 is 8. (1)

Worked solution

(a) Diploid means containing two sets of chromosomes. The haploid number is 8.

(b) A haploid nucleus contains one set of chromosomes. One set is half the diploid number, so 16 divided by 2 is 8.

23 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- A body-cell nucleus contains 46 chromosomes. (1)
- A gamete nucleus contains 23 chromosomes. (1)

(b) **[2]**

- The body-cell nucleus is diploid with two sets. (1)
- The gamete nucleus is haploid with one set. (1)

Worked solution

(a) A body-cell nucleus contains 46 chromosomes. A gamete nucleus contains 23 chromosomes.

(b) The body-cell nucleus is diploid with two sets. The gamete nucleus is haploid with one set.

Topical Hub
topical-hub.pages.dev

24 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- The gene remains in the nucleus. (1)
- An mRNA copy moves to the cytoplasm instead. (1)

(b) **[2]**

- A ribosome assembles amino acids to form the protein. (1)
- The sequence is specified by the mRNA rather than DNA being converted directly into protein. (1)

Worked solution

(a) The gene remains in the nucleus. An mRNA copy moves to the cytoplasm instead.

(b) A ribosome assembles amino acids to form the protein. The sequence is specified by the mRNA rather than DNA being converted directly into protein.

25 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- DNA controls production of the receptor protein. (1)
- Whether the receptor is present affects the cell's ability to detect the signal. (1)

(b) **[2]**

- The receptor gene is expressed in one cell type. (1)
- It is not expressed, or is expressed much less, in the other cell type. (1)

Worked solution

(a) DNA controls production of the receptor protein. Whether the receptor is present affects the cell's ability to detect the signal.

(b) The receptor gene is expressed in one cell type. It is not expressed, or is expressed much less, in the other cell type.

Topical Hub
topical-hub.pages.dev

26 ? Part 1 Core Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- Some sperm contain X. (1)
- Other sperm contain Y. (1)

(b) **[2]**

- Fertilisation by an X-bearing or Y-bearing sperm is a new event. (1)
- Each outcome remains approximately equally likely regardless of earlier children. (1)

Worked solution

(a) Some sperm contain X. Other sperm contain Y.

(b) Fertilisation by an X-bearing or Y-bearing sperm is a new event. Each outcome remains approximately equally likely regardless of earlier children.

27 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- The sequence of bases in the gene specifies amino-acid order. (1)
- The two individuals may have different alleles of that gene. (1)

(b) **[2]**

- Different amino-acid sequences can produce different protein shapes. (1)
- A change in shape may affect binding or transport of the usual substance. (1)

Worked solution

(a) The sequence of bases in the gene specifies amino-acid order. The two individuals may have different alleles of that gene.

(b) Different amino-acid sequences can produce different protein shapes. A change in shape may affect binding or transport of the usual substance.

Topical Hub
topical-hub.pages.dev

28 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- The amino acids could be assembled in different sequences. (1)
- The correct order must be specified by genetic information. (1)

(b) **[2]**

- The mRNA base sequence supplies the sequence information. (1)
- The ribosome reads that information while assembling the amino acids. (1)

Worked solution

(a) The amino acids could be assembled in different sequences. The correct order must be specified by genetic information.

(b) The mRNA base sequence supplies the sequence information. The ribosome reads that information while assembling the amino acids.

29 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- P has 10 divided by 2, giving 5. (1)
- Q has 28 divided by 2, giving 14. (1)

(b) **[2]**

- Diploid describes two complete sets, regardless of the number in each set. (1)
- Both species have chromosome pairs, although the number of pairs differs. (1)

Worked solution

(a) P has 10 divided by 2, giving 5. Q has 28 divided by 2, giving 14.

(b) Diploid describes two complete sets, regardless of the number in each set. Both species have chromosome pairs, although the number of pairs differs.

Topical Hub
topical-hub.pages.dev

30 ? Part 1 Supplement Chromosomes, genes and proteins [4]

Mark scheme

(a) **[2]**

- A chromosome is made of DNA. (1)
- A gene is a length of that DNA coding for a protein. (1)

(b) **[2]**

- The changed gene can alter production of the enzyme protein. (1)
- Less enzyme may reduce the rate of the metabolic reaction it catalyses. (1)

Worked solution

(a) A chromosome is made of DNA. A gene is a length of that DNA coding for a protein.

(b) The changed gene can alter production of the enzyme protein. Less enzyme may reduce the rate of the metabolic reaction it catalyses.

1 ? Part 2 Supplement Mitosis [1]

Answer: B

Explanation

Mitosis gives genetically identical nuclei and maintains chromosome number. Halving is associated with meiosis and fusion with fertilisation.

2 ? Part 2 Supplement Mitosis [1]

Answer: D

Explanation

Exact replication provides copies that can be separated into daughter nuclei. Destruction, fertilisation and reduction division are different processes.

3 ? Part 2 Supplement Mitosis [1]

Answer: A

Explanation

Stem cells divide by mitosis and their daughter cells can specialise. They contain chromosomes and are not defined by pollen production.

Topical Hub
topical-hub.pages.dev

4 ? Part 2

Supplement | Meiosis | [1]

Answer: C

Explanation

Meiosis produces gametes with reduced chromosome number. Fertilisation joins gametes, while pollination and digestion have other roles.

5 ? Part 2

Supplement | Meiosis | [1]

Answer: D

Explanation

Meiosis halves 32 to 16 and produces genetically different cells. Neither chromosome doubling nor repeated halving to eight describes the process.

6 ? Part 2

Core | Monohybrid inheritance | [1]

Answer: B

Explanation

Phenotype means observable features. Rr is the genotype; R and r are alleles, and a gamete is a reproductive cell.

7 ? Part 2

Core | Monohybrid inheritance | [1]

Answer: C

Explanation

Heterozygous means two different alleles of one gene, here T and t. TT and tt are homozygous; the unrelated notation is not a genotype for this gene.

8 ? Part 2

Core | Monohybrid inheritance | [1]

Answer: A

Explanation

The recessive feature is expressed when no dominant allele is present, as in bb. BB and Bb contain B; one allele alone is not the stated diploid genotype.

Topical Hub
topical-hub.pages.dev

9 ? Part 2

Core | Monohybrid inheritance | [1]

Answer: B

Explanation

Pp produces P and p gametes; pp produces p gametes only. The offspring are equally likely Pp or pp, giving 1 : 1.

10 ? Part 2

Core | Monohybrid inheritance | [1]

Answer: C

Explanation

A 3 : 1 ratio under complete dominance includes one homozygous recessive combination among four equally likely combinations. Both other genotypes express the dominant feature.

11 ? Part 2

Supplement | Monohybrid inheritance | [1]

Answer: D

Explanation

A recessive partner allows recessive offspring to reveal a recessive allele in the unknown parent. A dominant partner can mask that distinction.

12 ? Part 2

Supplement | Monohybrid inheritance | [1]

Answer: A

Explanation

In codominance both alleles contribute to the heterozygote's phenotype. Neither allele is simply absent or completely masked.

13 ? Part 2

Supplement | Monohybrid inheritance | [1]

Answer: C

Explanation

Group O requires two I^o alleles. I^A and I^B are each dominant over I^o, and together they give group AB.

Topical Hub
topical-hub.pages.dev

14 ? Part 2 Supplement Monohybrid inheritance [1]

Answer: A

Explanation

A male's single X can express a recessive colour-blindness allele without another X carrying a dominant allele to mask it.

15 ? Part 2 Core Monohybrid inheritance [1]

Answer: B

Explanation

Two aa parents produce only a gametes, so all offspring are aa. Crosses containing heterozygotes can produce more than one genotype.

16 ? Part 2 Supplement Mitosis [4]

Mark scheme

(a) **[2]**

- Each daughter nucleus has 24 chromosomes. (1)
- Copies of each replicated chromosome separate so each nucleus receives a complete set. (1)

(b) **[2]**

- Growth by increasing cell number. (1)
- Asexual reproduction or replacement of worn-out cells. (1)

Worked solution

(a) Each daughter nucleus has 24 chromosomes. Copies of each replicated chromosome separate so each nucleus receives a complete set.

(b) Growth by increasing cell number. Asexual reproduction or replacement of worn-out cells.

Topical Hub
topical-hub.pages.dev

17 ? Part 2 Supplement Mitosis [4]

Mark scheme

(a) **[2]**

- Each daughter nucleus needs a complete set of genetic information. (1)
- Exact copying helps both daughter nuclei receive the same information as the parent. (1)

(b) **[2]**

- The copies separate. (1)
- Each daughter nucleus receives one copy of each chromosome. (1)

Worked solution

(a) Each daughter nucleus needs a complete set of genetic information. Exact copying helps both daughter nuclei receive the same information as the parent.

(b) The copies separate. Each daughter nucleus receives one copy of each chromosome.

18 ? Part 2 Supplement Mitosis [4]

Mark scheme

(a) **[2]**

- Stem cells are unspecialised cells. (1)
- They divide by mitosis to produce daughter cells. (1)

(b) **[2]**

- The daughter cells become specialised. (1)
- They develop the features needed for that tissue's particular function. (1)

Worked solution

(a) Stem cells are unspecialised cells. They divide by mitosis to produce daughter cells.

(b) The daughter cells become specialised. They develop the features needed for that tissue's particular function.

Topical Hub
topical-hub.pages.dev

19 ? Part 2

Supplement | Meiosis | [4]

Mark scheme

(a) **[2]**

- Each gamete nucleus has 6 chromosomes. (1)
- Gametes are genetically different from one another. (1)

(b) **[2]**

- It halves the number of chromosome sets. (1)
- It changes a diploid nucleus to haploid nuclei. (1)

Worked solution

(a) Each gamete nucleus has 6 chromosomes. Gametes are genetically different from one another.

(b) It halves the number of chromosome sets. It changes a diploid nucleus to haploid nuclei.

20 ? Part 2

Core | Monohybrid inheritance | [4]

Mark scheme

(a) **[2]**

- Hh is its genotype, describing the alleles present. (1)
- Hairy leaves are its phenotype, an observable feature. (1)

(b) **[2]**

- Inheritance is transmission of genetic information from generation to generation. (1)
- Alleles of genes are passed from parents to offspring. (1)

Worked solution

(a) Hh is its genotype, describing the alleles present. Hairy leaves are its phenotype, an observable feature.

(b) Inheritance is transmission of genetic information from generation to generation. Alleles of genes are passed from parents to offspring.

Topical Hub
topicalhub.pages.dev

21 ? Part 2

Core | Monohybrid inheritance | [4]

Mark scheme

(a) **[2]**

- BB and bb are homozygous because each has two identical alleles. (1)
- Bb is heterozygous because its alleles differ. (1)

(b) **[2]**

- Bb is black because the dominant B allele is expressed. (1)
- bb is white because no dominant B allele is present. (1)

Worked solution

(a) BB and bb are homozygous because each has two identical alleles. Bb is heterozygous because its alleles differ.

(b) Bb is black because the dominant B allele is expressed. bb is white because no dominant B allele is present.

22 ? Part 2

Core | Monohybrid inheritance | [4]

Mark scheme

(a) **[2]**

- Each parent produces S and s gametes. (1)
- The offspring combinations are SS, Ss, Ss and ss. (1)

(b) **[2]**

- The ratio is 3 smooth : 1 wrinkled. (1)
- Only ss offspring are wrinkled because the dominant S allele is absent. (1)

Worked solution

(a) Each parent produces S and s gametes. The offspring combinations are SS, Ss, Ss and ss.

(b) The ratio is 3 smooth : 1 wrinkled. Only ss offspring are wrinkled because the dominant S allele is absent.

Topical Hub
topical-hub.pages.dev

23 ? Part 2

Core | Monohybrid inheritance | [4]

Mark scheme

(a) **[2]**

- The Ll parent produces L and l gametes. (1)
- The ll parent produces l gametes only. (1)

(b) **[2]**

- Offspring are equally likely Ll and ll. (1)
- The ratio is 1 long-winged : 1 short-winged. (1)

Worked solution

(a) The Ll parent produces L and l gametes. The ll parent produces l gametes only.

(b) Offspring are equally likely Ll and ll. The ratio is 1 long-winged : 1 short-winged.

24 ? Part 2

Supplement | Monohybrid inheritance | [4]

Mark scheme

(a) **[2]**

- Cross the red-flowered plant with a white-flowered rr plant. (1)
- Record the flower colours of many offspring. (1)

(b) **[2]**

- White offspring show the unknown parent supplied r and is Rr. (1)
- A small sample could by chance contain only red offspring even if the parent is Rr. (1)

Worked solution

(a) Cross the red-flowered plant with a white-flowered rr plant. Record the flower colours of many offspring.

(b) White offspring show the unknown parent supplied r and is Rr. A small sample could by chance contain only red offspring even if the parent is Rr.

Topical Hub
topicalhub.pages.dev

25 ? Part 2

Supplement | Monohybrid inheritance | [4]

Mark scheme

(a) **[2]**

- Both parents produce C^R and C^W gametes. (1)
- Offspring genotypes are C^RC^R, C^RC^W and C^WC^W. (1)

(b) **[2]**

- The ratio is 1 red : 2 mixed red-and-white : 1 white. (1)
- Both alleles contribute to the heterozygote's phenotype because they are codominant. (1)

Worked solution

(a) Both parents produce C^R and C^W gametes. Offspring genotypes are C^RC^R, C^RC^W and C^WC^W.

(b) The ratio is 1 red : 2 mixed red-and-white : 1 white. Both alleles contribute to the heterozygote's phenotype because they are codominant.

26 ? Part 2

Supplement | Monohybrid inheritance | [4]

Mark scheme

(a) **[2]**

- The first produces I^A and I^o gametes. (1)
- The second produces I^B and I^o gametes. (1)

(b) **[2]**

- The combinations give AB, A, B and O. (1)
- These groups occur in an expected ratio of 1 : 1 : 1 : 1. (1)

Worked solution

(a) The first produces I^A and I^o gametes. The second produces I^B and I^o gametes.

(b) The combinations give AB, A, B and O. These groups occur in an expected ratio of 1 : 1 : 1 : 1.

Topical Hub
topical-hub.pages.dev

27 ? Part 2

Supplement | Monohybrid inheritance | [4]

Mark scheme

(a) **[2]**

- X^NX^N. (1)
- X^NX^n. (1)

(b) **[2]**

- Half the sons are expected to be X^nY. (1)
- A son receives Y from his father and has an equal chance of receiving either maternal X. (1)

Worked solution

(a) X^NX^N. X^NX^n.

(b) Half the sons are expected to be X^nY. A son receives Y from his father and has an equal chance of receiving either maternal X.

28 ? Part 2

Core | Monohybrid inheritance | [4]

Mark scheme

(a) **[2]**

- PP is consistent with the line. (1)
- Two PP parents pass on P only, so all offspring are PP and purple. (1)

(b) **[2]**

- Both parents can pass on p. (1)
- Some offspring will be pp and have white flowers. (1)

Worked solution

(a) PP is consistent with the line. Two PP parents pass on P only, so all offspring are PP and purple.

(b) Both parents can pass on p. Some offspring will be pp and have white flowers.

Topical Hub
topical-hub.pages.dev

29 ? Part 2 Supplement Monohybrid inheritance [4]

Mark scheme

(a) **[2]**

- The AB parent is I^AI^B and produces I^A or I^B gametes. (1)
- The O parent is I^oI^o and produces I^o gametes only. (1)

(b) **[2]**

- Half are expected to be A and half B. (1)
- Each child receives I^o from the O parent, so no child receives both I^A and I^B. (1)

Worked solution

(a) The AB parent is I^AI^B and produces I^A or I^B gametes. The O parent is I^oI^o and produces I^o gametes only.

(b) Half are expected to be A and half B. Each child receives I^o from the O parent, so no child receives both I^A and I^B.

30 ? Part 2 Core Monohybrid inheritance [4]

Mark scheme

(a) **[2]**

- A heterozygote cross predicts three quarters dominant, so 200 × 3 / 4 = 150. (1)
- One quarter recessive gives 200 × 1 / 4 = 50. (1)

(b) **[2]**

- Which gametes fuse is a chance event. (1)
- An expected probability does not guarantee an exact ratio in a finite sample. (1)

Worked solution

(a) A heterozygote cross predicts three quarters dominant, so 200 × 3 / 4 = 150. One quarter recessive gives 200 × 1 / 4 = 50.

(b) Which gametes fuse is a chance event. An expected probability does not guarantee an exact ratio in a finite sample.

Topical Hub
topical-hub.pages.dev