Reproduction

Mark scheme and worked solutions for 60 questions: 30 multiple choice, 30 structured.

1 · Part 1

Core · Asexual reproduction · [1]

Answer: B

Explanation

Asexual reproduction uses one parent and produces genetically identical offspring. Neither gamete fusion nor the mixing of two parents' genes occurs.

2 · Part 1

Core · Asexual reproduction · [1]

Answer: D

Explanation

Division from one parent without gamete fusion is asexual reproduction. Pollination transfers pollen; fertilisation and sexual reproduction involve gamete nuclei.

3 · Part 1

Extended · Asexual reproduction · [1]

Answer: A

Explanation

Cloning retains the parent's genetic combination. It does not create broad variation or guarantee disease resistance, and it does not require pollination.

4 · Part 1

Core · Sexual reproduction · [1]

Answer: C

Explanation

Sexual reproduction involves fusion of gamete nuclei. Growth, cell multiplication and production of offspring also occur in asexual reproduction.

Topical Hub
topical-hub.pages.dev

5 · Part 1 Core Sexual reproduction [1]

Answer: D

Explanation

Fertilisation is nuclear fusion. Landing on the stigma is pollination, while tube growth enables the nuclei to meet but is not itself fertilisation.

6 · Part 1 Extended Sexual reproduction [1]

Answer: C

Explanation

Two haploid sets of 12 combine, giving 12 + 12 = 24 chromosomes. Fertilisation adds sets rather than halving or doubling each set separately.

7 · Part 1 Core Sexual reproduction in plants [1]

Answer: C

Explanation

The anther produces pollen. The stigma receives it, the ovary contains ovules, and petals may attract pollinators.

8 · Part 1 Core Sexual reproduction in plants [1]

Answer: A

Explanation

A spiky (or sticky) surface helps pollen attach to an insect's body. Smooth, small and light grains are characteristic of wind-pollinated flowers and are less likely to be carried by insects.

9 · Part 1 Core Sexual reproduction in plants [1]

Answer: B

Explanation

Pollination transfers pollen from an anther to a stigma. Movement to an ovule occurs later, involving a pollen tube rather than transfer of whole grains.

Topical Hub
topical-hub.pages.dev

10 · Part 1 Core Sexual reproduction in plants [1]

Answer: C

Explanation

Exposed anthers release pollen into moving air and a feathery stigma traps airborne grains. Nectar, scent and showy petals chiefly attract animals.

11 · Part 1 Core Sexual reproduction in plants [1]

Answer: D

Explanation

Water is required for germination. Light is not a universal requirement, so darkness alone does not prevent all viable seeds from germinating.

12 · Part 1 Extended Sexual reproduction in plants [1]

Answer: A

Explanation

Both flowers belong to the same plant, so this is self-pollination. Cross-pollination requires a different plant of the same species.

13 · Part 1 Extended Sexual reproduction in plants [1]

Answer: C

Explanation

Cross-pollination transfers pollen to a flower on another plant of the same species. Neither within-plant transfer nor seed movement fits this definition.

14 · Part 1 Extended Sexual reproduction in plants [1]

Answer: A

Explanation

Cross-pollination requires transfer between plants, so reproduction can become vulnerable to failure of that transfer. It does not abolish fertilisation or embryo formation.

Topical Hub
topical-hub.pages.dev

15 · Part 1 Extended Sexual reproduction in plants [1]

Answer: B

Explanation

The tube grows from the stigma through the style towards an ovule, carrying the male nucleus. The other routes do not deliver it to the female nucleus.

16 · Part 1 Extended Asexual reproduction [4]

Mark scheme

(a) **[2]**

- The plants have the same genetic information for relevant inherited features. (1)
- They are therefore likely to produce similar crop characteristics under the same conditions. (1)

(b) **[2]**

- The plants have little genetic variation. (1)
- If the cloned genotype is susceptible, many plants may be affected together. (1)

Worked solution

(a) The plants have the same genetic information for relevant inherited features. They are therefore likely to produce similar crop characteristics under the same conditions.

(b) The plants have little genetic variation. If the cloned genotype is susceptible, many plants may be affected together.

Topical Hub
topical-hub.pages.dev

17 · Part 1 Extended Asexual reproduction [4]

Mark scheme

(a) **[2]**

- Only one parent is required, so no mate is needed. (1)
- Offspring can be produced without the delay of finding another reproductive individual. (1)

ALLOW no pollinator, gametes or fertilisation needed; many offspring produced in a short time

(b) **[2]**

- The offspring are genetically identical apart from mutation. (1)
- There may be no individuals with inherited features suited to the changed conditions. (1)

Worked solution

(a) Only one parent is required, so no mate is needed. Offspring can be produced without the delay of finding another reproductive individual.

(b) The offspring are genetically identical apart from mutation. There may be no individuals with inherited features suited to the changed conditions.

18 · Part 1 Core Sexual reproduction [4]

Mark scheme

(a) **[2]**

- Fertilisation is fusion of the nuclei of two gametes. (1)
- The resulting cell is a zygote. (1)

(b) **[2]**

- Each offspring receives genetic information from both gametes. (1)
- Offspring are genetically different from one another. (1)

Worked solution

(a) Fertilisation is fusion of the nuclei of two gametes. The resulting cell is a zygote.

(b) Each offspring receives genetic information from both gametes. Offspring are genetically different from one another.

Topical Hub
topical-hub.pages.dev

19 · Part 1 Extended Sexual reproduction [4]

Mark scheme

(a) **[2]**

- A gamete has 9 chromosomes. (1)
- A gamete is haploid and has one set, half the diploid body-cell number. (1)

(b) **[2]**

- The zygote has 18 chromosomes. (1)
- The two gamete nuclei contribute 9 chromosomes each during fusion. (1)

Worked solution

(a) A gamete has 9 chromosomes. A gamete is haploid and has one set, half the diploid body-cell number.

(b) The zygote has 18 chromosomes. The two gamete nuclei contribute 9 chromosomes each during fusion.

20 · Part 1 Extended Sexual reproduction [4]

Mark scheme

(a) **[2]**

- Sexual reproduction produces genetic variation among offspring. (1)
- Some offspring may possess a combination of alleles better suited to the new climate. (1)

(b) **[2]**

- The offspring may not retain all desirable parental characteristics. (1)
- Time and resources are needed for sexual reproduction and subsequent selection. (1)

Worked solution

(a) Sexual reproduction produces genetic variation among offspring. Some offspring may possess a combination of alleles better suited to the new climate.

(b) The offspring may not retain all desirable parental characteristics. Time and resources are needed for sexual reproduction and subsequent selection.

Topical Hub
topical-hub.pages.dev

21 · Part 1 Extended Sexual reproduction [4]

Mark scheme

(a) **[2]**

- Some individuals may have inherited features that help them avoid the predator. (1)
- These individuals are more likely to survive and reproduce. (1)

(b) **[2]**

- Individuals must obtain compatible gametes, normally by finding a mate. (1)
- Separation reduces the chance of mating and therefore successful fertilisation. (1)

Worked solution

(a) Some individuals may have inherited features that help them avoid the predator. These individuals are more likely to survive and reproduce.

(b) Individuals must obtain compatible gametes, normally by finding a mate. Separation reduces the chance of mating and therefore successful fertilisation.

22 · Part 1 Core Sexual reproduction in plants [4]

Mark scheme

(a) **[2]**

- The stigma receives pollen grains. (1)
- The ovary contains ovules. (1)

ALLOW stigma has a sticky surface that traps pollen; ovary protects the ovules / develops into the fruit

(b) **[2]**

- The damaged stigma may receive less pollen, reducing successful fertilisation. (1)
- The intact anthers can still produce pollen grains. (1)

Worked solution

(a) The stigma receives pollen grains. The ovary contains ovules.

(b) The damaged stigma may receive less pollen, reducing successful fertilisation. The intact anthers can still produce pollen grains.

Topical Hub
topical-hub.pages.dev

23 · Part 1 Core Sexual reproduction in plants [4]

Mark scheme

(a) **[2]**

- A is insect-pollinated because its nectar or colour attracts insects. (1)
- B is wind-pollinated because exposed hanging anthers release pollen into air. (1)

(b) **[2]**

- A is likely to have stickier or spikier pollen than B. (1)
- B is likely to produce smaller, lighter pollen grains than A. (1)

ALLOW B produces larger numbers of pollen grains; A has larger pollen grains; B pollen is smooth

Worked solution

(a) A is insect-pollinated because its nectar or colour attracts insects. B is wind-pollinated because exposed hanging anthers release pollen into air.

(b) A is likely to have stickier or spikier pollen than B. B is likely to produce smaller, lighter pollen grains than A.

24 · Part 1 Core Sexual reproduction in plants [4]

Mark scheme

(a) **[2]**

- Pollination is transfer of pollen grains from anther to stigma. (1)
- Fertilisation is fusion of a pollen nucleus with a nucleus in an ovule. (1)

(b) **[2]**

- The pollen may be incompatible with the receiving flower. (1)
- The pollen grain may fail to germinate or form a functioning pollen tube. (1)

ALLOW pollen from a different species; pollen not viable; stigma not receptive or not mature; pollen tube fails to reach the ovule

Worked solution

(a) Pollination is transfer of pollen grains from anther to stigma. Fertilisation is fusion of a pollen nucleus with a nucleus in an ovule.

(b) The pollen may be incompatible with the receiving flower. The pollen grain may fail to germinate or form a functioning pollen tube.

Topical Hub
topical-hub.pages.dev

25 · Part 1 Core Sexual reproduction in plants [4]

Mark scheme

(a) **[2]**

- The condition being tested is water availability. (1)
- More seeds are expected to germinate in A than in B. (1)

(b) **[2]**

- The temperature must be the same. (1)
- The availability of oxygen must be the same. (1)

ALLOW same number, species or age of seeds; same light conditions; same length of time

Worked solution

(a) The condition being tested is water availability. More seeds are expected to germinate in A than in B.

(b) The temperature must be the same. The availability of oxygen must be the same.

26 · Part 1 Core Sexual reproduction in plants [4]

Mark scheme

(a) **[2]**

- Seeds in air germinate while those without oxygen do not. (1)
- Oxygen is required for germination under these conditions. (1)

(b) **[2]**

- Use seeds of the same species and similar viability. (1)
- Provide the same water availability in both containers. (1)

ALLOW same number of seeds; same light conditions; same length of time; same volume of container or gas

Worked solution

(a) Seeds in air germinate while those without oxygen do not. Oxygen is required for germination under these conditions.

(b) Use seeds of the same species and similar viability. Provide the same water availability in both containers.

Topical Hub
topical-hub.pages.dev

27 · Part 1 Extended Sexual reproduction in plants [4]

Mark scheme

(a) **[2]**

- X to Y is self-pollination because both flowers are on one plant. (1)
- X to Z is cross-pollination because the flowers are on different plants of the same species. (1)

(b) **[2]**

- The pollen starts on an anther. (1)
- It is received by a stigma. (1)

Worked solution

(a) X to Y is self-pollination because both flowers are on one plant. X to Z is cross-pollination because the flowers are on different plants of the same species.

(b) The pollen starts on an anther. It is received by a stigma.

28 · Part 1 Extended Sexual reproduction in plants [4]

Mark scheme

(a) **[2]**

- Q is more likely to have a wider range of combinations. (1)
- Cross-pollination combines genetic information from different plants. (1)

(b) **[2]**

- P may still reproduce if self-pollination can occur without insects. (1)
- Q may produce fewer seeds because pollen transfer between plants is reduced. (1)

Worked solution

(a) Q is more likely to have a wider range of combinations. Cross-pollination combines genetic information from different plants.

(b) P may still reproduce if self-pollination can occur without insects. Q may produce fewer seeds because pollen transfer between plants is reduced.

Topical Hub
topical-hub.pages.dev

29 · Part 1 Extended Sexual reproduction in plants [4]

Mark scheme

(a) **[2]**

- A pollen tube grows through the style towards an ovule. (1)
- The male nucleus travels along the pollen tube into the ovule. (1)

ALLOW pollen tube grows from the stigma, through the style, into the ovary and enters the ovule

(b) **[2]**

- The male nucleus fuses with a nucleus in the ovule. (1)
- This forms a zygote. (1)

Worked solution

(a) A pollen tube grows through the style towards an ovule. The male nucleus travels along the pollen tube into the ovule.

(b) The male nucleus fuses with a nucleus in the ovule. This forms a zygote.

30 · Part 1 Core Sexual reproduction in plants [4]

Mark scheme

(a) **[2]**

- Exposed anthers allow pollen to be released into the wind. (1)
- Large feathery stigmas increase the chance of trapping airborne pollen. (1)

ALLOW anthers or stigmas hang outside the flower on long filaments or styles; production of large numbers of small, light pollen grains that are easily carried by wind

(b) **[2]**

- Oxygen must be available. (1)
- The temperature must be suitable for the species. (1)

Worked solution

(a) Exposed anthers allow pollen to be released into the wind. Large feathery stigmas increase the chance of trapping airborne pollen.

(b) Oxygen must be available. The temperature must be suitable for the species.

Topical Hub
topical-hub.pages.dev

1 · Part 2 Core Sexual reproduction in humans [1]

Answer: B

Explanation

The acrosome contains enzymes that help penetration. The flagellum provides movement and the nucleus carries genetic information.

2 · Part 2 Core Sexual reproduction in humans [1]

Answer: D

Explanation

The jelly coat changes after fertilisation, preventing further sperm entry. Nutrient loss and new motility structures are not the protective response.

3 · Part 2 Core Sexual reproduction in humans [1]

Answer: A

Explanation

Eggs are larger and have energy stores; sperm are small, motile and produced in much greater numbers.

4 · Part 2 Core Sexual reproduction in humans [1]

Answer: C

Explanation

The zygote divides to form an embryo, which implants in the uterine lining. Anthers are plant structures and implantation normally occurs in the uterus.

5 · Part 2 Extended Sexual reproduction in humans [1]

Answer: D

Explanation

Carbon dioxide and urea are fetal waste products. Oxygen and dissolved nutrients normally move towards the fetus.

Topical Hub
topical-hub.pages.dev

6 · Part 2 Extended Sexual reproduction in humans [1]

Answer: B

Explanation

The placenta allows exchange and is not a complete barrier to toxins and pathogens. Maternal and fetal blood are separate circulations.

7 · Part 2 Core Sex hormones in humans [1]

Answer: C

Explanation

Testosterone promotes male secondary sexual characteristics. Oestrogen promotes female secondary sexual characteristics; the other listed hormones have different reproductive roles.

8 · Part 2 Core Sex hormones in humans [1]

Answer: A

Explanation

Menstruation is the breakdown and shedding of the uterus lining when no embryo implants. Thickening of the lining happens after menstruation, in the first half of the cycle; release of an egg is ovulation, which happens in an ovary; and during menstruation progesterone is falling, not rising, which is why the lining is not maintained.

Topical Hub
topical-hub-pages.dev

9 · Part 2 Extended Sex hormones in humans [1]

Answer: B

Explanation

The rise in LH triggers ovulation. The other hormones do not provide this menstrual-cycle signal.

10 · Part 2 Extended Sex hormones in humans [1]

Answer: C

Explanation

The corpus luteum in the ovary produces progesterone after ovulation. The pituitary releases FSH and LH rather than progesterone.

11 · Part 2 Core Sexually transmitted infections [1]

Answer: D

Explanation

STIs are transmitted through sexual contact. Inherited disorders and nutritional conditions are not infections, and coughing alone does not define an STI.

12 · Part 2 Core Sexually transmitted infections [1]

Answer: A

Explanation

HIV is a virus and infection may progress to AIDS. AIDS is a condition, not a pathogen, and many STIs have other causes.

13 · Part 2 Core Sexually transmitted infections [1]

Answer: C

Explanation

Contaminated needles can transfer infected blood. Ordinary social contact such as handshaking or sharing desks does not transmit HIV.

14 · Part 2 Core Sexually transmitted infections [1]

Answer: A

Explanation

A condom provides a barrier to infected body fluids. Antibiotics do not prevent HIV, a viral infection; the other actions do not provide this barrier.

15 · Part 2 Core Sexually transmitted infections [1]

Answer: B

Explanation

HIV can be transmitted from an infected mother to her infant in breast milk (and also across the placenta or during birth). Hugging, breathing the same air and sharing cups or cutlery do not transmit HIV because the virus is carried in body fluids such as blood, semen and breast milk, not in air, saliva or skin contact.

Topical Hub
topical-hub.pages.dev

16 · Part 2 Core Sexual reproduction in humans [4]

Mark scheme

(a) **[2]**

- The sperm nucleus fuses with the egg nucleus. (1)
- The cell formed is a zygote. (1)

(b) **[2]**

- The flagellum propels the sperm. (1)
- Mitochondria release energy by respiration for movement. (1)

Worked solution

(a) The sperm nucleus fuses with the egg nucleus. The cell formed is a zygote.

(b) The flagellum propels the sperm. Mitochondria release energy by respiration for movement.

17 · Part 2 Core Sexual reproduction in humans [4]

Mark scheme

(a) **[2]**

- It supplies nutrients for early development. (1)
- The stored nutrients can be used in respiration to release energy. (1)

(b) **[2]**

- Sperm are motile whereas eggs do not swim using a flagellum. (1)
- Sperm are produced in much greater numbers than eggs. (1)

Worked solution

(a) It supplies nutrients for early development. The stored nutrients can be used in respiration to release energy.

(b) Sperm are motile whereas eggs do not swim using a flagellum. Sperm are produced in much greater numbers than eggs.

Topical Hub
topicalhub.pages.dev

18 · Part 2 Core Sexual reproduction in humans [4]

Mark scheme

(a) **[2]**

- The zygote divides repeatedly. (1)
- The embryo becomes a ball of cells. (1)

(b) **[2]**

- Implantation occurs in the lining of the uterus. (1)
- The embryo becomes embedded in that lining. (1)

Worked solution

(a) The zygote divides repeatedly. The embryo becomes a ball of cells.

(b) Implantation occurs in the lining of the uterus. The embryo becomes embedded in that lining.

19 · Part 2 Extended Sexual reproduction in humans [4]

Mark scheme

(a) **[2]**

- Glucose is transferred. (1)
- Amino acids are transferred. (1)

ALLOW any other named dissolved nutrient, e.g. mineral ions, vitamins, fatty acids

REJECT oxygen (a gas, not a nutrient)

(b) **[2]**

- Carbon dioxide moves towards the mother. (1)
- Urea moves towards the mother. (1)

Worked solution

(a) Glucose is transferred. Amino acids are transferred.

(b) Carbon dioxide moves towards the mother. Urea moves towards the mother.

Topical Hub
topical-hub.pages.dev

20 · Part 2 Extended Sexual reproduction in humans [4]

Mark scheme

(a) **[2]**

- The placenta is a site of exchange between maternal and fetal blood. (1)
- Blood in the umbilical cord carries exchanged substances between placenta and fetus. (1)

(b) **[2]**

- Some toxins can cross the placenta into fetal blood. (1)
- The toxin can therefore reach fetal tissues and affect development. (1)

Worked solution

(a) The placenta is a site of exchange between maternal and fetal blood. Blood in the umbilical cord carries exchanged substances between placenta and fetus.

(b) Some toxins can cross the placenta into fetal blood. The toxin can therefore reach fetal tissues and affect development.

21 · Part 2 Core Sexual reproduction in humans [4]

Mark scheme

(a) **[2]**

- The acrosome normally provides enzymes. (1)
- Reduced enzyme release makes penetration of the egg coverings less likely. (1)

(b) **[2]**

- Energy stores support early development. (1)
- The jelly coat changes after fertilisation to prevent additional sperm entering. (1)

Worked solution

(a) The acrosome normally provides enzymes. Reduced enzyme release makes penetration of the egg coverings less likely.

(b) Energy stores support early development. The jelly coat changes after fertilisation to prevent additional sperm entering.

Topical Hub
topical-hub.pages.dev

22 · Part 2 Core Sexual reproduction in humans [4]

Mark scheme

(a) **[2]**

- Sperm have a flagellum whereas eggs do not. (1)
- Eggs have much larger nutrient stores and cytoplasm than sperm. (1)

> **ALLOW** sperm have an acrosome / many mitochondria whereas eggs do not; eggs have a jelly coat; eggs are much larger than sperm

(b) **[2]**

- Many sperm are produced compared with relatively few eggs. (1)
- Sperm swim using their flagella whereas eggs do not swim independently. (1)

Worked solution

(a) Sperm have a flagellum whereas eggs do not. Eggs have much larger nutrient stores and cytoplasm than sperm.

(b) Many sperm are produced compared with relatively few eggs. Sperm swim using their flagella whereas eggs do not swim independently.

---

23 · Part 2 Core Sex hormones in humans [4]

Mark scheme

(a) **[2]**

- Testosterone is the hormone. (1)
- A valid example is deepening of the voice or growth of facial hair. (1)

(b) **[2]**

- Oestrogen is the hormone. (1)
- A valid example is development of breasts or widening of the hips. (1)

Worked solution

(a) Testosterone is the hormone. A valid example is deepening of the voice or growth of facial hair.

(b) Oestrogen is the hormone. A valid example is development of breasts or widening of the hips.

Topical Hub
topical-hub.pages.dev

24 · Part 2 Core Sex hormones in humans [4]

Mark scheme

(a) **[2]**

- An egg matures in a follicle. (1)
- Ovulation releases the egg around the middle of the cycle. (1)

ALLOW the empty follicle becomes a corpus luteum after ovulation

(b) **[2]**

- The lining thickens during the cycle. (1)
- It breaks down and is shed if pregnancy does not occur. (1)

Worked solution

(a) An egg matures in a follicle. Ovulation releases the egg around the middle of the cycle.

(b) The lining thickens during the cycle. It breaks down and is shed if pregnancy does not occur.

25 · Part 2 Extended Sex hormones in humans [4]

Mark scheme

(a) **[2]**

- The corpus luteum produces progesterone. (1)
- Progesterone maintains the uterine lining. (1)

(b) **[2]**

- It inhibits secretion of FSH and LH. (1)
- This reduces further follicle maturation and an additional ovulation. (1)

Worked solution

(a) The corpus luteum produces progesterone. Progesterone maintains the uterine lining.

(b) It inhibits secretion of FSH and LH. This reduces further follicle maturation and an additional ovulation.

Topical Hub
topical-hub.pages.dev

26 · Part 2 Extended Sex hormones in humans [4]

Mark scheme

(a) **[2]**

- FSH stimulates maturation of an ovarian follicle. (1)
- It stimulates oestrogen secretion by the developing follicle. (1)

(b) **[2]**

- Oestrogen stimulates rebuilding or thickening of the uterine lining. (1)
- A sufficiently high oestrogen concentration stimulates the LH surge that triggers ovulation. (1)

Worked solution

(a) FSH stimulates maturation of an ovarian follicle. It stimulates oestrogen secretion by the developing follicle.

(b) Oestrogen stimulates rebuilding or thickening of the uterine lining. A sufficiently high oestrogen concentration stimulates the LH surge that triggers ovulation.

27 · Part 2 Core Sexually transmitted infections [4]

Mark scheme

(a) **[2]**

- An STI is an infection transmitted through sexual contact. (1)
- HIV is one example of an STI pathogen. (1)

(b) **[2]**

- Education can encourage people to avoid risky sexual contact and seek testing. (1)
- Condoms form a barrier that reduces transfer of infected body fluids. (1)

Worked solution

(a) An STI is an infection transmitted through sexual contact. HIV is one example of an STI pathogen.

(b) Education can encourage people to avoid risky sexual contact and seek testing. Condoms form a barrier that reduces transfer of infected body fluids.

Topical Hub
topical-hub.pages.dev

28 · Part 2 Core Sexually transmitted infections [4]

Mark scheme

(a) **[2]**

- Screening reduces the chance of transfusing blood containing HIV. (1)
- Single-use needles prevent transfer of infected blood between patients. (1)

(b) **[2]**

- HIV may pass during birth. (1)
- HIV may pass through breast milk. (1)

ALLOW HIV may pass across the placenta before birth

Worked solution

(a) Screening reduces the chance of transfusing blood containing HIV. Single-use needles prevent transfer of infected blood between patients.

(b) HIV may pass during birth. HIV may pass through breast milk.

29 · Part 2 Core Sexually transmitted infections [4]

Mark scheme

(a) **[2]**

- HIV is a virus and a pathogen. (1)
- HIV infection may lead to the condition called AIDS. (1)

(b) **[2]**

- Transmission requires a suitable route involving infected body fluids. (1)
- Ordinary contact such as shaking hands does not provide that route. (1)

Worked solution

(a) HIV is a virus and a pathogen. HIV infection may lead to the condition called AIDS.

(b) Transmission requires a suitable route involving infected body fluids. Ordinary contact such as shaking hands does not provide that route.

Topical Hub
topical-hub.pages.dev

30 · Part 2 Core Sexually transmitted infections [4]

Mark scheme

(a) **[2]**

- Shared equipment may transfer infected blood. (1)
- Unprotected sexual contact may transfer infected sexual fluids. (1)

(b) **[2]**

- Use sterile single-use injection equipment and do not share it. (1)
- Use a condom correctly during sexual contact. (1)

Worked solution

(a) Shared equipment may transfer infected blood. Unprotected sexual contact may transfer infected sexual fluids.

(b) Use sterile single-use injection equipment and do not share it. Use a condom correctly during sexual contact.

Topical Hub
topical-hub.pages.dev