

## Differentiation

Imagine a bug that moves with constant speed on a circular path of radius  $r$  around the origin. The angle of the bug's position vector with the  $+x$  axis can be written as

$$\theta = \omega t + a.$$

Assume  $a = 0$ , so that the bug is on the  $+x$  axis at time 0. Then the position vector of the bug is

$$X(t) = (r \cos(\omega t), r \sin(\omega t)).$$

Now imagine the bug lives in  $\mathbb{R}^3$  with

$$X(t) = (\cos(t), \sin(t), t).$$

This lifts the circular path into a helix.

In general, a **parametrized curve**  $X : I \rightarrow \mathbb{R}^n$  is a vector-valued function that maps points from an interval  $I$  into  $n$ -space. In the examples above,  $I$  is the entire real line  $\mathbb{R}$  (which we consider to be an interval). We can write  $X(t)$  as its individual coordinate functions

$$X(t) = (x_1(t), \dots, x_n(t)).$$

Just as with ordinary real-valued function, we can take derivatives by looking at the limit

$$\lim_{h \rightarrow 0} \frac{X(t+h) - X(t)}{h}.$$

Here, dividing by  $h$  really means scaling the vector by  $1/h$ . Writing out components, this is simply

$$\lim_{h \rightarrow 0} \frac{(x_1(t+h) - x_1(t), \dots, x_n(t+h) - x_n(t))}{h}.$$

If the individual components are all differentiable, we obtain a new vector-valued function

$$X'(t) = (x'_1(t), \dots, x'_n(t)).$$

$X'(t)$  is called the **derivative** or **velocity** of  $X(t)$ .

So for the example  $X(t) = (\cos(t), \sin(t), t)$ , we have

$$X'(t) = (-\sin(t), \cos(t), 1).$$

The velocity is parallel to the direction of instantaneous motion.

**Example.** Find a parametric equation of the tangent line to the curve  $X(t) = (\sin t, \cos t)$  at  $t = \pi/3$ .

We need two pieces of information: a point on the line, and a direction vector of the line. These are supplied by  $X(\pi/3)$  and  $X'(\pi/3)$  respectively. The tangent line  $L(t)$  can

thus be written

$$\begin{aligned} L(s)|_{t=\pi/3} &= X(\pi/3) + sX'(\pi/3) \\ &= \left( \frac{\sqrt{3}}{2} + \frac{1}{2}s, \frac{1}{2} - \frac{\sqrt{3}}{2}s \right). \end{aligned}$$

We used the parameter  $s$  for the line to avoid confusion with the already defined  $X(t)$  above.

The **speed** of the curve  $X(t)$ , denoted  $v(t)$ , is defined to be

$$v(t) = \|X'(t)\|.$$

**acceleration** is the second derivative  $X''(t)$ .