

Final Exam/Homework

Exercise 1

Let $T : \mathbb{P}_n \rightarrow \mathbb{P}_n$ be the linear transformation defined by $T(p(t)) = 2p(t) + 3p'(t) - 4f''(t)$. Find the matrix of T with respect to the basis $1, t, t^2, \dots, t^n$ of $\mathbb{P}_n$. You can express the matrix by specifying what each $a_{i,j}$ is, where $a_{i,j}$ is the entry in the i -th row and j -th column of the matrix (or whatever is easiest to write, as long as you give a complete description).

Exercise 2

Find the change of coordinates matrix from the basis $1, 1+t$ to the basis $1-t, 2t$ in $\mathbb{P}_1$.

Exercise 3

Let $Perm(n)$ be the set of permutations of $\{1, 2, \dots, n\}$. Let $\tau \in Perm(n)$. Show that the map from $Perm(n)$ to $Perm(n)$ defined by $\sigma \mapsto \sigma\tau$ is a bijection (one-to-one and onto). In other words, composition with a fixed permutation τ takes the list of all permutations and rearranges it, but does not duplicate or remove any (it permutes the permutations).

Exercise 4

Show that the determinant of a matrix is the product of its eigenvalues (counting multiplicities). **Hint:** First show that the characteristic polynomial $\det(A - \lambda I) = (\lambda_1 - \lambda)(\lambda_2 - \lambda) \cdots (\lambda_n - \lambda)$ (note the λ_i 's occur before λ), where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of A (with multiplicity). Then plug in $\lambda = 0$. Note that in general, a polynomial of degree n can be written as $p(\lambda) = c(\lambda - r_1)(\lambda - r_2) \cdots (\lambda - r_n)$ (note the λ comes before each λ_i), so the first step is to show that this c is $(-1)^n$. If you use the permutation definition of the determinant, there is exactly one term that is of degree n in λ , and it comes from the identity permutation.

Exercise 5

Suppose the entries of A and A^{-1} are integers. What are the possible values of $\det(A)$? **Hint:** Use the fact that $\det(A) \cdot \det(A^{-1}) = 1$.

Exercise 6

A matrix A is called *nilpotent* if $A^k = 0$ for some positive integer k . Show that the determinant of a nilpotent matrix is 0.

Exercise 7

How are the determinants of A and B related if

(a)

$$A = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix}, B = \begin{pmatrix} 2a_1 & 3a_2 & 5a_3 \\ 2b_1 & 3b_2 & 5b_3 \\ 2c_1 & 3c_2 & 5c_3 \end{pmatrix}$$

(b)

$$A = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix}, B = \begin{pmatrix} 3a_1 & 4a_2 + 5a_1 & 5a_3 \\ 3b_1 & 4b_2 + 5b_1 & 5b_3 \\ 3c_1 & 4c_2 + 5c_1 & 5c_3 \end{pmatrix}$$

Exercise 8

Recall that the Fibonacci sequence is defined by $\varphi_0 = 0$, $\varphi_1 = 1$, and $\varphi_{n+2} = \varphi_{n+1} + \varphi_n$ for $n \geq 0$ (the first few terms are 0, 1, 1, 2, 3, 5, 8, 13, ...). You will now find a formula for the n -th Fibonacci number.

- (a) Find a 2×2 matrix A such that

$$\begin{pmatrix} \varphi_{n+2} \\ \varphi_{n+1} \end{pmatrix} = A \begin{pmatrix} \varphi_{n+1} \\ \varphi_n \end{pmatrix}.$$

- (b) Diagonalize A and find a formula for A^n .

- (c) Noticing that

$$\begin{pmatrix} \varphi_{n+1} \\ \varphi_n \end{pmatrix} = A^n \begin{pmatrix} \varphi_1 \\ \varphi_0 \end{pmatrix} = A^n \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

use your formula for A^n to find a formula for φ_n .