
```

% Zain Bhaila
% Math 401
% Special Assignment #2

syms a b c d e;
% transition matrix given by problem
% ab,c,d,e are the values for teh transitions from S5
M = [3/12 5/12 0/12 2/12 a ;
      5/12 7/12 2/12 2/12 b ;
      3/12 0/12 8/12 2/12 c ;
      0/12 0/12 0/12 4/12 d ;
      1/12 0/12 2/12 2/12 e ]

% combinations with repetition
% number of ways to divide 12 items into 5 groups
V = 0:12;
N = 5;
[Y{N:-1:1}] = ndgrid(1:numel(V));
I = reshape(cat(N+1, Y{:}), [], N);
P = V(I); % contains every possible permutation with repetition with
           range 0-12 of 5 selections
[h, w] = size(P) % dimensions of P, h is the number of permutations
F = []; % filter P so that total tickets is 12
for i = 1:h % for each permutation
    if P(i,1) + P(i,2) + P(i,3) + P(i, 4) + P(i,5) == 12 % if the
        total # of tickets is twelve, keep the row
        F = [F;P(i,:)]; %#ok<*AGROW>
    end
end
[h, w] = size(F) % dimensions of F, h is the number of permutations
               where the total ticket amount is 12

min = 2147483647; % store minimum variance, set to max int to start
               (2^32 - 1)
loc = 0; % row of F with minimum variance distribution of tickets
S = []; % steady state
for i = 1:h % for each permutation in F
    Q = M; % set Q to M
    Q(1,5) = F(i,1)/12; % set values of Q to reflect current ticket
    distribution
    Q(2,5) = F(i,2)/12;
    Q(3,5) = F(i,3)/12;
    Q(4,5) = F(i,4)/12;
    Q(5,5) = F(i,5)/12;
    T = Q^4; % 4 is the longest possible path between states
    if all(T, 'all') % if T has no 0 values
        [V,D] = eig(Q); % get eigenvalue/eigenvector set
        [row,col] = find(D == 1); % index of eigenvalue = 1, row will
        be the same as col
        V(:,col) = V(:,col)/sum(V(:,col)); % divide corresponding
        eigenvector by sum of its values
        m = (V(1,col) + V(2,col) + V(3,col) + V(4,col) + V(5,col))/5;
    end
end

```

```

        s = (V(1,col) - m)^2 + (V(2,col) - m)^2 + (V(3,col) - m)^2 +
        (V(4,col) - m)^2 + (V(5,col) - m)^2;
        if (s < min) % if this variance is the new minimum
            min = s; % store variance as new minimum
            loc = i; % save the row this occurs at
            S = V(:,col); % save the steady state
        end
    end
end

% output answers
min % minimum variance
F(loc,:) % ticket distribution to S1-S5 from S5 in order
S % steady state
Q = M;
Q(1,5) = F(loc,1)/12;
Q(2,5) = F(loc,2)/12;
Q(3,5) = F(loc,3)/12;
Q(4,5) = F(loc,4)/12;
Q(5,5) = F(loc,5)/12 % final transition matrix
Q^4 % proof of regularity
% since the matrix is regular, it will always converge to the same
    steady
% state corresponding to eigenvalue = 1

M =

[ 1/4, 5/12, 0, 1/6, a]
[ 5/12, 7/12, 1/6, 1/6, b]
[ 1/4, 0, 2/3, 1/6, c]
[ 0, 0, 0, 1/3, d]
[ 1/12, 0, 1/6, 1/6, e]

h =

    371293

w =

    5

h =

    1820

w =

    5

```

`min =`

`3076/132845`

`ans =`

`0 0 0 3 9`

`S =`

`30/163`

`48/163`

`30/163`

`15/163`

`40/163`

`Q =`

`[1/4, 5/12, 0, 1/6, 0]`

`[5/12, 7/12, 1/6, 1/6, 0]`

`[1/4, 0, 2/3, 1/6, 0]`

`[0, 0, 0, 1/3, 1/4]`

`[1/12, 0, 1/6, 1/6, 3/4]`

`ans =`

`[2233/10368, 155/576, 707/5184, 101/576, 337/3456]`

`[403/1152, 2189/5184, 20/81, 491/1728, 481/3456]`

`[89/432, 625/3456, 1315/5184, 19/96, 197/1728]`

`[89/1728, 145/6912, 13/144, 31/324, 1435/6912]`

`[919/5184, 2215/20736, 707/2592, 20/81, 3053/6912]`

Published with MATLAB® R2019a