Results First Draft

Zachary Serby

November 13, 2026

Main

Summary Statistics

Table 1—Summary Statistics, Discrete-Dose Samples

	Panel A: WAZ, age ≤10 in 2005 and 2009					
	Mean	Std. dev.	Min.	Median	Max.	Obs.
Outcomes (MxFLS-2, 2005)						
Weight-for-age z-score (WAZ)	-0.30	1.42	-5.07	-0.36	4.51	1,045
Outcome change, 2005 to 2009						
Change in WAZ	0.55	1.41	-4.88	0.46	9.10	1,045
Dose levels (CHIRPS)						
2009 rainfall, z	-0.50	0.83	-2.24	-0.39	0.96	1,045
Near normal ($-1 < z < 1$)	0.65	0.48	0.00	1.00	1.00	1,045
Moderately dry ($-1.5 < z \leq -1$)	0.21	0.41	0.00	0.00	1.00	1,045
Severely dry or worse ($z \leq -1.5$)	0.14	0.35	0.00	0.00	1.00	1,045

	Panel B: HAZ and BMIZ, age ≤15 in 2005 and 2009					
	Mean	Std. dev.	Min.	Median	Max.	Obs.
Outcomes (MxFLS-2, 2005)						
Height-for-age z-score (HAZ)	-0.58	1.49	-5.46	-0.63	5.65	2,351
BMI-for-age z-score (BMIZ)	0.36	1.43	-4.95	0.30	4.86	2,310
Outcome change, 2005 to 2009						
Change in HAZ	0.07	1.37	-5.93	0.07	6.35	2,351
Change in BMIZ	0.23	1.31	-5.90	0.17	7.59	2,295
Dose levels (CHIRPS)						
2009 rainfall, z	-0.55	0.84	-2.24	-0.53	0.96	2,351
Near normal ($-1 < z < 1$)	0.64	0.48	0.00	1.00	1.00	2,351
Moderately dry ($-1.5 < z \leq -1$)	0.21	0.41	0.00	0.00	1.00	2,351
Severely dry or worse ($z \leq -1.5$)	0.16	0.36	0.00	0.00	1.00	2,351

Notes: This table is for the discrete-dose specification, the main model. Obs. counts children. The sample is children with the outcome measured in MxFLS-2 (2005) and MxFLS-3 (2009), under the panel's age cutoff in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009. I use November 2009 as the cutoff because the 2009 rainfall deficit had not fully occurred before then. Dropping them from both groups equalizes treated and control measurement timing. Children in wet municipalities (z of 1 or above) are excluded. Panel A uses the WAZ sample, Panel B the HAZ and BMIZ sample. The BMIZ sample is marginally smaller than the HAZ sample shown in Panel B, at 2,300 children. Statistics are at the 2005 baseline, except the change in WAZ, HAZ and BMIZ from 2005 to 2009, the outcome of the discrete-dose regressions, and 2009 rainfall. z is 2009 May–October CHIRPS rainfall, the maize growing season (Jessoe, Manning, and Taylor 2018), in the child's 2005 municipality in standard deviations from its 1981–2004 mean, grouped into dose levels with the SPI cutoffs (McKee, Doesken, and Kleist 1993; WMO 2012). The 2009 rainfall z is continuous. The three dose-group indicators have means equal to the share of children in each dose group. The discrete-dose regressions have no controls, as the contdid package does not yet accommodate covariates (Callaway, Goodman-Bacon, and Sant'Anna 2026).

Regression Results

Table 2—2009 Rainfall Deficits and Child Anthropometrics, Discrete Dose

	HAZ		WAZ		BMIZ	
	(1)	(2)	(3)	(4)	(5)	(6)
Dry ($z \leq -1$)	-0.041		-0.197		-0.163**	
	(0.100)		(0.117)		(0.076)	
	[0.704]		[0.104]		[0.036]	
Moderately dry ($-1.5 < z \leq -1$)		0.018		-0.230		-0.221*
		(0.139)		(0.164)		(0.106)
		[0.915]		[0.220]		[0.062]
Severely dry or worse ($z \leq -1.5$)		-0.117		-0.148		-0.086
		(0.112)		(0.119)		(0.068)
		[0.308]		[0.238]		[0.218]
P-val: pre-trend 2002–2005, moderately dry		0.933		0.393		0.230
P-val: pre-trend 2002–2005, severely dry or worse		0.791		0.237		0.040
Children	2,351	2,351	1,045	1,045	2,300	2,300
Dry 2005 municipalities	40		39		40	
Near-normal 2005 municipalities	89	89	80	80	89	89

Notes: This table compares children in dry 2005 municipalities with children in near-normal ones. Regressions are at the child level, on the 2005 to 2009 change in the WHO z-score, for the binary DID estimation samples without children in wet municipalities (z of 1 or above). z is 2009 May–October CHIRPS rainfall, the maize growing season (Jessoe, Manning, and Taylor 2018), in the child's 2005 municipality in standard deviations from its 1981–2004 mean, grouped into dose levels with the SPI cutoffs. The SPI cutoffs are near normal for z between -1 and 1, moderately dry between -1.5 and -1, and severely dry or worse for -1.5 or below (McKee, Doesken, and Kleist 1993; WMO 2012). Cols. 1, 3 and 5 compare dry (z of -1 or below) with near normal, and Cols. 2, 4 and 6 split dry into moderately dry and severely dry or worse. Each dose group is only compared with near normal, which needs parallel trends, and not with the other dose group, which would need strong parallel trends (Callaway, Goodman-Bacon, and Sant'Anna 2024). Since the contdid package does not yet support discrete doses, the dose groups are estimated with the discrete-dose regression in equation 4.1 (Callaway, Goodman-Bacon, and Sant'Anna 2024). The regressions have no controls, as the contdid package does not yet accommodate covariates (Callaway, Goodman-Bacon, and Sant'Anna 2026). The pre-trend p-values test the 2002 leads of the discrete-dose event study, on the same children with child and wave fixed effects. Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are in brackets, which tend to be conservative with few treated municipalities (MacKinnon, Nielsen, and Webb 2023). Stars use the clustered p-values, and where a wild bootstrap p-value is shown both must pass. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 3—2009 Drought and Height-for-Age, Binary DID

	Height-for-age z-score (HAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.072	-0.035	0.021	0.101	-0.005
	(0.096)	(0.103)	(0.113)	(0.168)	(0.118)
Drought × post × 2002 remittances					-0.153
					(0.148)
2002 remittances × post					-0.165*
					(0.090)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	2,432	2,432	1,755	734	1,622

Notes: This table reports the effect of the 2009 drought on the WHO height-for-age z-score (HAZ) with the binary DID. Regressions are at the child-wave level. The sample is children with the outcome measured in both MxFLS-2 (2005) and MxFLS-3 (2009), aged 15 or younger in both survey waves. Cols. 3–5 also keep only children with all of their regressors observed. Children have a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Drought equals 1 if May–October CHIRPS rainfall in 2009, the maize growing season (Jessoe, Manning, and Taylor 2018), in the child's 2005 municipality was below its 1981–2004 20th percentile (Shah and Steinberg 2017). Post equals 1 in MxFLS-3. Remittances equal 1 if the household received cash in the last 12 months from parents, siblings, or children living outside the household in MxFLS-1 (2002), a proxy for remittances. Cols. 1–4 report the average drought effect. Col. 1 is the basic DID, Col. 2 adds the 2005–2008 droughts, and Col. 3 adds 2002 household controls (electricity, floor, roof, ventilation, sleeping rooms per person, log expenditure per capita, rural) and 2005 child and mother controls (child sex and full vaccination, mother's height, schooling, age at birth and employment). Col. 4 adds the 2005 father and birth controls (father's height, schooling, age at birth and employment, birth order, oldest child, birth weight, delivered at home, weeks predominantly breastfed, iron in pregnancy, premature), which are missing for many children, often because the father migrated. All controls are interacted with post. Col. 5 is the triple difference: it keeps the controls of Col. 3 and adds 2002 remittances × post and drought × post × 2002 remittances, the difference in the drought effect for children in 2002 remittance households. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). All regressions include child and interview year-month fixed effects, children in singleton cells are dropped, and standard errors are clustered by 2005 municipality, where the drought is assigned. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 4—2009 Drought and Weight-for-Age, Binary DID

	Weight-for-age z-score (WAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.200**	-0.196**	-0.299**	-0.181	-0.119
	(0.101)	(0.095)	(0.125)	(0.193)	(0.147)
Drought × post × 2002 remittances					-0.376*
					(0.205)
2002 remittances × post					0.096
					(0.119)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,078	1,078	700	224	634

Notes: This table reports the effect of the 2009 drought on the WHO weight-for-age z-score (WAZ) with the binary DID. Regressions are at the child-wave level. The sample is children with the outcome measured in both MxFLS-2 (2005) and MxFLS-3 (2009), aged 10 or younger in both survey waves. Cols. 3–5 also keep only children with all of their regressors observed. Children have a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Drought equals 1 if May–October CHIRPS rainfall in 2009, the maize growing season (Jessoe, Manning, and Taylor 2018), in the child's 2005 municipality was below its 1981–2004 20th percentile (Shah and Steinberg 2017). Post equals 1 in MxFLS-3. Remittances equal 1 if the household received cash in the last 12 months from parents, siblings, or children living outside the household in MxFLS-1 (2002), a proxy for remittances. Cols. 1–4 report the average drought effect. Col. 1 is the basic DID, Col. 2 adds the 2005–2008 droughts, and Col. 3 adds 2002 household controls (electricity, floor, roof, ventilation, sleeping rooms per person, log expenditure per capita, rural) and 2005 child and mother controls (child sex and full vaccination, mother's height, schooling, age at birth and employment). Col. 4 adds the 2005 father and birth controls (father's height, schooling, age at birth and employment, birth order, oldest child, birth weight, delivered at home, weeks predominantly breastfed, iron in pregnancy, premature), which are missing for many children, often because the father migrated. All controls are interacted with post. Col. 5 is the triple difference: it keeps the controls of Col. 3 and adds 2002 remittances × post and drought × post × 2002 remittances, the difference in the drought effect for children in 2002 remittance households. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). All regressions include child and interview year-month fixed effects, children in singleton cells are dropped, and standard errors are clustered by 2005 municipality, where the drought is assigned. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 5—2009 Drought and BMI-for-Age, Binary DID

	BMI-for-age z-score (BMIZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.145**	-0.180**	-0.304***	-0.212*	-0.266**
	(0.071)	(0.087)	(0.095)	(0.112)	(0.110)
Drought × post × 2002 remittances					0.027
					(0.217)
2002 remittances × post					0.010
					(0.074)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	2,379	2,379	1,714	712	1,587

Notes: This table reports the effect of the 2009 drought on the WHO BMI-for-age z-score (BMIZ) with the binary DID. Regressions are at the child-wave level. The sample is children with the outcome measured in both MxFLS-2 (2005) and MxFLS-3 (2009), aged 15 or younger in both survey waves. Cols. 3–5 also keep only children with all of their regressors observed. Children have a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Drought equals 1 if May–October CHIRPS rainfall in 2009, the maize growing season (Jessoe, Manning, and Taylor 2018), in the child's 2005 municipality was below its 1981–2004 20th percentile (Shah and Steinberg 2017). Post equals 1 in MxFLS-3. Remittances equal 1 if the household received cash in the last 12 months from parents, siblings, or children living outside the household in MxFLS-1 (2002), a proxy for remittances. Cols. 1–4 report the average drought effect. Col. 1 is the basic DID, Col. 2 adds the 2005–2008 droughts, and Col. 3 adds 2002 household controls (electricity, floor, roof, ventilation, sleeping rooms per person, log expenditure per capita, rural) and 2005 child and mother controls (child sex and full vaccination, mother's height, schooling, age at birth and employment). Col. 4 adds the 2005 father and birth controls (father's height, schooling, age at birth and employment, birth order, oldest child, birth weight, delivered at home, weeks predominantly breastfed, iron in pregnancy, premature), which are missing for many children, often because the father migrated. All controls are interacted with post. Col. 5 is the triple difference: it keeps the controls of Col. 3 and adds 2002 remittances × post and drought × post × 2002 remittances, the difference in the drought effect for children in 2002 remittance households. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). All regressions include child and interview year-month fixed effects, children in singleton cells are dropped, and standard errors are clustered by 2005 municipality, where the drought is assigned. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 6—Dose Groups and Baseline Characteristics, BMIZ Sample

	(1) Electricity	(2) Good floor	(3) Good roof	(4) Enough ventilation	(5) Rooms per person	(6) Log expenditure	(7) Expenditure (pesos)	(8) Rural	(9) PROGRESA	(10) Remittances	(11) Male	(12) Age	(13) Fully vaccinated	(14) Mother's height (cm)	(15) Mother's schooling	(16) Mother's age at birth
Moderately dry	-0.048*	-0.038	-0.048	0.101*	-0.026	-0.078	293.396	0.038	0.095	-0.087*	0.018	-0.146	0.034	0.526	-0.600	0.818
	(0.028)	(0.074)	(0.060)	(0.053)	(0.027)	(0.168)	(628.937)	(0.165)	(0.094)	(0.049)	(0.025)	(0.160)	(0.054)	(1.051)	(0.641)	(0.507)
Severely dry or worse	-0.011	0.104**	0.065	0.080	0.071**	0.204	433.655	-0.257*	-0.120**	-0.087*	-0.012	0.169	0.074	0.575	1.181**	0.100
	(0.010)	(0.042)	(0.074)	(0.069)	(0.033)	(0.163)	(590.207)	(0.134)	(0.059)	(0.045)	(0.029)	(0.228)	(0.054)	(1.610)	(0.528)	(0.537)
Observations	1,914	1,914	1,914	1,899	1,911	1,886	1,886	1,914	1,870	1,734	2,300	2,300	2,184	2,172	2,199	2,219
Mean of dep. var.	0.981	0.828	0.808	0.565	0.385	6.514	1302.934	0.466	0.228	0.369	0.494	5.918	0.482	154.329	7.190	26.386
Fixed effects	None															

	(17) Mother employed	(18) Father's height (cm)	(19) Father's schooling	(20) Father's age at birth	(21) Father employed	(22) Birth order	(23) Oldest child	(24) Birth weight (kg)	(25) Born at home	(26) Weeks breastfed	(27) Iron in pregnancy	(28) Premature	(29) BMIZ	(30) Long-run rain mean	(31) Long-run rain SD
Moderately dry	0.021	-0.467	-0.487	1.152*	0.007	0.030	-0.027	-0.010	0.085	-1.959	-0.035	0.013	0.007	123.123	1.368
	(0.041)	(1.028)	(0.645)	(0.587)	(0.016)	(0.258)	(0.038)	(0.049)	(0.085)	(3.837)	(0.072)	(0.022)	(0.158)	(133.660)	(22.653)
Severely dry or worse	0.071	0.310	0.852	-0.073	0.021	-0.345*	0.034	0.003	-0.103***	-11.307***	0.124**	0.006	0.178	195.378**	2.366
	(0.060)	(1.432)	(0.657)	(0.780)	(0.015)	(0.206)	(0.034)	(0.048)	(0.027)	(3.017)	(0.049)	(0.030)	(0.120)	(89.347)	(17.421)
Observations	2,199	1,527	1,820	1,884	1,824	1,640	1,640	1,496	1,823	1,611	1,621	1,617	2,300	2,300	2,300
Mean of dep. var.	0.288	166.426	7.398	29.894	0.958	2.551	0.359	3.254	0.111	33.460	0.745	0.140	0.354	723.185	123.855
Fixed effects	None														

Notes: The sample is the 2,300 children in the BMIZ discrete-dose regressions. Each column regresses a baseline characteristic on indicators for moderately dry ($-1.5 < z \leq -1$) and severely dry or worse ($z \leq -1.5$) municipalities. Each coefficient is the mean difference in the characteristic between children in that dry dose group and children in the near-normal dose group ($-1 < z < 1$). Columns 1–10 are from MxFLS-1 (2002) and columns 11–29 from MxFLS-2 (2005). Of particular interest are the measures of household poverty. Expenditure per person is a proxy for income and is shown in logs and in 2002 pesos per month. PROGRESA equals 1 if any household member was in the program in the last 12 months, which marks the household as poor by the program's targeting. The dwelling characteristics also reflect household poverty. Long-run rainfall is the municipality's 1981–2004 May–October mean and SD in millimeters. The table has no fixed effects because the discrete-dose regressions use each child's 2005-to-2009 change without fixed effects or covariates. Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are not reported. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Figure 1—2009 Rainfall by Municipality, Dose Levels






Figure 2—Event Study by Dose Level, BMI-for-Age, Discrete Dose




Notes: Moderately dry and severely dry or worse compared with near-normal. The 2009 coefficients are the same as in the discrete-dose table. Here I include both the 2002 lead and the 2009 coefficient. Similar to the discrete-dose table, there are no controls. Included are child and survey wave fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 3—Event Study by Dose Level, Height-for-Age, Discrete Dose




Notes: Moderately dry and severely dry or worse compared with near-normal. The 2009 coefficients are the same as in the discrete-dose table. Here I include both the 2002 lead and the 2009 coefficient. Similar to the discrete-dose table, there are no controls. Included are child and survey wave fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 4—Event Study by Dose Level, Weight-for-Age, Discrete Dose




Notes: Moderately dry and severely dry or worse compared with near-normal. The 2009 coefficients are the same as in the discrete-dose table. Here I include both the 2002 lead and the 2009 coefficient. Similar to the discrete-dose table, there are no controls. Included are child and survey wave fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 5—Event Study by Dose Level, BMI-for-Age, Discrete Dose, Children in All Three Waves



Notes: Moderately dry and severely dry or worse compared with near-normal. The sample is the children in the discrete-dose table who were measured in all three survey waves, so the 2002 lead and the 2009 coefficient come from the same children. Similar to the discrete-dose table, there are no controls. Included are child and survey wave fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 6—Overlap of Baseline Characteristics, BMI-for-Age, Dose Groups



Notes: The sample is the children in the discrete-dose regressions. Kernel densities of the propensity score. That is, the predicted probability of being in the dry dose group from a logit on baseline characteristics. Baseline characteristics are the 2002 household controls (electricity, floor, roof, ventilation, rooms per person, log expenditure per person, rural) and the 2005 child and mother controls (sex, full vaccination, mother's height, schooling, age at birth, and employment), and the child's age. Missing values are set to the sample mean and flagged with an indicator. Each panel compares one dry dose group with near-normal ($-1 < z < 1$) and is estimated separately. Moderately dry is $-1.5 < z \leq -1$, and severely dry or worse is $z \leq -1.5$. In each box, children and mean score are shown as dry dose group/near-normal. Outside support is the number of dry dose group children whose score lies below the lowest or above the highest near-normal score.

Appendix

Table 7—Summary Statistics, Binary DID Model 3 Samples

	Panel A: WAZ, age ≤10 in 2005 and 2009					
	Mean	Std. dev.	Min.	Median	Max.	Obs.
Outcomes (MxFLS-2, 2005)						
Weight-for-age z-score (WAZ)	-0.31	1.40	-4.92	-0.34	4.51	700
Binary DID treatment (CHIRPS)						
2009 drought (indicator)	0.36	0.48	0.00	0.00	1.00	700
2005 drought (indicator)	0.68	0.47	0.00	1.00	1.00	700
2006 drought (indicator)	0.03	0.17	0.00	0.00	1.00	700
2007 drought (indicator)	0.15	0.36	0.00	0.00	1.00	700
2008 drought (indicator)	0.13	0.34	0.00	0.00	1.00	700
Remittances (MxFLS-1, 2002)						
Household received remittances, proxy (indicator)	0.41	0.49	0.00	0.00	1.00	634
Household (MxFLS-1, 2002)						
Dwelling has electricity (indicator)	0.97	0.16	0.00	1.00	1.00	700
Good floor quality (indicator)	0.85	0.36	0.00	1.00	1.00	700
Good roof quality (indicator)	0.81	0.39	0.00	1.00	1.00	700
Enough ventilation (indicator)	0.57	0.50	0.00	1.00	1.00	700
Sleeping rooms per household member	0.41	0.25	0.11	0.33	2.33	700
Ln (monthly expenditure per capita, pesos), proxy	6.56	0.86	4.21	6.57	11.43	700
Rural locality (indicator)	0.43	0.50	0.00	0.00	1.00	700
Child and mother (MxFLS-2, 2005)						
Age (years, not in regression)	2.52	1.68	0.00	3.00	6.00	700
Male (indicator)	0.53	0.50	0.00	1.00	1.00	700
Fully vaccinated (indicator)	0.39	0.49	0.00	0.00	1.00	700
Mother's height (centimeters)	154.79	6.91	136.00	154.50	181.00	700
Mother's years of schooling	7.40	3.42	0.00	8.00	18.00	700
Mother's age at birth (years)	26.93	6.52	14.00	27.00	59.00	700
Mother employed (indicator)	0.24	0.42	0.00	0.00	1.00	700
Father and birth (MxFLS-1 and MxFLS-2)						
Father's height (centimeters)	166.81	6.98	144.00	167.00	189.40	465
Father's years of schooling	7.51	3.77	0.00	8.00	18.00	569
Father's age at birth (years)	30.74	7.82	16.00	30.00	75.00	581
Father employed (indicator)	0.96	0.19	0.00	1.00	1.00	569
Birth order	2.31	1.68	1.00	2.00	13.00	401
Oldest child (indicator)	0.40	0.49	0.00	0.00	1.00	401
Birth weight (kilograms)	3.26	0.59	1.55	3.20	5.50	523
Delivered at home (indicator)	0.08	0.27	0.00	0.00	1.00	560
Weeks predominantly breastfed	31.98	29.20	0.00	26.09	157.29	542
Iron during pregnancy (indicator)	0.76	0.43	0.00	1.00	1.00	549
Premature, mother reported (indicator)	0.19	0.39	0.00	0.00	1.00	546

	Panel B: HAZ and BMIZ, age ≤15 in 2005 and 2009					
	Mean	Std. dev.	Min.	Median	Max.	Obs.
Outcomes (MxFLS-2, 2005)						
Height-for-age z-score (HAZ)	-0.60	1.47	-5.17	-0.62	5.20	1,755
BMI-for-age z-score (BMIZ)	0.40	1.41	-4.95	0.35	4.86	1,721
Binary DID treatment (CHIRPS)						
2009 drought (indicator)	0.39	0.49	0.00	0.00	1.00	1,755
2005 drought (indicator)	0.66	0.47	0.00	1.00	1.00	1,755
2006 drought (indicator)	0.03	0.17	0.00	0.00	1.00	1,755
2007 drought (indicator)	0.16	0.36	0.00	0.00	1.00	1,755
2008 drought (indicator)	0.14	0.34	0.00	0.00	1.00	1,755
Remittances (MxFLS-1, 2002)						
Household received remittances, proxy (indicator)	0.36	0.48	0.00	0.00	1.00	1,624
Household (MxFLS-1, 2002)						
Dwelling has electricity (indicator)	0.98	0.15	0.00	1.00	1.00	1,755
Good floor quality (indicator)	0.83	0.37	0.00	1.00	1.00	1,755
Good roof quality (indicator)	0.81	0.39	0.00	1.00	1.00	1,755
Enough ventilation (indicator)	0.56	0.50	0.00	1.00	1.00	1,755
Sleeping rooms per household member	0.38	0.23	0.11	0.33	2.33	1,755
Ln (monthly expenditure per capita, pesos), proxy	6.52	0.86	3.75	6.51	12.24	1,755
Rural locality (indicator)	0.46	0.50	0.00	0.00	1.00	1,755
Child and mother (MxFLS-2, 2005)						
Age (years, not in regression)	6.08	3.22	0.00	6.00	12.00	1,755
Male (indicator)	0.50	0.50	0.00	0.00	1.00	1,755
Fully vaccinated (indicator)	0.50	0.50	0.00	0.00	1.00	1,755
Mother's height (centimeters)	154.39	7.14	133.40	154.20	190.00	1,755
Mother's years of schooling	7.13	3.66	0.00	6.00	18.00	1,755
Mother's age at birth (years)	26.80	6.46	14.00	26.00	59.00	1,755
Mother employed (indicator)	0.28	0.45	0.00	0.00	1.00	1,755
Father and birth (MxFLS-1 and MxFLS-2)						
Father's height (centimeters)	166.42	7.15	144.00	166.30	192.00	1,212
Father's years of schooling	7.33	3.92	0.00	7.00	18.00	1,443
Father's age at birth (years)	30.36	7.85	15.00	29.00	75.00	1,482
Father employed (indicator)	0.96	0.20	0.00	1.00	1.00	1,447
Birth order	2.66	1.89	1.00	2.00	13.00	1,392
Oldest child (indicator)	0.32	0.47	0.00	0.00	1.00	1,392
Birth weight (kilograms)	3.27	0.60	0.85	3.25	6.00	1,256
Delivered at home (indicator)	0.12	0.32	0.00	0.00	1.00	1,544
Weeks predominantly breastfed	33.57	33.98	0.00	26.09	260.89	1,357
Iron during pregnancy (indicator)	0.72	0.45	0.00	1.00	1.00	1,366
Premature, mother reported (indicator)	0.14	0.35	0.00	0.00	1.00	1,362

Notes: This table is for the sample of Model 3, the main binary DID with controls. Obs. counts children. The children have the outcome measured in MxFLS-2 (2005) and MxFLS-3 (2009), under the panel's age cutoff in both survey waves, and a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Panel A uses the WAZ sample, Panel B the HAZ and BMIZ sample. The BMIZ sample is marginally smaller than the HAZ sample shown in Panel B, at 1,714 children. Statistics are at the 2005 baseline, except remittances and household variables (2002). Remittances enter Model 5, and the father and birth controls enter Model 4. Both are observed for fewer children. Drought is May–October CHIRPS rainfall, the maize growing season (Jessoe, Manning, and Taylor 2018), in the 2005 municipality below its 1981–2004 20th percentile (Shah and Steinberg 2017). Remittances are cash received in the last 12 months from parents, siblings, or children outside the household. Expenditure is in 2002 pesos per person per month. Rural is a locality under 2,500 inhabitants (INEGI 2024).

Table 8—2009 Drought and Child Anthropometrics, Binary DID Model 3 with All Covariates

	HAZ (1)	WAZ (2)	BMIZ (3)
2009 drought × post	0.021	-0.299**	-0.304***
	(0.113)	(0.125)	(0.095)
2005 drought × post	0.127	-0.193	-0.129
	(0.114)	(0.124)	(0.085)
2006 drought × post	-0.254	-0.343	-0.043
	(0.158)	(0.294)	(0.199)
2007 drought × post	-0.137	-0.181	-0.202*
	(0.118)	(0.152)	(0.120)
2008 drought × post	-0.070	0.037	0.350**
	(0.154)	(0.181)	(0.138)
2002 dwelling has electricity × post	-0.707**	-0.494**	0.016
	(0.321)	(0.211)	(0.239)
2002 good floor quality × post	-0.140	-0.031	0.137
	(0.114)	(0.156)	(0.102)
2002 good roof quality × post	0.083	-0.161	-0.155
	(0.093)	(0.160)	(0.099)
2002 enough ventilation × post	0.057	-0.010	0.140**
	(0.077)	(0.124)	(0.065)
2002 sleeping rooms per member × post	0.116	0.224	0.258*
	(0.187)	(0.255)	(0.146)
2002 ln expenditure per capita × post	0.023	0.147**	-0.044
	(0.053)	(0.064)	(0.042)
2002 rural locality × post	-0.137	-0.033	0.016
	(0.096)	(0.115)	(0.073)
Male × post	0.089	0.021	-0.000
	(0.069)	(0.103)	(0.061)
2005 mother's height (cm) × post	-0.009	-0.005	-0.001
	(0.006)	(0.008)	(0.005)
2005 mother's years of schooling × post	0.008	0.023	0.015
	(0.012)	(0.015)	(0.011)
Mother's age at birth × post	-0.002	0.005	-0.001
	(0.006)	(0.007)	(0.005)
2005 mother employed × post	-0.033	-0.171	-0.011
	(0.077)	(0.117)	(0.079)
2005 fully vaccinated × post	-0.020	0.282***	0.053
	(0.086)	(0.105)	(0.064)
Child FE	Y	Y	Y
Interview year-month FE	Y	Y	Y
Children	1,755	700	1,714
2005 municipalities (clusters)	116	101	116

Notes: This table reports every coefficient of Model 3, which is Col. 3 of the main binary DID tables, including all control variables. All regressions include child and interview year-month fixed effects, children in singleton cells are dropped, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 9—2009 Drought and Height-for-Age, Binary DID, All Three Waves

	Height-for-age z-score (HAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	0.023	0.048	0.144	0.099	0.181*
	(0.087)	(0.090)	(0.090)	(0.136)	(0.103)
Drought × post × 2002 remittances					-0.166
					(0.141)
2002 remittances × post					-0.084
					(0.092)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,293	1,293	1,084	559	1,003

Notes: This robustness check keeps only children measured in all three waves. The table reports the effect of the 2009 drought on HAZ with the binary DID for these children. They are aged 15 or younger in both survey waves. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 10—2009 Drought and Weight-for-Age, Binary DID, All Three Waves

	Weight-for-age z-score (WAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	0.025	-0.084	0.064	0.361	0.293
	(0.136)	(0.176)	(0.183)	(0.266)	(0.233)
Drought × post × 2002 remittances					-0.444
					(0.360)
2002 remittances × post					0.149
					(0.257)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	266	266	203	121	179

Notes: This robustness check keeps only children measured in all three waves. The table reports the effect of the 2009 drought on WAZ with the binary DID for these children. They are aged 10 or younger in both survey waves. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 11—2009 Drought and BMI-for-Age, Binary DID, All Three Waves

	BMI-for-age z-score (BMIZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.147	-0.223**	-0.250**	-0.171	-0.190
	(0.089)	(0.107)	(0.118)	(0.121)	(0.147)
Drought × post × 2002 remittances					-0.032
					(0.247)
2002 remittances × post					0.012
					(0.098)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,196	1,196	1,006	513	933

Notes: This robustness check keeps only children measured in all three waves. The table reports the effect of the 2009 drought on BMIZ with the binary DID for these children. They are aged 15 or younger in both survey waves. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 12—Event Study, 2009 Drought and Height-for-Age

	Height-for-age z-score (HAZ)	
	(1) Event study	(2) Triple difference
Drought × 2002 wave (lead)	0.115	0.192
	(0.116)	(0.128)
Drought × 2009 wave	0.015	-0.008
	(0.110)	(0.114)
Drought × 2002 remittances × 2002 wave (lead)		-0.297
		(0.180)
Drought × 2002 remittances × 2009 wave		-0.158
		(0.147)
2005–2008 droughts × wave	Y	Y
2002 household controls × wave	Y	Y
2005 child and mother controls × wave	Y	Y
Child FE	Y	Y
Interview year-month FE	Y	Y
Children	1,755	1,622

Notes: This table tests for pre-trends before the 2009 drought. Regressions are at the child-wave level with 2005 as the reference wave. The sample is children aged 15 or younger in each wave they are in. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Col. 1 keeps the children in Model 3 of the binary DID and Col. 2 those in Model 5, so each matches its DID sample. Col. 1 interacts the drought indicator with the 2002 and 2009 waves. The 2002 interaction is the lead. Col. 2 adds the drought terms times 2002 remittances. The 2002 remittance terms on their own are not shown. Baseline controls are interacted with wave to allow different trends by pre-treatment characteristics, following the same logic as the controls interacted with post in the binary DID. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 13—Event Study, 2009 Drought and Weight-for-Age

	Weight-for-age z-score (WAZ)	
	(1) Event study	(2) Triple difference
Drought × 2002 wave (lead)	-0.481**	-0.336
	(0.216)	(0.244)
Drought × 2009 wave	-0.298**	-0.105
	(0.126)	(0.145)
Drought × 2002 remittances × 2002 wave (lead)		-0.078
		(0.347)
Drought × 2002 remittances × 2009 wave		-0.373*
		(0.208)
2005–2008 droughts × wave	Y	Y
2002 household controls × wave	Y	Y
2005 child and mother controls × wave	Y	Y
Child FE	Y	Y
Interview year-month FE	Y	Y
Children	700	634

Notes: This table tests for pre-trends before the 2009 drought. Regressions are at the child-wave level with 2005 as the reference wave. The sample is children aged 10 or younger in each wave they are in. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Col. 1 keeps the children in Model 3 of the binary DID and Col. 2 those in Model 5, so each matches its DID sample. Col. 1 interacts the drought indicator with the 2002 and 2009 waves. The 2002 interaction is the lead. Col. 2 adds the drought terms times 2002 remittances. The 2002 remittance terms on their own are not shown. Baseline controls are interacted with wave to allow different trends by pre-treatment characteristics, following the same logic as the controls interacted with post in the binary DID. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 14—Event Study, 2009 Drought and BMI-for-Age

	BMI-for-age z-score (BMIZ)	
	(1) Event study	(2) Triple difference
Drought × 2002 wave (lead)	-0.387***	-0.423***
	(0.104)	(0.120)
Drought × 2009 wave	-0.293***	-0.262**
	(0.097)	(0.114)
Drought × 2002 remittances × 2002 wave (lead)		0.186
		(0.202)
Drought × 2002 remittances × 2009 wave		0.040
		(0.217)
2005–2008 droughts × wave	Y	Y
2002 household controls × wave	Y	Y
2005 child and mother controls × wave	Y	Y
Child FE	Y	Y
Interview year-month FE	Y	Y
Children	1,714	1,587

Notes: This table tests for pre-trends before the 2009 drought. Regressions are at the child-wave level with 2005 as the reference wave. The sample is children aged 15 or younger in each wave they are in. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Col. 1 keeps the children in Model 3 of the binary DID and Col. 2 those in Model 5, so each matches its DID sample. Col. 1 interacts the drought indicator with the 2002 and 2009 waves. The 2002 interaction is the lead. Col. 2 adds the drought terms times 2002 remittances. The 2002 remittance terms on their own are not shown. Baseline controls are interacted with wave to allow different trends by pre-treatment characteristics, following the same logic as the controls interacted with post in the binary DID. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Figure 7—2009 Drought by Municipality, Binary DID Treatment






Figure 8—Event Study, BMI-for-Age, Binary DID, With Controls




Notes: Drought compared with no drought. The coefficients are on the 2009 drought indicator interacted with the 2002 and 2009 survey waves. The sample is the children in Model 3 of the binary difference-in-differences, in each survey wave in which they are measured. The Model 3 controls are each interacted with the 2002 and 2009 survey waves. Included are child and interview year-month fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 9—Event Study, Height-for-Age, Binary DID, With Controls




Notes: Drought compared with no drought. The coefficients are on the 2009 drought indicator interacted with the 2002 and 2009 survey waves. The sample is the children in Model 3 of the binary difference-in-differences, in each survey wave in which they are measured. The Model 3 controls are each interacted with the 2002 and 2009 survey waves. Included are child and interview year-month fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 10—Event Study, Weight-for-Age, Binary DID, With Controls




Notes: Drought compared with no drought. The coefficients are on the 2009 drought indicator interacted with the 2002 and 2009 survey waves. The sample is the children in Model 3 of the binary difference-in-differences, in each survey wave in which they are measured. The Model 3 controls are each interacted with the 2002 and 2009 survey waves. Included are child and interview year-month fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 11—Event Study by Dose Level, Height-for-Age, Discrete Dose, Children in All Three Waves




Notes: Moderately dry and severely dry or worse compared with near-normal. The sample is the children in the discrete-dose table who were measured in all three survey waves, so the 2002 lead and the 2009 coefficient come from the same children. Similar to the discrete-dose table, there are no controls. Included are child and survey wave fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 12—Event Study by Dose Level, Weight-for-Age, Discrete Dose, Children in All Three Waves




Notes: Moderately dry and severely dry or worse compared with near-normal. The sample is the children in the discrete-dose table who were measured in all three survey waves, so the 2002 lead and the 2009 coefficient come from the same children. Similar to the discrete-dose table, there are no controls. Included are child and survey wave fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 13—Event Study for Pooled Dry, BMI-for-Age, Discrete Dose




Notes: Dry ($z \leq -1$) compared with near-normal, pooling the moderately dry and severely dry or worse dose groups. The figure looks at pre-trends for the pooled dry estimates in the discrete-dose table. Similar to the discrete-dose table, there are no controls. Included are child and survey wave fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 14—Event Study for Pooled Dry, Height-for-Age, Discrete Dose



Notes: Dry ($z \leq -1$) compared with near-normal, pooling the moderately dry and severely dry or worse dose groups. The figure looks at pre-trends for the pooled dry estimates in the discrete-dose table. Similar to the discrete-dose table, there are no controls. Included are child and survey wave fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Figure 15—Event Study for Pooled Dry, Weight-for-Age, Discrete Dose



Notes: Dry ($z \leq -1$) compared with near-normal, pooling the moderately dry and severely dry or worse dose groups. The figure looks at pre-trends for the pooled dry estimates in the discrete-dose table. Similar to the discrete-dose table, there are no controls. Included are child and survey wave fixed effects. 2005 is the reference wave. 95% confidence intervals use standard errors clustered by 2005 municipality.

Online Appendix

Table 15—Sample Sizes, Treated and Control Children

	Panel A: Binary DID, 2009 drought	
	Control	Treated
HAZ sample, children aged 15 or younger in 2005 and 2009		
Children, estimation sample	1,514	922
Children, Model 1 (basic DID)	1,512	920
Children, Model 3 (DID with controls)	1,077	678
Children, Model 5 (triple difference)	987	635
In 2002 remittance households	390	193
Households, estimation sample	944	578
2005 municipalities, estimation sample	87	46
WAZ sample, children aged 10 or younger in 2005 and 2009		
Children, estimation sample	696	388
Children, Model 1 (basic DID)	692	386
Children, Model 3 (DID with controls)	445	255
Children, Model 5 (triple difference)	400	234
In 2002 remittance households	168	90
Households, estimation sample	565	316
2005 municipalities, estimation sample	79	44
BMIZ sample, children aged 15 or younger in 2005 and 2009		
Children, estimation sample	1,487	896
Children, Model 1 (basic DID)	1,485	894
Children, Model 3 (DID with controls)	1,057	657
Children, Model 5 (triple difference)	968	619
In 2002 remittance households	381	189
Households, estimation sample	931	561
2005 municipalities, estimation sample	87	46

	Panel B: Continuous DID, any 2009 deficit	
	Control	Treated
HAZ sample, children aged 15 or younger in 2005 and 2009		
Children, continuous DID sample	829	1,607
In 2002 remittance households	236	438
In 2002 non-remittance households	437	738
Households	501	1,021
2005 municipalities	46	87
WAZ sample, children aged 10 or younger in 2005 and 2009		
Children, continuous DID sample	380	704
In 2002 remittance households	100	193
In 2002 non-remittance households	172	260
Households	304	577
2005 municipalities	42	81
BMIZ sample, children aged 15 or younger in 2005 and 2009		
Children, continuous DID sample	815	1,568
In 2002 remittance households	231	427
In 2002 non-remittance households	430	720
Households	496	996
2005 municipalities	46	87

	Panel C: Dose groups vs near normal	
	Control	Treated
Moderately dry ($-1.5 < z \leq -1$)		
Children, HAZ sample	1,493	488
Children, WAZ sample	683	217
Children, BMIZ sample	1,464	472
2005 municipalities, HAZ sample	89	23
2005 municipalities, WAZ sample	80	23
2005 municipalities, BMIZ sample	89	23
Severely dry or worse ($z \leq -1.5$)		
Children, HAZ sample	1,493	370
Children, WAZ sample	683	145
Children, BMIZ sample	1,464	364
2005 municipalities, HAZ sample	89	17
2005 municipalities, WAZ sample	80	16
2005 municipalities, BMIZ sample	89	17
Severely dry ($-2 < z \leq -1.5$)		
Children, HAZ sample	1,493	302
Children, WAZ sample	683	121
Children, BMIZ sample	1,464	296
2005 municipalities, HAZ sample	89	15
2005 municipalities, WAZ sample	80	14
2005 municipalities, BMIZ sample	89	15
Extremely dry ($z \leq -2$)		
Children, HAZ sample	1,493	68
Children, WAZ sample	683	24
Children, BMIZ sample	1,464	68
2005 municipalities, HAZ sample	89	2
2005 municipalities, WAZ sample	80	2
2005 municipalities, BMIZ sample	89	2

Notes: Counts are children, households or 2005 municipalities in each regression sample at the 2005 baseline. Estimation samples are children with the outcome measured in both MxFLS-2 (2005) and MxFLS-3 (2009) who have a 2005 municipality. They are aged 15 or younger for HAZ and BMIZ and 10 or younger for WAZ in both survey waves. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. In Panel A, treated children's 2005 municipality had a drought in 2009, meaning May–October CHIRPS rainfall in the maize growing season was below its 1981–2004 20th percentile (Jessoe, Manning, and Taylor 2018; Shah and Steinberg 2017). Model 1 drops children alone in their interview-month cell. Continuous DID uses the full sample. Model 3 adds 2005–2008 droughts and 2002 household and 2005 child and mother controls. Model 5 adds 2002 remittances times post and drought times post times 2002 remittances. Model 5 children not in remittance households are the Model 5 count minus the remittance row. In Panel B, treated children's 2005 municipality had 2009 rainfall below its 1981–2004 mean (z below 0), where z is 2009 rainfall in standard deviations from that mean. Panel B gives sample sizes for the continuous DID. Its two remittance rows are children in 2002 remittance and non-remittance households, the two groups compared in Panel B of Table **??**. Rows do not add up to the total because children with missing 2002 remittances are in neither row. Panel C compares each dose group with near normal (z between -1 and 1) in the discrete-dose analysis. Dose levels use standardized rainfall cutoffs of the SPI adopted by the World Meteorological Organization (McKee, Doesken, and Kleist 1993; WMO 2012). Children in wet municipalities (z of 1 or above) are not in the discrete-dose analysis. Remittances are cash received in the last 12 months from parents, siblings or children living outside the household, from MxFLS-1 (2002).

Table 16—Sample Attrition Across Waves

	Number of children (N)		
	MxFLS-1 (2002)	MxFLS-2 (2005)	MxFLS-3 (2009)
HAZ sample, children under 16			
DID (2005 and 2009)		3,582	2,436
Event studies (all three waves)	2,723	1,856	1,300
WAZ sample, children under 11			
DID (2005 and 2009)		1,895	1,084
Event studies (all three waves)	962	609	275
BMIZ sample, children under 16			
DID (2005 and 2009)		3,511	2,383
Event studies (all three waves)	2,569	1,719	1,203

Notes: Counts of children in each wave. The binary DID only uses MxFLS-2 (2005) and MxFLS-3 (2009). The continuous DID uses the same sample as the binary DID. The event studies use all three waves. Each row starts from the children who meet the regression sample rules in its first wave and could stay in the sample. They are under the age cutoff in every wave, 15 or younger for HAZ and BMIZ and 10 or younger for WAZ. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Later columns count the children still in the sample, so the drop across columns is attrition. This is not the same as the children the regressions drop as singletons, with only one wave in the sample, as these also include children who aged out or are only in a later wave. The DID 2009 count is the estimation sample, and the event study 2009 count is the children in all three waves. For children not measured in a later wave, I estimate their age in that wave to see whether they aged out. Their age comes from their birth date and their household's MxFLS-3 interview date, since MxFLS-3 interviews ran over several years. Children not in MxFLS-3 or whose birth date does not match their reported age get their last measured age plus the median gap between waves. This gap is 4 years from 2005 to 2009 and 8 years from 2002 to 2009. Among children measured in both waves, 92 and 86 percent of gaps are within a year of the median. A one-year error only matters for children near the age cutoff. This way, children who simply aged out are not counted as attrition.

Table 17—Effects of the 2009 Rainfall Deficit on Attrition, Discrete Dose

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Moderately dry ($-1.5 < z \leq -1$)	-0.023	-0.024	-0.011
	(0.044)	(0.045)	(0.044)
Severely dry or worse ($z \leq -1.5$)	-0.029	0.045	-0.035
	(0.038)	(0.043)	(0.038)
Mean attrition	0.319	0.426	0.320
Children	3,451	1,821	3,384

Notes: Each column regresses an indicator for attrition on the treatment. Each coefficient is the difference in the probability of attrition between children in that rainfall group and children in near-normal municipalities. Attrition equals 1 if the child is not in the 2009 estimation sample. The table compares moderately dry and severely dry or worse with near-normal ($-1 < z < 1$) municipalities. The sample omits children in wet municipalities (z of 1 or above). Specifically, children in the sample meet the regression sample rules in MxFLS-2 (2005). That is, they are aged 15 or younger for HAZ and BMIZ and 10 or younger for WAZ in both survey waves, have a 2005 municipality, and were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Children who aged out are not counted as attrition. There are no fixed effects or controls. Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are not reported. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 18—Effects of the 2009 Rainfall Deficit on Attrition, Binary DID

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Drought	-0.034	0.004	-0.027
	(0.032)	(0.034)	(0.032)
Mean attrition	0.320	0.428	0.321
Children	3,582	1,895	3,511

Notes: See Table **??** notes. The coefficient is the difference in the probability of attrition between children in drought and non-drought municipalities. This specification includes wet municipalities (z of 1 or above) and uses the binary drought instead of the dose groups. Drought is May–October 2009 CHIRPS rainfall below the municipality's 1981–2004 20th percentile (Shah and Steinberg 2017). $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 19—Baseline Outcomes of Attrited and Non-Attrited Children, Discrete Dose

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Attrited	0.074	0.167**	0.082
	(0.069)	(0.071)	(0.062)
Mean of 2005 outcome	-0.557	-0.227	0.381
Children	3,451	1,821	3,384

Notes: This table tests whether the children lost to attrition differ at baseline from those who stay. Each column regresses the 2005 WHO z-score on an indicator for attrition, for all children. Attrition equals 1 if the child is not in the 2009 estimation sample. The sample is as in Table **??**. The treatment does not enter. The dose and binary versions differ only in that the discrete-dose version excludes children in wet municipalities (z of 1 or above). There are no fixed effects or controls. Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are not reported. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 20—Baseline Outcomes of Attrited and Non-Attrited Children, Binary DID

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Attrited	0.101	0.188***	0.091
	(0.071)	(0.071)	(0.060)
Mean of 2005 outcome	-0.565	-0.236	0.385
Children	3,582	1,895	3,511

Notes: See Table **??** notes. This version includes children in wet municipalities. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 21—Baseline Outcomes of Children Lost to Attrition, Discrete Dose

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Moderately dry ($-1.5 < z \leq -1$)	0.061	-0.196	0.083
	(0.162)	(0.160)	(0.159)
Severely dry or worse ($z \leq -1.5$)	0.290*	-0.046	0.073
	(0.152)	(0.131)	(0.134)
Mean of 2005 outcome	-0.507	-0.131	0.436
Children	1,100	776	1,084

Notes: This table tests whether the children lost to attrition from moderately dry and severely dry or worse municipalities differ at baseline from those lost from near-normal ($-1 < z < 1$) municipalities. It keeps only the children lost to attrition, without children in wet municipalities (z of 1 or above). Each column regresses their 2005 WHO z-score on the treatment. The sample is as in Table **??**. There are no fixed effects or controls. Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are not reported. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 22—Baseline Outcomes of Children Lost to Attrition, Binary DID

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Drought	0.140	-0.104	0.064
	(0.122)	(0.113)	(0.112)
Mean of 2005 outcome	-0.496	-0.128	0.446
Children	1,146	811	1,128

Notes: See Table **??** notes. This version compares the children lost to attrition from drought municipalities with those lost from all other municipalities, including wet ones. Drought is May–October 2009 CHIRPS rainfall below the municipality's 1981–2004 20th percentile (Shah and Steinberg 2017). $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 23—2009 Drought and Height-for-Age, Binary DID, Boys

	Height-for-age z-score (HAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.123	-0.108	-0.108	0.228	-0.075
	(0.123)	(0.130)	(0.146)	(0.237)	(0.142)
Drought × post × 2002 remittances					-0.301
					(0.257)
2002 remittances × post					-0.254**
					(0.123)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,201	1,201	869	353	806

Notes: Estimating the effect separately for boys and girls tests for gender disparities in the impact of the drought, as households facing an income shock may protect the nutrition and health of sons and daughters differently. The table reports the effect of the 2009 drought on the WHO height-for-age z-score (HAZ) with the binary DID for boys, using sex reported in the 2005 household roster, aged 15 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include child sex. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 24—2009 Drought and Weight-for-Age, Binary DID, Boys

	Weight-for-age z-score (WAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.219	-0.175	-0.264	-0.330	-0.064
	(0.153)	(0.149)	(0.182)	(0.358)	(0.196)
Drought × post × 2002 remittances					-0.225
					(0.283)
2002 remittances × post					0.084
					(0.179)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	551	551	367	103	331

Notes: Estimating the effect separately for boys and girls tests for gender disparities in the impact of the drought, as households facing an income shock may protect the nutrition and health of sons and daughters differently. The table reports the effect of the 2009 drought on the WHO weight-for-age z-score (WAZ) with the binary DID for boys, using sex reported in the 2005 household roster, aged 10 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include child sex. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 25—2009 Drought and BMI-for-Age, Binary DID, Boys

	BMI-for-age z-score (BMIZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.170*	-0.168	-0.269**	-0.262	-0.336*
	(0.099)	(0.119)	(0.131)	(0.180)	(0.175)
Drought × post × 2002 remittances					0.416
					(0.322)
2002 remittances × post					-0.057
					(0.111)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,170	1,170	847	346	787

Notes: Estimating the effect separately for boys and girls tests for gender disparities in the impact of the drought, as households facing an income shock may protect the nutrition and health of sons and daughters differently. The table reports the effect of the 2009 drought on the WHO BMI-for-age z-score (BMIZ) with the binary DID for boys, using sex reported in the 2005 household roster, aged 15 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include child sex. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 26—2009 Drought and Height-for-Age, Binary DID, Girls

	Height-for-age z-score (HAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.046	0.020	0.111	-0.055	0.035
	(0.101)	(0.116)	(0.131)	(0.202)	(0.139)
Drought × post × 2002 remittances					-0.060
					(0.176)
2002 remittances × post					-0.143
					(0.115)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,220	1,220	878	377	811

Notes: Estimating the effect separately for boys and girls tests for gender disparities in the impact of the drought, as households facing an income shock may protect the nutrition and health of sons and daughters differently. The table reports the effect of the 2009 drought on the WHO height-for-age z-score (HAZ) with the binary DID for girls, using sex reported in the 2005 household roster, aged 15 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include child sex. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 27—2009 Drought and Weight-for-Age, Binary DID, Girls

	Weight-for-age z-score (WAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.259**	-0.282**	-0.379**	-0.184	-0.265
	(0.111)	(0.119)	(0.167)	(0.288)	(0.197)
Drought × post × 2002 remittances					-0.401
					(0.302)
2002 remittances × post					0.097
					(0.189)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	520	520	324	115	295

Notes: Estimating the effect separately for boys and girls tests for gender disparities in the impact of the drought, as households facing an income shock may protect the nutrition and health of sons and daughters differently. The table reports the effect of the 2009 drought on the WHO weight-for-age z-score (WAZ) with the binary DID for girls, using sex reported in the 2005 household roster, aged 10 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include child sex. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 28—2009 Drought and BMI-for-Age, Binary DID, Girls

	BMI-for-age z-score (BMIZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.112	-0.160	-0.313***	-0.200	-0.184
	(0.077)	(0.097)	(0.116)	(0.167)	(0.114)
Drought × post × 2002 remittances					-0.342*
					(0.200)
2002 remittances × post					0.065
					(0.098)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,199	1,199	861	364	795

Notes: Estimating the effect separately for boys and girls tests for gender disparities in the impact of the drought, as households facing an income shock may protect the nutrition and health of sons and daughters differently. The table reports the effect of the 2009 drought on the WHO BMI-for-age z-score (BMIZ) with the binary DID for girls, using sex reported in the 2005 household roster, aged 15 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include child sex. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 29—2009 Drought and Height-for-Age, Binary DID, Rural Children

	Height-for-age z-score (HAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.045	0.189	0.190	0.472	0.273*
	(0.162)	(0.173)	(0.163)	(0.289)	(0.159)
Drought × post × 2002 remittances					-0.278
					(0.288)
2002 remittances × post					-0.006
					(0.130)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,016	1,016	732	272	714

Notes: Estimating the effect separately for rural and urban children tests whether the drought hits harder where household incomes depend more directly on rain-fed agriculture, while urban households are mainly exposed through food prices. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). The table reports the effect of the 2009 drought on the WHO height-for-age z-score (HAZ) with the binary DID for rural children (2005 household locality), aged 15 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include rural. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 30—2009 Drought and Weight-for-Age, Binary DID, Rural Children

	Weight-for-age z-score (WAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.178	-0.074	-0.276	0.532	-0.131
	(0.180)	(0.172)	(0.264)	(0.334)	(0.281)
Drought × post × 2002 remittances					-0.362
					(0.332)
2002 remittances × post					0.216
					(0.225)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	414	414	265	85	260

Notes: Estimating the effect separately for rural and urban children tests whether the drought hits harder where household incomes depend more directly on rain-fed agriculture, while urban households are mainly exposed through food prices. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). The table reports the effect of the 2009 drought on the WHO weight-for-age z-score (WAZ) with the binary DID for rural children (2005 household locality), aged 10 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include rural. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 31—2009 Drought and BMI-for-Age, Binary DID, Rural Children

	BMI-for-age z-score (BMIZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.096	-0.138	-0.208	-0.244	-0.197
	(0.112)	(0.127)	(0.180)	(0.254)	(0.204)
Drought × post × 2002 remittances					-0.035
					(0.255)
2002 remittances × post					0.086
					(0.095)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	998	998	719	264	701

Notes: Estimating the effect separately for rural and urban children tests whether the drought hits harder where household incomes depend more directly on rain-fed agriculture, while urban households are mainly exposed through food prices. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). The table reports the effect of the 2009 drought on the WHO BMI-for-age z-score (BMIZ) with the binary DID for rural children (2005 household locality), aged 15 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include rural. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 32—2009 Drought and Height-for-Age, Binary DID, Urban Children

	Height-for-age z-score (HAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.102	-0.136	-0.088	-0.021	-0.243*
	(0.106)	(0.110)	(0.125)	(0.200)	(0.122)
Drought × post × 2002 remittances					0.032
					(0.167)
2002 remittances × post					-0.331***
					(0.121)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,412	1,412	1,020	455	904

Notes: Estimating the effect separately for rural and urban children tests whether the drought hits harder where household incomes depend more directly on rain-fed agriculture, while urban households are mainly exposed through food prices. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). The table reports the effect of the 2009 drought on the WHO height-for-age z-score (HAZ) with the binary DID for urban children (2005 household locality), aged 15 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include rural. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 33—2009 Drought and Weight-for-Age, Binary DID, Urban Children

	Weight-for-age z-score (WAZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.282**	-0.370***	-0.446***	-0.213	-0.201
	(0.114)	(0.108)	(0.135)	(0.254)	(0.175)
Drought × post × 2002 remittances					-0.486*
					(0.261)
2002 remittances × post					0.035
					(0.145)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	654	654	427	137	365

Notes: Estimating the effect separately for rural and urban children tests whether the drought hits harder where household incomes depend more directly on rain-fed agriculture, while urban households are mainly exposed through food prices. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). The table reports the effect of the 2009 drought on the WHO weight-for-age z-score (WAZ) with the binary DID for urban children (2005 household locality), aged 10 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include rural. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 34—2009 Drought and BMI-for-Age, Binary DID, Urban Children

	BMI-for-age z-score (BMIZ)				
	(1)	(2)	(3)	(4)	(5)
Drought × post	-0.189**	-0.259***	-0.382***	-0.192	-0.316**
	(0.078)	(0.080)	(0.096)	(0.141)	(0.128)
Drought × post × 2002 remittances					0.043
					(0.327)
2002 remittances × post					0.049
					(0.107)
2005–2008 droughts × post	N	Y	Y	Y	Y
2002 household controls × post	N	N	Y	Y	Y
2005 child and mother controls × post	N	N	Y	Y	Y
2005 father and birth controls × post	N	N	N	Y	N
Child FE	Y	Y	Y	Y	Y
Interview year-month FE	Y	Y	Y	Y	Y
Children	1,377	1,377	992	441	882

Notes: Estimating the effect separately for rural and urban children tests whether the drought hits harder where household incomes depend more directly on rain-fed agriculture, while urban households are mainly exposed through food prices. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). The table reports the effect of the 2009 drought on the WHO BMI-for-age z-score (BMIZ) with the binary DID for urban children (2005 household locality), aged 15 or younger in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Columns, variables and controls are as in the main binary DID tables. The controls do not include rural. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 35—Continuous DID and Remittances as a Moderator

	HAZ (1)	WAZ (2)	BMIZ (3)
Panel A: Continuous DID, any 2009 rainfall deficit			
Average ATT	-0.077	0.028	-0.045
	(0.101)	(0.097)	(0.069)
	[0.462]	[0.777]	[0.518]
Average ACRT	-0.007	-0.033	0.029
	(0.137)	(0.173)	(0.130)
Children	2,436	1,084	2,383
Panel B: Average ATT by 2002 remittances			
No 2002 remittances	-0.080	0.050	-0.042
	(0.107)	(0.152)	(0.091)
2002 remittances	-0.279	-0.030	0.031
	(0.182)	(0.190)	(0.116)
Difference, remittances minus none	-0.199	-0.081	0.074
	(0.179)	(0.218)	(0.145)
Children	1,849	725	1,808
Panel C: Dry vs near normal by 2002 remittances			
Dry ($z \leq -1$)	-0.054	-0.160	-0.251**
	(0.114)	(0.200)	(0.109)
	[0.657]	[0.543]	[0.027]
Dry × 2002 remittances	-0.135	-0.220	0.100
	(0.133)	(0.216)	(0.207)
	[0.317]	[0.329]	[0.653]
2002 remittances	-0.040	0.088	0.039
	(0.109)	(0.134)	(0.070)
	[0.722]	[0.524]	[0.577]
Children	1,773	693	1,734

Notes: Coefficients are effects on the 2005 to 2009 change in the WHO z-score for the binary DID estimation samples. The dose is the 2009 rainfall deficit, max(0, -z), where z is 2009 May–October CHIRPS rainfall in the child's 2005 municipality in standard deviations from its 1981–2004 mean (Jessoe, Manning, and Taylor 2018). Panel A compares children with any 2009 deficit (z ¡ 0) with children without one using the contdid package (Callaway, Goodman-Bacon, and Sant'Anna 2026). Panel B gives the average ATT for children in 2002 remittance and non-remittance households and their difference. Panel C interacts dry ($z \leq -1$) versus near-normal rainfall with 2002 remittances using the discrete-dose regression in equation 4.1 (Callaway, Goodman-Bacon, and Sant'Anna 2024). Remittances are cash received in the last 12 months from parents, siblings, or children outside the household in MxFLS-1 (2002). The average ATT requires parallel trends and the average ACRT requires strong parallel trends (Callaway, Goodman-Bacon, and Sant'Anna 2024). No regression includes controls or fixed effects. Standard errors clustered by 2005 municipality are in parentheses. For the average ACRT they come from a multiplier bootstrap of the contdid influence functions. Wild cluster bootstrap p-values are in brackets (MacKinnon, Nielsen, and Webb 2023). Stars use the clustered p-values, and where a wild bootstrap p-value is shown both must pass. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 36—Continuous DID by Subgroup, Height-for-Age

	Change in the height-for-age z-score (HAZ)			
	Boys (1)	Girls (2)	Rural (3)	Urban (4)
Any 2009 deficit	-0.050	-0.108	0.101	-0.180
	(0.114)	(0.116)	(0.181)	(0.101)
	[0.672]	[0.376]	[0.614]	[0.104]
Dry vs near normal ($z \leq -1$)	-0.073	-0.009	0.033	-0.095
	(0.114)	(0.115)	(0.180)	(0.106)
	[0.533]	[0.937]	[0.876]	[0.406]
Severely dry or worse vs near normal ($z \leq -1.5$)	-0.158	-0.077	0.181	-0.241*
	(0.158)	(0.106)	(0.191)	(0.115)
	[0.344]	[0.488]	[0.519]	[0.065]
Dry × 2002 remittances	-0.364	0.089	-0.197	-0.098
	(0.226)	(0.168)	(0.211)	(0.175)
	[0.121]	[0.610]	[0.375]	[0.586]
Children	1,208	1,228	1,020	1,416
Municipalities with no 2009 deficit	42	41	27	24

Notes: This table repeats the continuous DID and discrete-dose results for the WHO height-for-age z-score (HAZ) by subgroup. Cols. 1–2 test for gender disparities in the impact of the drought, using sex reported in the 2005 household roster, and Cols. 3–4 test whether it hits rural children harder, as rural household incomes depend more directly on rain-fed agriculture. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). Rows, samples and definitions are as in the main continuous DID and discrete-dose tables. Children are aged 15 or younger in both survey waves. The any deficit row uses the contdid package, and the dose-group rows use the discrete-dose regression in equation 4.1 (Callaway, Goodman-Bacon, and Sant'Anna 2024). Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are in brackets, which tend to be conservative with few treated municipalities (MacKinnon, Nielsen, and Webb 2023). Stars use the clustered p-values, and where a wild bootstrap p-value is shown both must pass. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 37—Continuous DID by Subgroup, Weight-for-Age

	Change in the weight-for-age z-score (WAZ)			
	Boys (1)	Girls (2)	Rural (3)	Urban (4)
Any 2009 deficit	0.139	-0.089	0.066	0.047
	(0.143)	(0.112)	(0.202)	(0.094)
	[0.348]	[0.436]	[0.756]	[0.622]
Dry vs near normal ($z \leq -1$)	-0.221	-0.179	-0.143	-0.218**
	(0.180)	(0.111)	(0.227)	(0.103)
	[0.262]	[0.122]	[0.639]	[0.044]
Severely dry or worse vs near normal ($z \leq -1.5$)	-0.152	-0.153	0.013	-0.237
	(0.198)	(0.170)	(0.124)	(0.154)
	[0.471]	[0.428]	[0.924]	[0.168]
Dry × 2002 remittances	-0.008	-0.456	-0.144	-0.327
	(0.301)	(0.274)	(0.348)	(0.250)
	[0.977]	[0.102]	[0.705]	[0.209]
Children	556	528	424	660
Municipalities with no 2009 deficit	36	32	23	23

Notes: This table repeats the continuous DID and discrete-dose results for the WHO weight-for-age z-score (WAZ) by subgroup. Cols. 1–2 test for gender disparities in the impact of the drought, using sex reported in the 2005 household roster, and Cols. 3–4 test whether it hits rural children harder, as rural household incomes depend more directly on rain-fed agriculture. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). Rows, samples and definitions are as in the main continuous DID and discrete-dose tables. Children are aged 10 or younger in both survey waves. The any deficit row uses the contdid package, and the dose-group rows use the discrete-dose regression in equation 4.1 (Callaway, Goodman-Bacon, and Sant'Anna 2024). Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are in brackets, which tend to be conservative with few treated municipalities (MacKinnon, Nielsen, and Webb 2023). Stars use the clustered p-values, and where a wild bootstrap p-value is shown both must pass. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 38—Continuous DID by Subgroup, BMI-for-Age

	Change in the BMI-for-age z-score (BMIZ)			
	Boys (1)	Girls (2)	Rural (3)	Urban (4)
Any 2009 deficit	-0.008	-0.082	-0.035	-0.054
	(0.093)	(0.076)	(0.090)	(0.095)
	[0.933]	[0.294]	[0.699]	[0.593]
Dry vs near normal ($z \leq -1$)	-0.202*	-0.124	-0.147	-0.174
	(0.101)	(0.080)	(0.094)	(0.111)
	[0.054]	[0.138]	[0.155]	[0.141]
Severely dry or worse vs near normal ($z \leq -1.5$)	-0.064	-0.106	-0.129	-0.077
	(0.105)	(0.102)	(0.090)	(0.085)
	[0.548]	[0.350]	[0.309]	[0.379]
Dry × 2002 remittances	0.506	-0.301	-0.030	0.216
	(0.324)	(0.180)	(0.263)	(0.304)
	[0.142]	[0.119]	[0.915]	[0.534]
Children	1,177	1,206	1,002	1,381
Municipalities with no 2009 deficit	42	41	27	24

Notes: This table repeats the continuous DID and discrete-dose results for the WHO BMI-for-age z-score (BMIZ) by subgroup. Cols. 1–2 test for gender disparities in the impact of the drought, using sex reported in the 2005 household roster, and Cols. 3–4 test whether it hits rural children harder, as rural household incomes depend more directly on rain-fed agriculture. Rural is a locality with fewer than 2,500 inhabitants (INEGI 2024). Rows, samples and definitions are as in the main continuous DID and discrete-dose tables. Children are aged 15 or younger in both survey waves. The any deficit row uses the contdid package, and the dose-group rows use the discrete-dose regression in equation 4.1 (Callaway, Goodman-Bacon, and Sant'Anna 2024). Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are in brackets, which tend to be conservative with few treated municipalities (MacKinnon, Nielsen, and Webb 2023). Stars use the clustered p-values, and where a wild bootstrap p-value is shown both must pass. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 39—Triple Difference by Sex, Binary DID

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Drought × post	0.104	-0.241	-0.306***
	(0.135)	(0.175)	(0.110)
Drought × post × male	-0.163	-0.107	0.004
	(0.146)	(0.236)	(0.130)
Male × post	0.152*	0.059	-0.002
	(0.081)	(0.123)	(0.076)
P-val: pre-trend 2002–2005	0.724	0.114	0.712
Children	1,755	700	1,714

Notes: This table tests whether the effect of the 2009 drought differs for boys with a triple difference on all children, using the binary DID. Male equals 1 for boys, using sex reported in the 2005 household roster. Children are aged 15 or younger for HAZ and BMIZ and 10 or younger for WAZ in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. The regressions keep the controls of Col. 3 of the main binary DID tables and add drought × post × male, the difference in the drought effect for boys. The 2005 child and mother controls do not include child sex, which enters as male × post. The pre-trend p-value tests the 2002 lead of the triple difference in the event study on the same children, with the controls interacted with the 2002 and 2009 waves. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 40—Triple Difference by Sex, Discrete Dose

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Dry ($z \leq -1$)	-0.009	-0.179	-0.124
	(0.115)	(0.111)	(0.080)
	[0.942]	[0.128]	[0.138]
Dry × male	-0.065	-0.041	-0.078
	(0.112)	(0.194)	(0.102)
	[0.576]	[0.834]	[0.451]
Male	0.099	0.109	0.065
	(0.064)	(0.119)	(0.064)
	[0.131]	[0.370]	[0.316]
Children	2,351	1,045	2,300

Notes: This table tests whether the effect of the 2009 drought differs for boys with a triple difference on all children, using the discrete dose. Male equals 1 for boys, using sex reported in the 2005 household roster. Children are aged 15 or younger for HAZ and BMIZ and 10 or younger for WAZ in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Regressions are on the 2005 to 2009 change in the WHO z-score, without children in wet municipalities (z of 1 or above). The omitted group is near-normal rainfall ($-1 < z < 1$). There are no controls or fixed effects. Dry × male is the difference in the effect of dry rainfall for boys. Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are in brackets, which tend to be conservative with few treated municipalities (MacKinnon, Nielsen, and Webb 2023). Stars use the clustered p-values, and where a wild bootstrap p-value is shown both must pass. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 41—Triple Difference by Rural Residence, Binary DID

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Drought × post	0.011	-0.334***	-0.296**
	(0.127)	(0.123)	(0.117)
Drought × post × rural	0.043	0.112	-0.028
	(0.192)	(0.240)	(0.153)
Rural × post	-0.100	-0.046	0.056
	(0.103)	(0.159)	(0.096)
P-val: pre-trend 2002–2005	0.700	0.286	0.572
Children	1,755	700	1,714

Notes: This table tests whether the effect of the 2009 drought differs for rural children with a triple difference on all children, using the binary DID. Rural equals 1 if the child's 2005 household locality had fewer than 2,500 inhabitants (INEGI 2024). Children are aged 15 or younger for HAZ and BMIZ and 10 or younger for WAZ in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. The regressions keep the controls of Col. 3 of the main binary DID tables and add drought × post × rural, the difference in the drought effect for rural children. The 2002 household controls do not include rural, which enters as 2005 rural × post. The pre-trend p-value tests the 2002 lead of the triple difference in the event study on the same children, with the controls interacted with the 2002 and 2009 waves. All regressions include child and interview year-month fixed effects, and standard errors are clustered by 2005 municipality. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 42—Triple Difference by Rural Residence, Discrete Dose

	HAZ	WAZ	BMIZ
	(1)	(2)	(3)
Dry ($z \leq -1$)	-0.095	-0.218**	-0.174
	(0.106)	(0.103)	(0.110)
	[0.393]	[0.049]	[0.141]
Dry × rural	0.128	0.075	0.027
	(0.206)	(0.248)	(0.143)
	[0.575]	[0.796]	[0.865]
Rural	-0.141	-0.213*	-0.014
	(0.110)	(0.114)	(0.075)
	[0.220]	[0.073]	[0.847]
Children	2,351	1,045	2,300

Notes: This table tests whether the effect of the 2009 drought differs for rural children with a triple difference on all children, using the discrete dose. Rural equals 1 if the child's 2005 household locality had fewer than 2,500 inhabitants (INEGI 2024). Children are aged 15 or younger for HAZ and BMIZ and 10 or younger for WAZ in both survey waves, with a 2005 municipality. They were not measured in MxFLS-3 before November 2009, for the same reason as in Table **??**. Regressions are on the 2005 to 2009 change in the WHO z-score, without children in wet municipalities (z of 1 or above). The omitted group is near-normal rainfall ($-1 < z < 1$). There are no controls or fixed effects. Dry × rural is the difference in the effect of dry rainfall for rural children. Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are in brackets, which tend to be conservative with few treated municipalities (MacKinnon, Nielsen, and Webb 2023). Stars use the clustered p-values, and where a wild bootstrap p-value is shown both must pass. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 43—Dose Groups and Baseline Characteristics, HAZ Sample

	(1) Electricity	(2) Good floor	(3) Good roof	(4) Enough ventilation	(5) Rooms per person	(6) Log expenditure	(7) Expenditure (pesos)	(8) Rural	(9) PROGRESA	(10) Remittances	(11) Male	(12) Age	(13) Fully vaccinated	(14) Mother's height (cm)	(15) Mother's schooling	(16) Mother's age at birth
Moderately dry	-0.049*	-0.032	-0.046	0.094*	-0.024	-0.072	287.797	0.042	0.094	-0.087*	0.015	-0.157	0.037	0.550	-0.602	0.848
	(0.027)	(0.071)	(0.060)	(0.052)	(0.027)	(0.164)	(605.606)	(0.164)	(0.094)	(0.047)	(0.025)	(0.156)	(0.051)	(1.047)	(0.644)	(0.515)
Severely dry or worse	-0.010	0.104**	0.060	0.067	0.070**	0.207	433.784	-0.255*	-0.117*	-0.089**	-0.014	0.171	0.080	0.609	1.187**	0.082
	(0.010)	(0.043)	(0.077)	(0.069)	(0.033)	(0.162)	(583.739)	(0.133)	(0.060)	(0.044)	(0.029)	(0.240)	(0.053)	(1.601)	(0.513)	(0.563)
Observations	1,959	1,959	1,959	1,944	1,956	1,929	1,929	1,959	1,912	1,773	2,351	2,351	2,231	2,219	2,247	2,269
Mean of dep. var.	0.981	0.827	0.808	0.568	0.384	6.513	1292.315	0.464	0.228	0.368	0.496	5.885	0.480	154.375	7.197	26.407
Fixed effects									None							

	(17) Mother employed	(18) Father's height (cm)	(19) Father's schooling	(20) Father's age at birth	(21) Father employed	(22) Birth order	(23) Oldest child	(24) Birth weight (kg)	(25) Born at home	(26) Weeks breastfed	(27) Iron in pregnancy	(28) Premature	(29) HAZ	(30) Long-run rain mean	(31) Long-run rain SD
Moderately dry	0.029	-0.509	-0.542	1.246**	0.009	0.025	-0.023	-0.010	0.081	-1.666	-0.036	0.021	-0.014	121.226	0.615
	(0.040)	(1.027)	(0.632)	(0.595)	(0.016)	(0.255)	(0.039)	(0.050)	(0.083)	(3.645)	(0.073)	(0.022)	(0.140)	(133.660)	(22.412)
Severely dry or worse	0.073	0.360	0.880	-0.074	0.022	-0.352*	0.039	0.001	-0.104***	-11.203***	0.124**	0.008	0.216	198.196**	2.694
	(0.058)	(1.438)	(0.634)	(0.796)	(0.014)	(0.206)	(0.034)	(0.047)	(0.027)	(2.937)	(0.050)	(0.030)	(0.154)	(90.460)	(17.307)
Observations	2,247	1,561	1,863	1,929	1,867	1,675	1,675	1,529	1,859	1,644	1,656	1,652	2,351	2,351	2,351
Mean of dep. var.	0.288	166.465	7.384	29.896	0.958	2.548	0.360	3.256	0.111	33.320	0.743	0.138	-0.580	723.775	123.672
Fixed effects								None							

Notes: The sample is the 2,351 children in the HAZ discrete-dose regressions. Each column regresses a baseline characteristic on indicators for moderately dry ($-1.5 < z \leq -1$) and severely dry or worse ($z \leq -1.5$) municipalities. Each coefficient is the mean difference in the characteristic between children in that dry dose group and children in the near-normal dose group ($-1 < z < 1$). Columns 1–10 are from MxFLS-1 (2002) and columns 11–29 from MxFLS-2 (2005). Of particular interest are the measures of household poverty. Expenditure per person is a proxy for income and is shown in logs and in 2002 pesos per month. PROGRESA equals 1 if any household member was in the program in the last 12 months, which marks the household as poor by the program's targeting. The dwelling characteristics also reflect household poverty. Long-run rainfall is the municipality's 1981–2004 May–October mean and SD in millimeters. The table has no fixed effects because the discrete-dose regressions use each child's 2005-to-2009 change without fixed effects or covariates. Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are not reported. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

Table 44—Dose Groups and Baseline Characteristics, WAZ Sample

	(1) Electricity	(2) Good floor	(3) Good roof	(4) Enough ventilation	(5) Rooms per person	(6) Log expenditure	(7) Expenditure (pesos)	(8) Rural	(9) PROGRESA	(10) Remittances	(11) Male	(12) Age	(13) Fully vaccinated	(14) Mother's height (cm)	(15) Mother's schooling	(16) Mother's age at birth
Moderately dry	-0.049*	-0.080	-0.056	0.056	-0.018	-0.141	-417.097	0.088	0.113	-0.034	0.041	-0.041	0.079	0.591	-0.446	0.548
	(0.029)	(0.072)	(0.048)	(0.058)	(0.035)	(0.187)	(412.753)	(0.165)	(0.086)	(0.063)	(0.047)	(0.140)	(0.052)	(1.115)	(0.652)	(0.798)
Severely dry or worse	-0.009	0.039	-0.000	-0.009	0.082*	0.123	-228.740	-0.228*	-0.070	0.011	0.047	-0.171	0.054	0.089	0.899*	0.813
	(0.015)	(0.057)	(0.105)	(0.060)	(0.048)	(0.177)	(432.013)	(0.137)	(0.053)	(0.058)	(0.061)	(0.138)	(0.040)	(1.579)	(0.503)	(0.753)
Observations	786	786	786	780	785	771	771	786	768	693	1,045	1,045	991	994	1,009	1,016
Mean of dep. var.	0.981	0.831	0.818	0.571	0.409	6.567	1279.152	0.443	0.185	0.413	0.513	2.442	0.361	154.773	7.568	26.027
Fixed effects									None							

	(17) Mother employed	(18) Father's height (cm)	(19) Father's schooling	(20) Father's age at birth	(21) Father employed	(22) Birth order	(23) Oldest child	(24) Birth weight (kg)	(25) Born at home	(26) Weeks breastfed	(27) Iron in pregnancy	(28) Premature	(29) WAZ	(30) Long-run rain mean	(31) Long-run rain SD
Moderately dry	0.021	-0.135	-0.435	1.035	0.011	0.252	-0.020	0.049	0.078	1.199	-0.031	0.021	0.179	113.864	-1.609
	(0.037)	(1.270)	(0.596)	(0.789)	(0.020)	(0.265)	(0.050)	(0.067)	(0.066)	(3.729)	(0.059)	(0.033)	(0.168)	(129.920)	(21.534)
Severely dry or worse	0.033	0.557	0.667	0.474	0.027	-0.226	0.016	-0.004	-0.048**	-1.792	0.108**	0.009	0.141	215.088**	2.542
	(0.064)	(1.510)	(0.589)	(0.981)	(0.020)	(0.183)	(0.051)	(0.067)	(0.020)	(3.195)	(0.046)	(0.052)	(0.116)	(99.713)	(16.946)
Observations	1,008	688	839	860	839	599	599	744	794	767	776	773	1,045	1,045	1,045
Mean of dep. var.	0.240	166.762	7.690	29.558	0.956	2.072	0.487	3.247	0.073	32.083	0.791	0.177	-0.298	715.435	122.355
Fixed effects								None							

Notes: The sample is the 1,045 children in the WAZ discrete-dose regressions. Each column regresses a baseline characteristic on indicators for moderately dry ($-1.5 < z \leq -1$) and severely dry or worse ($z \leq -1.5$) municipalities. Each coefficient is the mean difference in the characteristic between children in that dry dose group and children in the near-normal dose group ($-1 < z < 1$). Columns 1–10 are from MxFLS-1 (2002) and columns 11–29 from MxFLS-2 (2005). Of particular interest are the measures of household poverty. Expenditure per person is a proxy for income and is shown in logs and in 2002 pesos per month. PROGRESA equals 1 if any household member was in the program in the last 12 months, which marks the household as poor by the program's targeting. The dwelling characteristics also reflect household poverty. Long-run rainfall is the municipality's 1981–2004 May–October mean and SD in millimeters. The table has no fixed effects because the discrete-dose regressions use each child's 2005-to-2009 change without fixed effects or covariates. Standard errors clustered by 2005 municipality are in parentheses. Wild cluster bootstrap p-values are not reported. $^{*}p < .10$, $^{**}p < .05$, $^{***}p < .01$.

FIGURE 16—CHANGE IN BMI-FOR-AGE BY 2009 RAINFALL DEFICIT, CONTINUOUS DOSE




FIGURE 17—CHANGE IN HEIGHT-FOR-AGE BY 2009 RAINFALL DEFICIT, CONTINUOUS DOSE




Figure 18—Change in Weight-for-Age by 2009 Rainfall Deficit, Continuous Dose




Figure 19—ATT by 2009 Rainfall Deficit, BMI-for-Age, Continuous Dose




Figure 20—ATT by 2009 Rainfall Deficit, Height-for-Age, Continuous Dose




Figure 21—ATT by 2009 Rainfall Deficit, Weight-for-Age, Continuous Dose

