Ethics of Data Science – Part II

Randomised Controlled Trials

How many of these apply to non-clinical contexts?

- 1. Randomisation is one of many virtues the RCT has. Others include:
- 2. Pre-registering the hypotheses to be tested in a Statistical Analysis Plan
- 3. Formal reasoning on whether the trial has a large enough sample size
- 4. Obtaining informed consent from the subjects involved
- 5. An obligation to report negative results
- The concept of "double blinding"
- 7. Inability to casually change course midway through based on partial results
- 8. Prospective planning of what to measure as evidence of success ("endpoint")

Ethics of Data Science

3. Statistical Analysis Plan

5. Obligation to publish negative results

Pre-registering your intended analysis and testable hypotheses protects against p-hacking and related offences.



All trials registered. All results reported.

September 2013

The AllTrials campaign calls for all past and present clinical trials to be registered and their results reported.

4. Informed consent



THIS ARTICLE HAS BEEN CORRECTED +

In an experiment with people who use Facebook, we test whether emotional contagion occurs outside of in-person interaction between individuals by reducing the amount of emotional content in the News Feed. When positive expressions were reduced, people produced fewer positive posts and more negative posts; when negative expressions were reduced, the opposite pattern occurred. These results indicate that emotions expressed by others on Facebook influence our own emotions, constituting experimental evidence for massive-scale contagion via social networks.

Ethics of Data Science

4. Informed consent





Ethics of Data Science

6. Double blinding

- The subject does not know which treatment they are taking ——— Placebo effect
- The clinical staff performing the trial does not know which treatment they are giving to patients
- The principal investigator does not know which patients are ——— Confirmation bias taking which treatment

Different level of care depending on treatment

Placebo effect outside medicine: "You have been randomly selected to test a new social media feed algorithm. You will be asked to report on your level of satisfaction with the new algorithm in 3 months."

Clinical staff bias outside medicine: sales reps excited about using a novel piece of software end up more motivated/enthusiastic during client calls and hence more effective, but not because of the tool itself.

Principal Investigator bias can be counteracted by best practices around pre-registration.